

## $B^0$ – $\overline{B}^0$ MIXING

Updated April 2008 by O. Schneider (Ecole Polytechnique Fédérale de Lausanne).

There are two neutral  $B^0$ – $\overline{B}^0$  meson systems,  $B_d^0$ – $\overline{B}_d^0$  and  $B_s^0$ – $\overline{B}_s^0$  (generically denoted  $B_q^0$ – $\overline{B}_q^0$ ,  $q = s, d$ ), which exhibit particle-antiparticle mixing [1]. This mixing phenomenon is described in Ref. 2. In the following, we adopt the notation introduced in Ref. 2, and assume  $CPT$  conservation throughout. In each system, the light (L) and heavy (H) mass eigenstates,

$$|B_{L,H}\rangle = p|B_q^0\rangle \pm q|\overline{B}_q^0\rangle, \quad (1)$$

have a mass difference  $\Delta m_q = m_H - m_L > 0$ , and a total decay width difference  $\Delta\Gamma_q = \Gamma_L - \Gamma_H$ . In the absence of  $CP$  violation in the mixing,  $|q/p| = 1$ , these differences are given by  $\Delta m_q = 2|M_{12}|$  and  $|\Delta\Gamma_q| = 2|\Gamma_{12}|$ , where  $M_{12}$  and  $\Gamma_{12}$  are the off-diagonal elements of the mass and decay matrices [2]. The evolution of a pure  $|B_q^0\rangle$  or  $|\overline{B}_q^0\rangle$  state at  $t = 0$  is given by

$$|B_q^0(t)\rangle = g_+(t)|B_q^0\rangle + \frac{q}{p}g_-(t)|\overline{B}_q^0\rangle, \quad (2)$$

$$|\overline{B}_q^0(t)\rangle = g_+(t)|\overline{B}_q^0\rangle + \frac{p}{q}g_-(t)|B_q^0\rangle, \quad (3)$$

which means that the flavor states remain unchanged (+) or oscillate into each other (–) with time-dependent probabilities proportional to


$$|g_{\pm}(t)|^2 = \frac{e^{-\Gamma_q t}}{2} \left[ \cosh\left(\frac{\Delta\Gamma_q}{2}t\right) \pm \cos(\Delta m_q t) \right], \quad (4)$$

where  $\Gamma_q = (\Gamma_H + \Gamma_L)/2$ . In the absence of  $CP$  violation, the time-integrated mixing probability  $\int |g_-(t)|^2 dt / (\int |g_-(t)|^2 dt + \int |g_+(t)|^2 dt)$  is given by

$$\chi_q = \frac{x_q^2 + y_q^2}{2(x_q^2 + 1)}, \quad \text{where} \quad x_q = \frac{\Delta m_q}{\Gamma_q}, \quad y_q = \frac{\Delta\Gamma_q}{2\Gamma_q}. \quad (5)$$

### ***Standard Model predictions and phenomenology***

In the Standard Model, the transitions  $B_q^0 \rightarrow \overline{B}_q^0$  and  $\overline{B}_q^0 \rightarrow B_q^0$  are due to the weak interaction. They are described, at the lowest order, by box diagrams involving two  $W$  bosons and two



**Figure 1:** Dominant box diagrams for the  $B_q^0 \rightarrow \bar{B}_q^0$  transitions ( $q = d$  or  $s$ ). Similar diagrams exist where one or both  $t$  quarks are replaced with  $c$  or  $u$  quarks.

up-type quarks (see Fig. 1), as is the case for  $K^0-\bar{K}^0$  mixing. However, the long range interactions arising from intermediate virtual states are negligible for the neutral  $B$  meson systems, because the large  $B$  mass is off the region of hadronic resonances. The calculation of the dispersive and absorptive parts of the box diagrams yields the following predictions for the off-diagonal element of the mass and decay matrices [3],

$$M_{12} = -\frac{G_F^2 m_W^2 \eta_B m_{B_q} B_{B_q} f_{B_q}^2}{12\pi^2} S_0(m_t^2/m_W^2) (V_{tq}^* V_{tb})^2, \quad (6)$$

$$\Gamma_{12} = \frac{G_F^2 m_b^2 \eta'_B m_{B_q} B_{B_q} f_{B_q}^2}{8\pi} \times \left[ (V_{tq}^* V_{tb})^2 + V_{tq}^* V_{tb} V_{cq}^* V_{cb} \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right) + (V_{cq}^* V_{cb})^2 \mathcal{O}\left(\frac{m_c^4}{m_b^4}\right) \right], \quad (7)$$

where  $G_F$  is the Fermi constant,  $m_W$  the  $W$  boson mass, and  $m_i$  the mass of quark  $i$ ;  $m_{B_q}$ ,  $f_{B_q}$  and  $B_{B_q}$  are the  $B_q^0$  mass, weak decay constant and bag parameter, respectively. The known function  $S_0(x_t)$  can be approximated very well by  $0.784 x_t^{0.76}$  [4], and  $V_{ij}$  are the elements of the CKM matrix [5]. The QCD corrections  $\eta_B$  and  $\eta'_B$  are of order unity. The only non-negligible contributions to  $M_{12}$  are from box diagrams involving two top quarks. The phases of  $M_{12}$  and  $\Gamma_{12}$  satisfy

$$\phi_M - \phi_\Gamma = \pi + \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right), \quad (8)$$

implying that the mass eigenstates have mass and width differences of opposite signs. This means that, like in the  $K^0-\bar{K}^0$  system, the heavy state is expected to have a smaller decay width

than that of the light state:  $\Gamma_H < \Gamma_L$ . Hence,  $\Delta\Gamma = \Gamma_L - \Gamma_H$  is expected to be positive in the Standard Model.

Furthermore, the quantity

$$\left| \frac{\Gamma_{12}}{M_{12}} \right| \simeq \frac{3\pi}{2} \frac{m_b^2}{m_W^2} \frac{1}{S_0(m_t^2/m_W^2)} \sim \mathcal{O}\left(\frac{m_b^2}{m_t^2}\right) \quad (9)$$

is small, and a power expansion of  $|q/p|^2$  yields

$$\left| \frac{q}{p} \right|^2 = 1 + \left| \frac{\Gamma_{12}}{M_{12}} \right| \sin(\phi_M - \phi_\Gamma) + \mathcal{O}\left(\left| \frac{\Gamma_{12}}{M_{12}} \right|^2\right). \quad (10)$$

Therefore, considering both Eqs. (8) and (9), the  $CP$ -violating parameter

$$1 - \left| \frac{q}{p} \right|^2 \simeq \text{Im}\left(\frac{\Gamma_{12}}{M_{12}}\right) \quad (11)$$

is expected to be very small:  $\sim \mathcal{O}(10^{-3})$  for the  $B_d^0\text{--}\bar{B}_d^0$  system and  $\lesssim \mathcal{O}(10^{-4})$  for the  $B_s^0\text{--}\bar{B}_s^0$  system [6].

In the approximation of negligible  $CP$  violation in mixing, the ratio  $\Delta\Gamma_q/\Delta m_q$  is equal to the small quantity  $|\Gamma_{12}/M_{12}|$  of Eq. (9); it is hence independent of CKM matrix elements, *i.e.*, the same for the  $B_d^0\text{--}\bar{B}_d^0$  and  $B_s^0\text{--}\bar{B}_s^0$  systems. Recent calculations [7] yield  $\sim 5 \times 10^{-3}$  with a  $\sim 20\%$  uncertainty. Given the current experimental knowledge on the mixing parameter  $x_q$

$$\begin{cases} x_d = 0.776 \pm 0.008 & (B_d^0\text{--}\bar{B}_d^0 \text{ system}) \\ x_s = 26.1 \pm 0.5 & (B_s^0\text{--}\bar{B}_s^0 \text{ system}) \end{cases}, \quad (12)$$

the Standard Model thus predicts that  $\Delta\Gamma_d/\Gamma_d$  is very small (below 1%), but  $\Delta\Gamma_s/\Gamma_s$  considerably larger ( $\sim 10\%$ ). These width differences are caused by the existence of final states to which both the  $B_q^0$  and  $\bar{B}_q^0$  mesons can decay. Such decays involve  $b \rightarrow c\bar{c}q$  quark-level transitions, which are Cabibbo-suppressed if  $q = d$  and Cabibbo-allowed if  $q = s$ .

A recent and complete set of Standard Model predictions for all mixing parameters in both the  $B_d^0\text{--}\bar{B}_d^0$  and  $B_s^0\text{--}\bar{B}_s^0$  systems can be found in Ref. 7.

### ***Experimental issues and methods for oscillation analyses***

Time-integrated measurements of  $B^0\text{--}\bar{B}^0$  mixing were published for the first time in 1987 by UA1 [8] and ARGUS [9], and

since then by many other experiments. These measurements are typically based on counting same-sign and opposite-sign lepton pairs from the semileptonic decay of the produced  $b\bar{b}$  pairs. Such analyses cannot easily separate the contributions from the different  $b$ -hadron species, therefore, the clean environment of  $\Upsilon(4S)$  machines (where only  $B_d^0$  and charged  $B_u$  mesons are produced) is in principle best suited to measure  $\chi_d$ .

However, better sensitivity is obtained from time-dependent analyses aiming at the direct measurement of the oscillation frequencies  $\Delta m_d$  and  $\Delta m_s$ , from the proper time distributions of  $B_d^0$  or  $B_s^0$  candidates identified through their decay in (mostly) flavor-specific modes, and suitably tagged as mixed or unmixed. This is particularly true for the  $B_s^0\text{--}\bar{B}_s^0$  system, where the large value of  $x_s$  implies maximal mixing, *i.e.*,  $\chi_s \simeq 1/2$ . In such analyses, the  $B_d^0$  or  $B_s^0$  mesons are either fully reconstructed, partially reconstructed from a charm meson, selected from a lepton with the characteristics of a  $b \rightarrow \ell^-$  decay, or selected from a reconstructed displaced vertex. At high-energy colliders (LEP, SLC, Tevatron), the proper time  $t = \frac{m_B}{p}L$  is measured from the distance  $L$  between the production vertex and the  $B$  decay vertex, and from an estimate of the  $B$  momentum  $p$ . At asymmetric  $B$  factories (KEKB, PEP-II), producing  $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B_d^0\bar{B}_d^0$  events with a boost  $\beta\gamma$  ( $= 0.425, 0.55$ ), the proper time difference between the two  $B$  candidates is estimated as  $\Delta t \simeq \frac{\Delta z}{\beta\gamma c}$ , where  $\Delta z$  is the spatial separation between the two  $B$  decay vertices along the boost direction. In all cases, the good resolution needed on the vertex positions is obtained with silicon detectors.

The average statistical significance  $\mathcal{S}$  of a  $B_d^0$  or  $B_s^0$  oscillation signal can be approximated as [10]

$$\mathcal{S} \approx \sqrt{N/2} f_{\text{sig}} (1 - 2\eta) e^{-(\Delta m \sigma_t)^2/2}, \quad (13)$$

where  $N$  is the number of selected and tagged candidates,  $f_{\text{sig}}$  is the fraction of signal in that sample,  $\eta$  is the total mistag probability, and  $\sigma_t$  is the resolution on proper time (or proper time difference). The quantity  $\mathcal{S}$  decreases very quickly as  $\Delta m$  increases; this dependence is controlled by  $\sigma_t$ , which is therefore

a critical parameter for  $\Delta m_s$  analyses. At high-energy colliders, the proper time resolution  $\sigma_t \sim \frac{m_B}{\langle p \rangle} \sigma_L \oplus t \frac{\sigma_p}{p}$  includes a constant contribution due to the decay length resolution  $\sigma_L$  (typically 0.05–0.3 ps), and a term due to the relative momentum resolution  $\sigma_p/p$  (typically 10–20% for partially reconstructed decays), which increases with proper time. At  $B$  factories, the boost of the  $B$  mesons is estimated from the known beam energies, and the term due to the spatial resolution dominates (typically 1–1.5 ps because of the much smaller  $B$  boost).

In order to tag a  $B$  candidate as mixed or unmixed, it is necessary to determine its flavor both in the initial state and in the final state. The initial and final state mistag probabilities,  $\eta_i$  and  $\eta_f$ , degrade  $\mathcal{S}$  by a total factor  $(1 - 2\eta) = (1 - 2\eta_i)(1 - 2\eta_f)$ . In lepton-based analyses, the final state is tagged by the charge of the lepton from  $b \rightarrow \ell^-$  decays; the largest contribution to  $\eta_f$  is then due to  $\bar{b} \rightarrow \bar{c} \rightarrow \ell^-$  decays. Alternatively, the charge of a reconstructed charm meson ( $D^{*-}$  from  $B_d^0$  or  $D_s^-$  from  $B_s^0$ ), or that of a kaon hypothesized to come from a  $b \rightarrow c \rightarrow s$  decay [11], can be used. For fully-inclusive analyses based on topological vertexing, final-state tagging techniques include jet-charge [12] and charge-dipole [13,14] methods. At high-energy colliders, the methods to tag the initial state (*i.e.*, the state at production), can be divided into two groups: the ones that tag the initial charge of the  $\bar{b}$  quark contained in the  $B$  candidate itself (same-side tag), and the ones that tag the initial charge of the other  $b$  quark produced in the event (opposite-side tag). On the same side, the charge of a track from the primary vertex is correlated with the production state of the  $B$  if that track is a decay product of a  $B^{**}$  state or the first particle in the fragmentation chain [15,16]. Jet- and vertex-charge techniques work on both sides and on the opposite side, respectively. Finally, the charge of a lepton from  $b \rightarrow \ell^-$  or of a kaon from  $b \rightarrow c \rightarrow s$  can be used as opposite side tags, keeping in mind that their performance is degraded due to integrated mixing. At SLC, the beam polarization produced a sizeable forward-backward asymmetry in the  $Z \rightarrow b\bar{b}$  decays, and provided another very interesting and effective initial state

tag based on the polar angle of the  $B$  candidate [13]. Initial state tags have also been combined to reach  $\eta_i \sim 26\%$  at LEP [16,17], or even 22% at SLD [13] with full efficiency. In the case  $\eta_f = 0$ , this corresponds to an effective tagging efficiency  $Q = \epsilon D^2 = \epsilon(1 - 2\eta)^2$ , where  $\epsilon$  is the tagging efficiency, in the range 23 – 31%. The equivalent figure achieved by CDF during Tevatron Run I was  $\sim 3.5\%$  [18], reflecting the fact that tagging is more difficult at hadron colliders. The current CDF and DØ analyses of Tevatron Run II data reach  $\epsilon D^2 = (1.8 \pm 0.1)\%$  [19] and  $(2.5 \pm 0.2)\%$  [20] for opposite-side tagging, while same-side kaon tagging (for  $B_s^0$  oscillation analyses) is contributing an additional 3.7 – 4.8% at CDF [19], and pushes the combined performance to  $(4.5 \pm 0.9)\%$  at DØ [21].

At  $B$  factories, the flavor of a  $B_d^0$  meson at production cannot be determined, since the two neutral  $B$  mesons produced in a  $\Upsilon(4S)$  decay evolve in a coherent  $P$ -wave state where they keep opposite flavors at any time. However, as soon as one of them decays, the other follows a time-evolution given by Eqs. (2) or (3), where  $t$  is replaced with  $\Delta t$  (which will take negative values half of the time). Hence, the “initial state” tag of a  $B$  can be taken as the final-state tag of the other  $B$ . Effective tagging efficiencies  $Q$  of 30% are achieved by BABAR and Belle [22], using different techniques including  $b \rightarrow \ell^-$  and  $b \rightarrow c \rightarrow s$  tags. It is worth noting that, in this case, mixing of the other  $B$  (*i.e.*, the coherent mixing occurring before the first  $B$  decay) does not contribute to the mistag probability.

In the absence of experimental observation of a decay-width difference, oscillation analyses typically neglect  $\Delta\Gamma$  in Eq. (4), and describe the data with the physics functions  $\Gamma e^{-\Gamma t}(1 \pm \cos(\Delta m t))/2$  (high-energy colliders) or  $\Gamma e^{-\Gamma|\Delta t|}(1 \pm \cos(\Delta m \Delta t))/4$  (asymmetric  $\Upsilon(4S)$  machines). As can be seen from Eq. (4), a non-zero value of  $\Delta\Gamma$  would effectively reduce the oscillation amplitude with a small time-dependent factor that would be very difficult to distinguish from time resolution effects. Measurements of  $\Delta m_d$  are usually extracted from the data using a maximum likelihood fit. To extract information useful for the interpretation of  $B_s^0$  oscillation searches and for the combination of their results, a method [10] is followed in

which a  $B_s^0$  oscillation amplitude  $\mathcal{A}$  is measured as a function of a fixed test value of  $\Delta m_s$ , using a maximum likelihood fit based on the functions  $\Gamma_s e^{-\Gamma_s t} (1 \pm \mathcal{A} \cos(\Delta m_s t))/2$ . To a good approximation, the statistical uncertainty on  $\mathcal{A}$  is Gaussian and equal to  $1/\mathcal{S}$  from Eq. (13). If  $\Delta m_s$  is equal to its true value, one expects  $\mathcal{A} = 1$  within the total uncertainty  $\sigma_{\mathcal{A}}$ ; in case a signal is seen, its observed (or expected) significance will be defined as  $\mathcal{A}/\sigma_{\mathcal{A}}$  (or  $1/\sigma_{\mathcal{A}}$ ). However, if  $\Delta m_s$  is (far) below its true value, a measurement consistent with  $\mathcal{A} = 0$  is expected. A value of  $\Delta m_s$  can be excluded at 95% CL if  $\mathcal{A} + 1.645 \sigma_{\mathcal{A}} \leq 1$  (since the integral of a normal distribution from  $-\infty$  to 1.645 is equal to 0.95). Because of the proper time resolution, the quantity  $\sigma_{\mathcal{A}}(\Delta m_s)$  is a steadily increasing function of  $\Delta m_s$ . We define the sensitivity for 95% CL exclusion of  $\Delta m_s$  values (or for a  $3\sigma$  or  $5\sigma$  observation of  $B_s^0$  oscillations) as the value of  $\Delta m_s$  for which  $1/\sigma_{\mathcal{A}} = 1.645$  (or  $1/\sigma_{\mathcal{A}} = 3$  or  $5$ ).

### **$B_d^0$ mixing studies**

Many  $B_d^0\text{-}\bar{B}_d^0$  oscillations analyses have been published [23] by the ALEPH [24], BABAR [25], Belle [26], CDF [15], DØ [20], DELPHI [14,27], L3 [28], and OPAL [29,30] collaborations. Although a variety of different techniques have been used, the individual  $\Delta m_d$  results obtained at high-energy colliders have remarkably similar precision. Their average is compatible with the recent and more precise measurements from asymmetric  $B$  factories. The systematic uncertainties are not negligible; they are often dominated by sample composition, mistag probability, or  $b$ -hadron lifetime contributions. Before being combined, the measurements are adjusted on the basis of a common set of input values, including the  $b$ -hadron lifetimes and fractions published in this *Review*. Some measurements are statistically correlated. Systematic correlations arise both from common physics sources (fragmentation fractions, lifetimes, branching ratios of  $b$  hadrons), and from purely experimental or algorithmic effects (efficiency, resolution, tagging, background description). Combining all published measurements [14,15,20,24–30] and accounting for all identified correlations yields  $\Delta m_d = 0.507 \pm 0.003(\text{stat}) \pm 0.003(\text{syst}) \text{ ps}^{-1}$  [31], a result dominated by the  $B$  factories.

On the other hand, ARGUS and CLEO have published time-integrated measurements [32–34], which average to  $\chi_d = 0.182 \pm 0.015$ . Following Ref. 34, the width difference  $\Delta\Gamma_d$  could in principle be extracted from the measured value of  $\Gamma_d$  and the above averages for  $\Delta m_d$  and  $\chi_d$  (see Eq. (5)), provided that  $\Delta\Gamma_d$  has a negligible impact on the  $\Delta m_d$  measurements. However, direct time-dependent studies published by DELPHI [14] and BABAR [35] provide stronger constraints, which can be combined to yield  $\text{sign}(\text{Re}\lambda_{CP})\Delta\Gamma_d/\Gamma_d = 0.009 \pm 0.037$  [31].

Assuming  $\Delta\Gamma_d = 0$  and no  $CP$  violation in mixing, and using the measured  $B_d^0$  lifetime of  $1.530 \pm 0.009$  ps, the  $\Delta m_d$  and  $\chi_d$  results are combined to yield the world average

$$\Delta m_d = 0.507 \pm 0.005 \text{ ps}^{-1} \quad (14)$$

or, equivalently,

$$\chi_d = 0.1878 \pm 0.0024. \quad (15)$$

Evidence for  $CP$  violation in  $B_d^0$  mixing has been searched for, both with flavor-specific and inclusive  $B_d^0$  decays, in samples where the initial flavor state is tagged. In the case of semileptonic (or other flavor-specific) decays, where the final-state tag is also available, the following asymmetry [2]

$$\mathcal{A}_{\text{SL}}^d = \frac{N(\overline{B}_d^0(t) \rightarrow \ell^+ \nu_\ell X) - N(B_d^0(t) \rightarrow \ell^- \bar{\nu}_\ell X)}{N(\overline{B}_d^0(t) \rightarrow \ell^+ \nu_\ell X) + N(B_d^0(t) \rightarrow \ell^- \bar{\nu}_\ell X)} \simeq 1 - |q/p|_d^2 \quad (16)$$

has been measured, either in time-integrated analyses at CLEO [34,36], CDF [37,38] and DØ [39], or in time-dependent analyses at LEP [30,40,41], BABAR [35,42,43] and Belle [44]. In the inclusive case, also investigated at LEP [40,41,45], no final-state tag is used, and the asymmetry [46]

$$\begin{aligned} & \frac{N(B_d^0(t) \rightarrow \text{all}) - N(\overline{B}_d^0(t) \rightarrow \text{all})}{N(B_d^0(t) \rightarrow \text{all}) + N(\overline{B}_d^0(t) \rightarrow \text{all})} \\ & \simeq \mathcal{A}_{\text{SL}}^d \left[ \frac{x_d}{2} \sin(\Delta m_d t) - \sin^2 \left( \frac{\Delta m_d t}{2} \right) \right] \end{aligned} \quad (17)$$

must be measured as a function of the proper time to extract information on  $CP$  violation. In all cases, asymmetries compatible with zero have been found, with a precision limited by the available statistics. A simple average of all



published results for the  $B_d^0$  meson [30,34–36,39,41,42,44,45] yields  $\mathcal{A}_{\text{SL}}^d = -0.0049 \pm 0.0038$ , under the assumption of no  $CP$  violation in  $B_s^0$  mixing. Published results at  $B$  factories only [34–36,42,44], where no  $B_s^0$  is produced, average to


$$\mathcal{A}_{\text{SL}}^d = -0.0005 \pm 0.0056, \text{ or } |q/p|_d = 1.0002 \pm 0.0028, \quad (18)$$

a result which does not yet constrain the Standard Model.

The  $\Delta m_d$  result of Eq. (14) provides an estimate of  $2|M_{12}|$ , and can be used, together with Eq. (6), to extract the magnitude of the CKM matrix element  $V_{td}$  within the Standard Model [47]. The main experimental uncertainties on the resulting estimate of  $|V_{td}|$  come from  $m_t$  and  $\Delta m_d$ ; however, the extraction is at present completely dominated by the uncertainty on the hadronic matrix element  $f_{B_d}\sqrt{B_{B_d}} = 244 \pm 26$  MeV obtained from lattice QCD calculations [48].

### **$B_s^0$ mixing studies**

In the decade before the Tevatron Run II results became available,  $B_s^0\text{--}\bar{B}_s^0$  oscillations have been the subject of many studies from ALEPH [49], CDF [50], DELPHI [14,17,51], OPAL [52] and SLD [13,53,54]. The most sensitive analyses appeared to be the ones based on inclusive lepton samples. Because of their better proper time resolution, the small data samples analyzed inclusively at SLD, as well as the fully reconstructed  $B_s$  decays at LEP were also very useful to explore the high  $\Delta m_s$  region. However, all results were limited by the available statistics. All published measurements of the  $B_s^0$  oscillation amplitude [13,14,17,49–53] are averaged [31] to yield the combined amplitudes  $\mathcal{A}$  shown in Fig. 2 (bottom) as a function of  $\Delta m_s$ . The individual results have been adjusted to common physics inputs, and all known correlations have been accounted for; the sensitivities of the inclusive analyses, which depend directly through Eq. (13) on the assumed fraction  $f_s$  of  $B_s^0$  mesons in an unbiased sample of weakly-decaying  $b$  hadrons, have also been rescaled to a common average of  $f_s = 0.105 \pm 0.009$ . The combined sensitivity for 95% CL exclusion of  $\Delta m_s$  values is found to be  $18.3 \text{ ps}^{-1}$ . All values of  $\Delta m_s$  below  $14.6 \text{ ps}^{-1}$  are excluded at 95% CL, while the values between  $14.6$  and  $21.7 \text{ ps}^{-1}$  cannot be excluded, because the data is compatible



**Figure 2:** Combined measurements of the  $B_s^0$  oscillation amplitude as a function of  $\Delta m_s$ . Top: CDF result based on Run II data, published in 2006 [19]. Middle: Average of all preliminary  $D\bar{O}$  results available at the end of 2007 [21]. Bottom: Average of all other results (mainly from LEP and SLD) published between 1997 and 2004. All measurements are dominated by statistical uncertainties. Neighboring points are statistically correlated.

with a signal in this region. However, the largest deviation from  $\mathcal{A} = 0$  in this range is a  $1.9\sigma$  effect only, so no signal can be claimed.

Tevatron Run II results based on  $1 \text{ fb}^{-1}$  of data became available in 2006. After  $D\bar{O}$  [55] reported  $17 < \Delta m_s < 21 \text{ ps}^{-1}$  (90% CL) and a most probable value of  $19 \text{ ps}^{-1}$  with an observed (expected) significance of  $2.5\sigma$  ( $0.9\sigma$ ), CDF [19] published the first direct evidence of  $B_s^0$  oscillations shortly followed by a  $> 5\sigma$  observation (shown at the top of Fig. 2). The measured value of  $\Delta m_s$  is

$$\Delta m_s = 17.77 \pm 0.10(\text{stat}) \pm 0.07(\text{syst}) \text{ ps}^{-1}, \quad (19)$$

based on samples of flavour-tagged hadronic and semileptonic  $B_s^0$  decays, partially or fully reconstructed in flavour-specific final states. More recently, DØ [21] obtained with  $2.4 \text{ fb}^{-1}$  an independent  $2.9\sigma$  preliminary evidence for  $B_s^0$  oscillations (middle of Fig. 2) at  $\Delta m_s = 18.53 \pm 0.93(\text{stat}) \pm 0.30(\text{syst}) \text{ ps}^{-1}$  [56], consistent with the CDF measurement.

The information on  $|V_{ts}|$  obtained in the framework of the Standard Model is hampered by the hadronic uncertainty, as in the  $B_d^0$  case. However, several uncertainties cancel in the frequency ratio

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{B_s}}{m_{B_d}} \xi^2 \left| \frac{V_{ts}}{V_{td}} \right|^2, \quad (20)$$

where  $\xi = (f_{B_s} \sqrt{B_{B_s}})/(f_{B_d} \sqrt{B_{B_d}}) = 1.210^{+0.047}_{-0.035}$  is an SU(3) flavor-symmetry breaking factor obtained from lattice QCD calculations [48]. Using the measurements of Eqs. (14) and (19), one can extract

$$\left| \frac{V_{td}}{V_{ts}} \right| = 0.2060 \pm 0.0012(\text{exp})^{+0.0080}_{-0.0060}(\text{lattice}), \quad (21)$$

in good agreement with (but much more precise than) the recent results obtained by the Belle [57] and BABAR [58] collaborations based on the observation of the  $b \rightarrow d\gamma$  transition. The CKM matrix can be constrained using experimental results on observables such as  $\Delta m_d$ ,  $\Delta m_s$ ,  $|V_{ub}/V_{cb}|$ ,  $\epsilon_K$ , and  $\sin(2\beta)$  together with theoretical inputs and unitarity conditions [47,59,60]. The constraint from our knowledge on the ratio  $\Delta m_s/\Delta m_d$  is presently more effective in limiting the position of the apex of the CKM unitarity triangle than the one obtained from the  $\Delta m_d$  measurements alone, due to the reduced hadronic uncertainty in Eq. (20). We also note that the measured value of  $\Delta m_s$  is consistent with the Standard Model prediction obtained from CKM fits where no experimental information on  $\Delta m_s$  is used, *e.g.*,  $20.6 \pm 2.6 \text{ ps}^{-1}$  [59] or  $17.7^{+6.4}_{-2.1} \text{ ps}^{-1}$  [60].

Information on  $\Delta\Gamma_s$  can be obtained by studying the proper time distribution of untagged  $B_s^0$  samples [61]. In the case of an inclusive  $B_s^0$  selection [62], or a semileptonic (or flavour-specific)  $B_s^0$  decay selection [17,63,64], both the short- and

long-lived components are present, and the proper time distribution is a superposition of two exponentials with decay constants  $\Gamma_{L,H} = \Gamma_s \pm \Delta\Gamma_s/2$ . In principle, this provides sensitivity to both  $\Gamma_s$  and  $(\Delta\Gamma_s/\Gamma_s)^2$ . Ignoring  $\Delta\Gamma_s$  and fitting for a single exponential leads to an estimate of  $\Gamma_s$  with a relative bias proportional to  $(\Delta\Gamma_s/\Gamma_s)^2$ . An alternative approach, which is directly sensitive to first order in  $\Delta\Gamma_s/\Gamma_s$ , is to determine the lifetime of  $B_s^0$  candidates decaying to  $CP$  eigenstates; measurements exist for  $B_s^0 \rightarrow K^+K^-$  [65],  $B_s^0 \rightarrow J/\psi\phi$  [66,67], and  $B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}$  [68], which are mostly  $CP$ -even states [69]. However, in the case of  $B_s^0 \rightarrow J/\psi\phi$ , this technique has now been replaced by more sensitive time-dependent angular analyses that allow the simultaneous extraction of  $\Delta\Gamma_s/\Gamma_s$  and the  $CP$ -even and  $CP$ -odd amplitudes [70,71]. Estimates of  $\Delta\Gamma_s/\Gamma_s$  have also been obtained directly from measurements of the  $B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}$  branching ratio [68,72], under the assumption that these decays account for all the  $CP$ -even final states (however, no systematic uncertainty due to this assumption is given, so the averages quoted below will not include these estimates).

Applying the combination procedure of Ref. 31 (including the constraint from the flavour-specific lifetime measurements) on the published results [17,63,66,68,70,71] yields

$$\Delta\Gamma_s/\Gamma_s = +0.069_{-0.062}^{+0.058} \quad \text{and} \quad 1/\Gamma_s = 1.470_{-0.027}^{+0.026} \text{ ps}, \quad (22)$$

or equivalently

$$1/\Gamma_L = 1.419_{-0.038}^{+0.039} \text{ ps} \quad \text{and} \quad 1/\Gamma_H = 1.525_{-0.063}^{+0.062} \text{ ps}, \quad (23)$$

under the assumption of no  $CP$  violation in  $B_s^0$  mixing.

Recent studies also consider  $CP$  violation, either in untagged [70,71] or tagged [73,74]  $B_s^0 \rightarrow J/\psi\phi$  decays, and start to constrain the phase difference  $-2\beta_s$  between the  $B_s^0$  mixing diagram and the  $b \rightarrow c\bar{c}s$  tree decay diagram. In the Standard Model,  $\beta_s = \arg(-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*))$  is expected to be about one degree [7]. A proper combination of the current Tevatron constraints on  $\beta_s$  requires extra information not available in

the original publications and is being prepared in collaboration between CDF and DØ.

On the other hand  $CP$  violation in  $B_s^0$  mixing has been investigated through the asymmetry between positive and negative same-sign muon pairs from semi-leptonic decays of  $b\bar{b}$  pairs [37–39], and directly through the charge asymmetry of  $B_s^0 \rightarrow D_s \mu \nu X$  decays [75]. Combining all published results [37,39,75] with the knowledge of  $CP$  violation in  $B_d^0$  mixing from Eq. (18) leads to

$$\mathcal{A}_{\text{SL}}^s = -0.0030 \pm 0.0101, \text{ or } |q/p|_s = 1.0015 \pm 0.0051. \quad (24)$$

A large New Physics phase could possibly contribute to both  $CP$  violation in  $B_s^0 \rightarrow J/\psi\phi$ , and to the mixing phase difference of Eq. (8) on which  $\mathcal{A}_{\text{SL}}^s$  depends. Combined fits [76,77] of  $\beta_s$  and  $\mathcal{A}_{\text{SL}}^s$  measurements already yield interesting constraints on this New Physics phase. A deviation from the Standard Model, with a significance of more than  $3\sigma$ , has recently been claimed [77], based on a preliminary analysis including the latest Tevatron  $B_s^0$  mixing results [19,38,39,73–75].

***Average  $b$ -hadron mixing probability and  $b$ -hadron production fractions in  $Z$  decays and at high energy***

Mixing measurements can significantly improve our knowledge on the fractions  $f_u$ ,  $f_d$ ,  $f_s$ , and  $f_{\text{baryon}}$ , defined as the fractions of  $B_u$ ,  $B_d^0$ ,  $B_s^0$  and  $b$ -baryon in an unbiased sample of weakly decaying  $b$  hadrons produced in high-energy collisions. Indeed, time-integrated mixing analyses performed with lepton pairs from  $b\bar{b}$  events at high energy measure the quantity

$$\bar{\chi} = f'_d \chi_d + f'_s \chi_s, \quad (25)$$

where  $f'_d$  and  $f'_s$  are the fractions of  $B_d^0$  and  $B_s^0$  hadrons in a sample of semileptonic  $b$ -hadron decays. Assuming that all  $b$  hadrons have the same semileptonic decay width implies  $f'_q = f_q/(\Gamma_q \tau_b)$  ( $q = s, d$ ), where  $\tau_b$  is the average  $b$ -hadron lifetime. Hence  $\bar{\chi}$  measurements, together with the  $\chi_d$  average of Eq. (15) and the very good approximation  $\chi_s = 1/2$  (in fact  $\chi_s = 0.49927 \pm 0.00003$  from Eqs. (5), (19) and (22)), provide constraints on the fractions  $f_d$  and  $f_s$ .

The LEP experiments have measured  $f_s \times \text{BR}(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell X)$  [78],  $\text{BR}(b \rightarrow \Lambda_b^0) \times \text{BR}(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell X)$  [79], and  $\text{BR}(b \rightarrow \Xi_b^-) \times \text{BR}(\Xi_b^- \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X)$  [80] from partially reconstructed final states including a lepton,  $f_{\text{baryon}}$  from protons identified in  $b$  events [81], and the production rate of charged  $b$  hadrons [82]. The  $b$ -hadron fractions measured at CDF using double semileptonic  $K^* \mu \mu$  and  $\phi \mu \mu$  final states [83] and electron-charm final states [84] are at slight discrepancy with the ones measured at LEP. Furthermore the averages of the  $\bar{\chi}$  values measured at LEP,  $0.1259 \pm 0.0042$  [85], and at Tevatron,  $0.147 \pm 0.011$  [39,86], show a  $1.8\sigma$  deviation with respect to each other. This may be a hint that the fractions at the Tevatron might be different from the ones in  $Z$  decays. Combining [31] all the available information under the constraints  $f_u = f_d$  and  $f_u + f_d + f_s + f_{\text{baryon}} = 1$  yields the two set of averages shown in Table 1. The second set, obtained using both LEP and Tevatron results, has larger errors than the first set, obtained using LEP results only, because we have applied scale factors as advocated by the PDG for the treatment of marginally consistent data.

**Table 1:**  $\bar{\chi}$  and  $b$ -hadron fractions (see text).

	in $Z$ decays	at high energy
$\bar{\chi}$	$0.1259 \pm 0.0042$	$0.1284 \pm 0.0069$
$f_u = f_d$	$0.402 \pm 0.009$	$0.399 \pm 0.011$
$f_s$	$0.104 \pm 0.009$	$0.110 \pm 0.012$
$f_{\text{baryon}}$	$0.091 \pm 0.015$	$0.092 \pm 0.019$

### *Summary and prospects*

$B^0-\bar{B}^0$  mixing has been and still is a field of intense study. While fairly little experimental progress was recently achieved in the  $B_d^0$  sector, impressive new  $B_s^0$  results are becoming available from Run II of the Tevatron.  $B_s^0$  oscillations are now established and the mass difference in the  $B_s^0-\bar{B}_s^0$  system is measured very accurately, with a central value compatible with the Standard Model (SM) expectation and a relative precision

(0.7%) matching that in the  $B_d^0\text{--}\bar{B}_d^0$  system (0.9%). However, the extraction of  $|V_{td}/V_{ts}|$  from these measurements in the SM framework is limited by the hadronic uncertainty, which will be an important challenge to reduce in the future. New time-dependent angular analyses of  $B_s^0 \rightarrow J/\psi\phi$  decays and measurements of time-integrated  $B_s^0$  asymmetries at CDF and DØ are improving our knowledge of the other  $B_s^0$  mixing parameters: while  $CP$  violation in  $B_s^0\text{--}\bar{B}_s^0$  mixing is consistent with zero, with an uncertainty still large compared to the SM prediction, the relative decay width difference  $\Delta\Gamma_s/\Gamma_s$  is now determined to an absolute precision of  $\sim 6\%$ , smaller than the central value of the SM prediction. The data prefer  $\Gamma_L > \Gamma_H$  as predicted in the SM.

Improved  $B_s^0$  results are still to come, with very promising short-term prospects, both for  $\Delta\Gamma_s$  and  $CP$ -violating phases induced by mixing such as  $\beta_s$  and  $\arg(-M_{12}/\Gamma_{12})$ . Although first interesting experimental constraints have been published, very little is known yet about these phases, which are predicted to be very small in the SM. A full search for New Physics effects in these observables will require statistics beyond that of the Tevatron. These will eventually become available at CERN’s Large Hadron Collider, scheduled to start operation in 2008, where LHCb expects to be able to measure  $\beta_s$  down to the SM value after many years of operations [87].

$B$  mixing may still reveal a surprize, but much effort is needed for this, both on the experimental and theoretical sides, in particular to further reduce the hadronic uncertainties of lattice QCD calculations. In the long term, a stringent check of the consistency of the  $B_d^0$  and  $B_s^0$  mixing amplitudes (magnitudes and phases) with all other measured flavour-physics observables will be possible within the SM, leading to very tight limits (or otherwise new interesting knowledge!) on New Physics.

## References

1. T.D. Lee and C.S. Wu, *Ann. Rev. Nucl. Sci.* **16**, 511 (1966); I.I. Bigi and A.I. Sanda, *CP Violation*, Cambridge Univ. Press, 2000; G.C. Branco, L. Lavoura, and J.P. Silva, *CP Violation*, Clarendon Press Oxford, 1999.

2. See the review on “ $CP$  violation in meson decays,” by D. Kirkby and Y. Nir in this publication.
3. A.J. Buras, W. Slominski, and H. Steger, Nucl. Phys. **B245**, 369 (1984).
4. T. Inami and C.S. Lim, Prog. Theor. Phys. **65**, 297 (1981); for the power-like approximation, see A.J. Buras and R. Fleischer, page 91 in *Heavy Flavours II*, eds., A.J. Buras and M. Lindner, Singapore World Scientific, 1998.
5. M. Kobayashi and K. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
6. I.I. Bigi *et al.*, in *CP violation*, ed. C. Jarlskog, Singapore World Scientific, 1989.
7. A. Lenz and U. Nierste, JHEP **0607**, 072 (2007), arXiv: hep-ph/0612167v3.
8. C. Albajar *et al.* (UA1 Collab.), Phys. Lett. **B186**, 247 (1987).
9. H. Albrecht *et al.* (Argus Collab.), Phys. Lett. **B192**, 245 (1987).
10. H.-G. Moser and A. Roussarie, Nucl. Instrum. Methods **384**, 491 (1997).
11. SLD Collab., SLAC-PUB-7228, SLAC-PUB-7229, and SLAC-PUB-7230, *28th Int. Conf. on High Energy Physics*, Warsaw, 1996; J. Wittlin, PhD thesis, SLAC-R-582, 2001.
12. Aleph Collab., *Int. Europhysics Conf. on High Energy Physics*, Jerusalem, 1997.
13. K. Abe *et al.* (SLD Collab.), Phys. Rev. **D67**, 012006 (2003).
14. J. Abdallah *et al.* (DELPHI Collab.), Eur. Phys. J. **C28**, 155 (2003).
15. F. Abe *et al.* (CDF Collab.), Phys. Rev. Lett. **80**, 2057 (1998) and Phys. Rev. **D59**, 032001 (1999); Phys. Rev. **D60**, 051101 (1999); Phys. Rev. **D60**, 072003 (1999); T. Affolder *et al.* (CDF Collab.), Phys. Rev. **D60**, 112004 (1999).
16. R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C4**, 367 (1998); Eur. Phys. J. **C7**, 553 (1999).
17. P. Abreu *et al.* (DELPHI Collab.), Eur. Phys. J. **C16**, 555 (2000); Eur. Phys. J. **C18**, 229 (2000).
18. See tagging summary on page 160 of K. Anikeev *et al.*, “ $B$  physics at the Tevatron: Run II and beyond,” FERMILAB-PUB-01/97, hep-ph/0201071, and references therein.



19. A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **97**, 242003 (2006); this result supersedes A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **97**, 062003 (2006).
20. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. **D74**, 112002 (2006).
21. DØCollab., DØ note 5474-CONF, August 2007, and DØ note 5254-CONF, October 2006.
22. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **94**, 161803 (2005);  
K.-F. Chen *et al.* (Belle Collab.), Phys. Rev. **D72**, 012004 (2005).
23. Throughout this paper, we omit references of results that have been superseded by new published measurements.
24. D. Buskulic *et al.* (Aleph Collab.), Z. Phys. **C75**, 397 (1997).
25. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **88**, 221802 (2002) and Phys. Rev. **D66**, 032003 (2002); Phys. Rev. Lett. **88**, 221803 (2002); Phys. Rev. **D67**, 072002 (2003); Phys. Rev. **D73**, 012004 (2006).
26. N.C.Hastings *et al.* (Belle Collab.), Phys. Rev. **D67**, 052004 (2003); Y. Zheng *et al.* (Belle Collab.), Phys. Rev. **D67**, 092004 (2003); K. Abe *et al.* (Belle Collab.), Phys. Rev. **D71**, 072003 (2005).
27. P. Abreu *et al.* (DELPHI Collab.), Z. Phys. **C76**, 579 (1997).
28. M. Acciarri *et al.* (L3 Collab.), Eur. Phys. J. **C5**, 195 (1998).
29. G. Alexander *et al.* (OPAL Collab.), Z. Phys. **C72**, 377 (1996); K. Ackerstaff *et al.* (OPAL Collab.), Z. Phys. **C76**, 417 (1997); G. Abbiendi *et al.* (OPAL Collab.), Phys. Lett. **B493**, 266 (2000).
30. K. Ackerstaff *et al.* (OPAL Collab.), Z. Phys. **C76**, 401 (1997).
31. E. Barberio *et al.* (HFAG Collab.), “Averages of  $b$ -hadron properties at the end of 2006,” [arXiv:0704.3575v1](https://arxiv.org/abs/0704.3575v1) [[hep-ex](https://arxiv.org/abs/0704.3575v1)], April 2007; the combined results on  $b$ -hadron fractions, lifetimes and mixing parameters published in this *Review* have been obtained by the  $B$  oscillations working group of the Heavy Flavour Averaging Group (HFAG), using the methods and procedures described in Chapter 3 of the above paper, after updating the list of inputs; for more information, see <http://www.slac.stanford.edu/xorg/hfag/osc/>.

32. H. Albrecht *et al.* (ARGUS Collab.), *Z. Phys.* **C55**, 357 (1992); *Phys. Lett.* **B324**, 249 (1994).
33. J. Bartelt *et al.* (CLEO Collab.), *Phys. Rev. Lett.* **71**, 1680 (1993).
34. B.H. Behrens *et al.* (CLEO Collab.), *Phys. Lett.* **B490**, 36 (2000).
35. B. Aubert *et al.* (BaBar Collab.), *Phys. Rev. Lett.* **92**, 181801 (2004) and *Phys. Rev.* **D70**, 012007 (2004).
36. D.E. Jaffe *et al.* (CLEO Collab.), *Phys. Rev. Lett.* **86**, 5000 (2001).
37. F. Abe *et al.* (CDF Collab.), *Phys. Rev.* **D55**, 2546 (1997).
38. CDF Collab., CDF note 9015, October 2007.
39. V.M. Abazov *et al.* (D0 Collab.), *Phys. Rev.* **D74**, 092001 (2006).
40. DELPHI Collab., contrib. 449 to *Int. Europhysics Conf. on High Energy Physics*, Jerusalem, 1997.
41. R. Barate *et al.* (Aleph Collab.), *Eur. Phys. J.* **C20**, 431 (2001).
42. B. Aubert *et al.* (BaBar Collab.), *Phys. Rev. Lett.* **96**, 251802 (2006).
43. BABAR Collab., [arXiv:hep-ex/0607091v1](https://arxiv.org/abs/hep-ex/0607091v1), *33rd Int. Conf. on High Energy Physics*, Moscow, 2006.
44. E. Nakano *et al.* (Belle Collab.), *Phys. Rev.* **D73**, 112002 (2006).
45. G. Abbiendi *et al.* (OPAL Collab.), *Eur. Phys. J.* **C12**, 609 (2000).
46. M. Beneke, G. Buchalla, and I. Dunietz, *Phys. Lett.* **B393**, 132 (1997); I. Dunietz, *Eur. Phys. J.* **C7**, 197 (1999).
47. See the review on the CKM quark-mixing matrix by A. Ceccucci, Z. Ligeti, and Y. Sakai in this publication.
48. M. Okamoto, *XXIIIth Int. Symp. on Lattice Field Theory*, Dublin, July 2005, [hep-lat/0510113](https://arxiv.org/abs/hep-lat/0510113); these estimates are obtained by combining the unquenched lattice QCD calculations from A. Gray *et al.* (HPQCD Collab.), *Phys. Rev. Lett.* **95**, 212001 (2005) and S. Aoki *et al.* (JLQCD Collab.), *Phys. Rev. Lett.* **91**, 212001 (2003).
49. A. Heister *et al.* (Aleph Collab.), *Eur. Phys. J.* **C29**, 143 (2003).
50. F. Abe *et al.* (CDF Collab.), *Phys. Rev. Lett.* **82**, 3576 (1999).
51. J. Abdallah *et al.* (DELPHI Collab.), *Eur. Phys. J.* **C35**, 35 (2004).

52. G. Abbiendi *et al.* (OPAL Collab.), *Eur. Phys. J.* **C11**, 587 (1999); *Eur. Phys. J.* **C19**, 241 (2001).
53. K. Abe *et al.* (SLD Collab.), *Phys. Rev.* **D66**, 032009 (2002).
54. SLD Collab., SLAC-PUB-8568, *30th Int. Conf. on High Energy Physics*, Osaka, 2000.
55. V.M. Abazov *et al.* (D0 Collab.), *Phys. Rev. Lett.* **97**, 021802 (2006).
56. D0 Collab., D0 note 5618-CONF v1.1, March 2008.
57. D. Mohapatra *et al.* (Belle Collab.), *Phys. Rev. Lett.* **96**, 221601 (2006), updated in M. Nakao, talk at *23rd Int. Symp. on Lepton and Photon Interactions at High Energy*, Daegu, 2007.
58. B. Aubert *et al.* (BaBar Collab.), *Phys. Rev. Lett.* **98**, 151802 (2007).
59. M. Bona *et al.* (UTfit Collab.), [arXiv:hep-ph/0606167v2](http://arxiv.org/abs/hep-ph/0606167v2) and updated results at <http://utfit.roma1.infn.it/>.
60. J. Charles *et al.* (CKMfitter Collab.), *Eur. Phys. J.* **C41**, 1 (2005) & updated results at <http://ckmfitter.in2p3.fr/>.
61. K. Hartkorn and H.-G. Moser, *Eur. Phys. J.* **C8**, 381 (1999).
62. M. Acciarri *et al.* (L3 Collab.), *Phys. Lett.* **B438**, 417 (1998).
63. D. Buskulic *et al.* (Aleph Collab.), *Phys. Lett.* **B377**, 205 (1996); K. Ackerstaff *et al.* (OPAL Collab.), *Phys. Lett.* **B426**, 161 (1998); F. Abe *et al.* (CDF Collab.), *Phys. Rev.* **D59**, 032004 (1999).
64. (CDF Collab.), CDF note 7386, March 2005; CDF note 7757, August 2005; V.M. Abazov *et al.* (D0 Collab.), *Phys. Rev. Lett.* **97**, 241801 (2006).
65. D. Tonelli for the (CDF Collab.), [arXiv:hep-ex/0605038v1](http://arxiv.org/abs/hep-ex/0605038v1), May 2006.
66. F. Abe *et al.* (CDF Collab.), *Phys. Rev.* **D57**, 5382 (1998).
67. V.M. Abazov *et al.* (D0 Collab.), *Phys. Rev. Lett.* **94**, 041001 (2005); (CDF Collab.), CDF note 7409, May 2004.
68. R. Barate *et al.* (Aleph Collab.), *Phys. Lett.* **B486**, 286 (2000).
69. R. Aleksan *et al.*, *Phys. Lett.* **B316**, 567 (1993).
70. T. Aaltonen *et al.* (CDF Collab.), *Phys. Rev. Lett.* **100**, 121803 (2008).
71. V.M. Abazov *et al.* (D0 Collab.), *Phys. Rev. Lett.* **98**, 121801 (2007).

72. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **99**, 241801 (2007); A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 021803 (2008).
73. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 161802 (2008).
74. V.M. Abazov *et al.* (D0 Collab.), arXiv:0802.2255v1 [hep-ex], submitted to Phys. Rev. Lett.
75. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **98**, 151801 (2007).
76. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. **D76**, 057101 (2007).
77. M. Bona *et al.* (UTfit Collab.), arXiv:0803.0659v1 [hep-ph], March 2008.
78. P. Abreu *et al.* (DELPHI Collab.), Phys. Lett. **B289**, 199 (1992); P.D. Acton *et al.* (OPAL Collab.), Phys. Lett. **B295**, 357 (1992); D. Buskulic *et al.* (Aleph Collab.), Phys. Lett. **B361**, 221 (1995).
79. P. Abreu *et al.* (DELPHI Collab.), Z. Phys. **C68**, 375 (1995); R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C2**, 197 (1998).
80. D. Buskulic *et al.* (Aleph Collab.), Phys. Lett. **B384**, 449 (1996); J. Abdallah *et al.* (DELPHI Collab.), Eur. Phys. J. **C44**, 299 (2005).
81. R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C5**, 205 (1998).
82. J. Abdallah *et al.* (DELPHI Collab.), Phys. Lett. **B576**, 29 (2003).
83. F. Abe *et al.* (CDF Collab.), Phys. Rev. **D60**, 092005 (1999).
84. T. Affolder *et al.* (CDF Collab.), Phys. Rev. Lett. **84**, 1663 (2000).
85. Aleph, DELPHI, L3, OPAL, and SLD Collabs., “Precision electroweak measurements on the  $Z$  resonance,” Physics Reports **427**, 257 (2006); we use the  $\bar{\chi}$  average given in Eq. (5.39).
86. D. Acosta *et al.* (CDF Collab.), Phys. Rev. **D69**, 012002 (2004).
87. L. Fernández for the LHCb Collab., LHCb note 2006-047, Acta Phys. Pol. **B38**, 931 (2007).