

$\chi_{c2}(2P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

$\chi_{c2}(2P)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3929±5±2	64	UEHARA 06	BELL	10.6 $e^+e^- \rightarrow e^+e^- D\bar{D}$

$\chi_{c2}(2P)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29±10±2	64	UEHARA 06	BELL	10.6 $e^+e^- \rightarrow e^+e^- D\bar{D}$

$\chi_{c2}(2P)$ DECAY MODES

Mode
$\Gamma_1 \quad \gamma\gamma$
$\Gamma_2 \quad D\bar{D}$
$\Gamma_3 \quad D^+D^-$
$\Gamma_4 \quad D^0\bar{D}^0$

$\chi_{c2}(2P)$ PARTIAL WIDTHS

————— $\chi_{c2}(2P) \Gamma(\gamma\gamma)\Gamma(i)/\Gamma(\text{total})$ —————

$\Gamma(\gamma\gamma) \times \Gamma(D\bar{D})/\Gamma_{\text{total}}$	<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_1\Gamma_2/\Gamma$
0.18±0.05±0.03	64	¹ UEHARA 06	BELL	10.6 $e^+e^- \rightarrow e^+e^- D\bar{D}$		

¹ Assuming $B(D^+D^-) = 0.89 B(D^0\bar{D}^0)$.

$\chi_{c2}(2P)$ BRANCHING RATIOS

$\Gamma(D^+D^-)/\Gamma(D^0\bar{D}^0)$	<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_4
0.74±0.43±0.16	64	UEHARA 06	BELL	10.6 $e^+e^- \rightarrow e^+e^- D\bar{D}$		

$\chi_{c2}(2P)$ REFERENCES

UEHARA 06 PRL 96 082003 S. Uehara *et al.* (BELLE Collab.)