

$\omega(1650)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\omega(1650)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670 ± 30 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1667 ± 13 ± 6		AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ
1645 ± 8	13	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → ωηγ
1660 ± 10 ± 2		AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
1770 ± 50 ± 60	1.2M	¹ ACHASOV	03D RVUE	0.44–2.00 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
1619 ± 5		² HENNER	02 RVUE	1.2–2.0 e ⁺ e ⁻ → ρπ, ωππ
1700 ± 20		EUGENIO	01 SPEC	18 π ⁻ p → ωηn
1705 ± 26	612	³ AKHMETSHIN	00D CMD2	e ⁺ e ⁻ → ωπ ⁺ π ⁻
1820 ⁺¹⁹⁰ ₋₁₅₀		⁴ ACHASOV	98H RVUE	e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
1840 ⁺¹⁰⁰ ₋₇₀		⁵ ACHASOV	98H RVUE	e ⁺ e ⁻ → ωπ ⁺ π ⁻
1780 ⁺¹⁷⁰ ₋₃₀₀		⁶ ACHASOV	98H RVUE	e ⁺ e ⁻ → K ⁺ K ⁻
~ 2100		⁷ ACHASOV	98H RVUE	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓]
1606 ± 9		⁸ CLEGG	94 RVUE	
1662 ± 13	750	⁹ ANTONELLI	92 DM2	1.34–2.4 e ⁺ e ⁻ → ρπ, ωππ
1670 ± 20		ATKINSON	83B OMEG	20–70 γp → 3πX
1657 ± 13		CORDIER	81 DM1	e ⁺ e ⁻ → ω2π
1679 ± 34	21	ESPOSITO	80 FRAM	e ⁺ e ⁻ → 3π
1652 ± 17		COSME	79 OSPK	e ⁺ e ⁻ → 3π

¹From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the π⁺π⁻π⁰ and ANTONELLI 92 on the ωπ⁺π⁻ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

²Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

³Using the data of AKHMETSHIN 00D and ANTONELLI 92. The ρπ dominance for the energy dependence of the ω(1420) and ω(1650) width assumed.

⁴Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

⁵Using the data from ANTONELLI 92.

⁶Using the data from IVANOV 81 and BISELLO 88B.

⁷Using the data from BISELLO 91C.

⁸From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁹From the combined fit of the ρπ and ωππ final states.

$\omega(1650)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315 ± 35 OUR ESTIMATE				

• • • We do not use the following data for averages, fits, limits, etc. • • •

222 ± 25 ± 20		AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ
114 ± 14	13	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → ωηγ
230 ± 30 ± 20		AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
490 ⁺²⁰⁰ ₋₁₅₀ ± 130	1.2M	¹⁰ ACHASOV	03D RVUE	0.44–2.00 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
250 ± 14		¹¹ HENNER	02 RVUE	1.2–2.0 e ⁺ e ⁻ → ρπ, ωππ
250 ± 50		EUGENIO	01 SPEC	18 π ⁻ p → ωηπ
370 ± 25	612	¹² AKHMETSHIN	00D CMD2	e ⁺ e ⁻ → ωπ ⁺ π ⁻
113 ± 20		¹³ CLEGG	94 RVUE	
280 ± 24	750	¹⁴ ANTONELLI	92 DM2	1.34–2.4e ⁺ e ⁻ → ρπ, ωππ
160 ± 20		ATKINSON	83B OMEG	20–70 γp → 3πX
136 ± 46		CORDIER	81 DM1	e ⁺ e ⁻ → ω2π
99 ± 49	21	ESPOSITO	80 FRAM	e ⁺ e ⁻ → 3π
42 ± 17		COSME	79 OSPK	e ⁺ e ⁻ → 3π

¹⁰From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the π⁺π⁻π⁰ and ANTONELLI 92 on the ωπ⁺π⁻ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

¹¹Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

¹²Using the data of AKHMETSHIN 00D and ANTONELLI 92. The ρπ dominance for the energy dependence of the ω(1420) and ω(1650) width assumed.

¹³From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹⁴From the combined fit of the ρπ and ωππ final states.

ω(1650) DECAY MODES

Mode	Fraction (Γ _i /Γ)
Γ ₁ ρπ	seen
Γ ₂ ωππ	seen
Γ ₃ ωη	seen
Γ ₄ e ⁺ e ⁻	seen

ω(1650) Γ(i)Γ(e⁺e⁻)/Γ²(total)

Γ(ρπ)/Γ_{total} × Γ(e⁺e⁻)/Γ_{total} Γ₁/Γ × Γ₄/Γ

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.3 ± 0.1 ± 0.1		AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
1.2 ^{+0.4} _{-0.1} ± 0.8	1.2M	^{15,16} ACHASOV	03D RVUE	0.44–2.00 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
0.921 ± 0.230		^{17,18} CLEGG	94 RVUE	
0.479 ± 0.050	750	^{19,20} ANTONELLI	92 DM2	1.34–2.4e ⁺ e ⁻ → ρπ, ωππ

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
7.0 ± 0.5		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 ± 0.9 ± 1.3	1.2M	15,16 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		21 AKHMETSHIN	00D CMD2	1.2–1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		17,18 CLEGG	94 RVUE	
6.07 ± 0.61	750	19,20 ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.57 ± 0.06		13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
<6	90		22 AKHMETSHIN	03B CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$

- ¹⁵ Calculated by us from the cross section at the peak.
- ¹⁶ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- ¹⁷ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- ¹⁸ From the partial and leptonic width given by the authors.
- ¹⁹ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.
- ²⁰ From the product of the leptonic width and partial branching ratio given by the authors.
- ²¹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
- ²² $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.

$\omega(1650)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 0.35	1.2M	23 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.620 ± 0.014		24 HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 0.65	1.2M	23 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.380 ± 0.014		24 HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE (units 10^{-7}) EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 18	1.2M	^{24,25} ACHASOV	03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
32 ± 1		²⁴ HENNER	02	RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

²³From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

²⁴Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

²⁵Calculated by us from the cross section at the peak.

$\omega(1650)$ REFERENCES

AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)