

$\Upsilon(4S)$
or $\Upsilon(10580)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\Upsilon(4S)$ MASS

<u>VALUE (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.5794 ± 0.0012 OUR AVERAGE			
10.5793 ± 0.0004 ± 0.0012	AUBERT	05Q	BABR $e^+e^- \rightarrow$ hadrons
10.5800 ± 0.0035	¹ BEBEK	87	CLEO $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10.5774 ± 0.0010	² LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BESSON 85.			
² No systematic error given.			

$\Upsilon(4S)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.5 ± 2.5 OUR AVERAGE			
20.7 ± 1.6 ± 2.5	AUBERT	05Q	BABR $e^+e^- \rightarrow$ hadrons
20 ± 2 ± 4	BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
25 ± 2.5	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

$\Upsilon(4S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $B\bar{B}$	> 96 %	95%
Γ_2 B^+B^-	(51.6 ± 0.6) %	
Γ_3 D_S^+ anything + c.c.	(17.8 ± 2.6) %	
Γ_4 $B^0\bar{B}^0$	(48.4 ± 0.6) %	
Γ_5 $J/\psi K_S^0 (J/\psi, \eta_c) K_S^0$	< 4 × 10 ⁻⁷	90%
Γ_6 non- $B\bar{B}$	< 4 %	95%
Γ_7 e^+e^-	(1.57 ± 0.08) × 10 ⁻⁵	
Γ_8 $\rho^+\rho^-$	< 5.7 × 10 ⁻⁶	90%
Γ_9 $J/\psi(1S)$ anything	< 1.9 × 10 ⁻⁴	95%
Γ_{10} D^{*+} anything + c.c.	< 7.4 %	90%
Γ_{11} ϕ anything	(7.1 ± 0.6) %	
Γ_{12} $\phi\eta$	< 1.8 × 10 ⁻⁶	90%
Γ_{13} $\phi\eta'$	< 4.3 × 10 ⁻⁶	90%
Γ_{14} $\rho\eta$	< 1.3 × 10 ⁻⁶	90%
Γ_{15} $\rho\eta'$	< 2.5 × 10 ⁻⁶	90%

Γ_{16}	$\Upsilon(1S)$ anything	< 4	$\times 10^{-3}$	90%
Γ_{17}	$\Upsilon(1S)\pi^+\pi^-$	(8.1 ± 0.6)	$\times 10^{-5}$	
Γ_{18}	$\Upsilon(1S)\eta$	(1.96 ± 0.11)	$\times 10^{-4}$	
Γ_{19}	$\Upsilon(2S)\pi^+\pi^-$	(8.6 ± 1.3)	$\times 10^{-5}$	
Γ_{20}	\bar{d} anything	< 1.3	$\times 10^{-5}$	90%

$\Upsilon(4S)$ PARTIAL WIDTHS


$\Gamma(e^+e^-)$

Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.272 ± 0.029 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
$0.321 \pm 0.017 \pm 0.029$	AUBERT	05Q	BABR $e^+e^- \rightarrow$ hadrons
$0.28 \pm 0.05 \pm 0.01$	³ ALBRECHT	95E	ARG $e^+e^- \rightarrow$ hadrons
$0.192 \pm 0.007 \pm 0.038$	BESSION	85	CLEO $e^+e^- \rightarrow$ hadrons
0.283 ± 0.037	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

³ Using LEYAOUANC 77 parametrization of $\Gamma(s)$.

WEIGHTED AVERAGE
 0.272 ± 0.029 (Error scaled by 1.5)



$\Upsilon(4S)$ BRANCHING RATIOS

$B\bar{B}$ DECAYS

The ratio of branching fraction to charged and neutral B mesons is often derived assuming isospin invariance in the decays, and relies on the knowledge of the B^+/B^0 lifetime ratio. "OUR EVALUATION" is obtained based on averages of rescaled data listed below. The average and

rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account the common dependence of the measurement on the value of the lifetime ratio.

$\Gamma(B^+ B^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>
0.516 ± 0.006 OUR EVALUATION	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$

$\Gamma(D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.178 ± 0.021 ± 0.016	⁴ ARTUSO 05B CLE3 $e^+ e^- \rightarrow D_s X$

⁴ ARTUSO 05B reports $[\Gamma(\Upsilon(4S) \rightarrow D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (8.0 \pm 0.2 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(B^0 \bar{B}^0)/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.484 ± 0.006 OUR EVALUATION	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.487 ± 0.010 ± 0.008	⁵ AUBERT,B 05H BABR $\Upsilon(4S) \rightarrow \bar{B}B \rightarrow D^* \ell \nu_\ell$

⁵ Direct measurement. This value is averaged with the value extracted from the $\Gamma(B^+ B^-) / \Gamma(B^0 \bar{B}^0)$ measurements.

$\Gamma(B^+ B^-)/\Gamma(B^0 \bar{B}^0)$	Γ_2/Γ_4
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.065 ± 0.026 OUR EVALUATION	

<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.006 ± 0.036 ± 0.031	⁶ AUBERT 04F BABR $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$
1.01 ± 0.03 ± 0.09	⁶ HASTINGS 03 BELL $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow \text{dileptons}$
1.058 ± 0.084 ± 0.136	⁷ ATHAR 02 CLEO $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow D^* \ell \nu$
1.10 ± 0.06 ± 0.05	⁸ AUBERT 02 BABR $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow (c\bar{c})K^*$
1.04 ± 0.07 ± 0.04	⁹ ALEXANDER 01 CLEO $\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K^*$

⁶ HASTINGS 03 and AUBERT 04F assume $\tau(B^+) / \tau(B^0) = 1.083 \pm 0.017$.

⁷ ATHAR 02 assumes $\tau(B^+) / \tau(B^0) = 1.074 \pm 0.028$. Supersedes BARISH 95.

⁸ AUBERT 02 assumes $\tau(B^+) / \tau(B^0) = 1.062 \pm 0.029$.

⁹ ALEXANDER 01 assumes $\tau(B^+) / \tau(B^0) = 1.066 \pm 0.024$.

$\Gamma(J/\psi K_S^0 (J/\psi, \eta_c) K_S^0)/\Gamma_{\text{total}}$	Γ_5/Γ
Forbidden by CP invariance.	

<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	¹⁰ TAJIMA	07A	BELL $\Upsilon(4S) \rightarrow B^0 \bar{B}^0$

¹⁰ $\Upsilon(4S)$ with $CP = +1$ decays to the final state with $CP = -1$.

————— non- $B\bar{B}$ DECAYS —————

$\Gamma(\text{non-}B\bar{B})/\Gamma_{\text{total}}$	Γ_6/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.04	95	BARISH	96B	CLEO $e^+ e^-$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ **Γ_7/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.57±0.08 OUR AVERAGE				
1.55±0.04±0.07		AUBERT	05Q BABR	$e^+e^- \rightarrow \text{hadrons}$
2.77±0.50±0.49		¹¹ ALBRECHT	95E ARG	$e^+e^- \rightarrow \text{hadrons}$

¹¹ Using LEYAOUANC 77 parametrization of $\Gamma(s)$.

$\Gamma(\rho^+\rho^-)/\Gamma_{\text{total}}$ **Γ_8/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.7 × 10⁻⁶	90	AUBERT	08BO BABR	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ **Γ_9/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.9	95	¹² ABE	02D BELL	$e^+e^- \rightarrow J/\psi X \rightarrow \ell^+\ell^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4.7	90	¹² AUBERT	01C BABR	$e^+e^- \rightarrow J/\psi X \rightarrow \ell^+\ell^-X$

¹² Uses $B(J/\psi \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(D^{*+} \text{ anything} + \text{ c.c.})/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.074	90	¹³ ALEXANDER	90C CLEO	e^+e^-

¹³ For $x > 0.473$.

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.1 ±0.1±0.6		HUANG	07 CLEO	$\Upsilon(4S) \rightarrow \phi X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.23	90	¹⁴ ALEXANDER	90C CLEO	e^+e^-

¹⁴ For $x > 0.52$.

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.8	90	¹⁵ BELOUS	09 BELL	$e^+e^- \rightarrow \phi\eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<2.5	90	AUBERT, BE	06F BABR	$e^+e^- \rightarrow \phi\eta$

¹⁵ Using all intermedite branching fraction values from PDG 08.

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.3	90	¹⁶ BELOUS	09 BELL	$e^+e^- \rightarrow \phi\eta'$

¹⁶ Using all intermedite branching fraction values from PDG 08.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	¹⁷ BELOUS	09 BELL	$e^+e^- \rightarrow \rho\eta$

¹⁷ Using all intermedite branching fraction values from PDG 08.

$\Gamma(\rho\eta')/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	¹⁸ BELOUS 09	BELL	$e^+e^- \rightarrow \rho\eta'$

¹⁸ Using all intermediate branching fraction values from PDG 08.

$\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	ALEXANDER 90C	CLEO	e^+e^-

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

8.1 ± 0.6 OUR AVERAGE

8.5 ± 1.3 ± 0.2	113 ± 16	¹⁹ SOKOLOV 09	BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
8.00 ± 0.64 ± 0.27	430	²⁰ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.8 ± 4.0 ± 0.3		^{21,22} SOKOLOV 07	BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
9.0 ± 1.5 ± 0.2	167 ± 19	²³ AUBERT 06R	BABR	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
<12	90	GLENN 99	CLE2	e^+e^-

¹⁹ SOKOLOV 09 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (0.211 \pm 0.030 \pm 0.014) \times 10^{-5}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

²¹ SOKOLOV 07 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (4.42 \pm 0.81 \pm 0.56) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²² According to the authors, systematic errors were underestimated.

²³ Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (2.23 \pm 0.25 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.96 ± 0.06 ± 0.09	56	²⁴ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\ell^+\ell^-$

²⁴ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ **Γ_{18}/Γ_{17}**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.41 ± 0.40 ± 0.12	56	²⁵ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-(\pi^0)\ell^+\ell^-$
--------------------	----	---------------------------	------	--

²⁵ Not independent of other values reported by AUBERT 08BP.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{19}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$0.86 \pm 0.11 \pm 0.07$		220	²⁶ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
--	--	-----	---------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.88 \pm 0.17 \pm 0.08$		97 ± 15	²⁷ AUBERT 06R	BABR	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
< 3.9		90	GLENN 99	CLE2	e^+e^-

²⁶ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.

²⁷ Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = (1.69 \pm 0.26 \pm 0.20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ **Γ_{19}/Γ_{17}**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

$1.16 \pm 0.16 \pm 0.14$	220	²⁸ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
--------------------------	-----	---------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

²⁸ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

$\Gamma(\bar{d} \text{ anything})/\Gamma_{\text{total}}$ **Γ_{20}/Γ**

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.3	90	ASNER 07	CLEO	$e^+e^- \rightarrow \bar{d}X$
------------------------------	----	----------	------	-------------------------------

$\Upsilon(4S)$ REFERENCES

BELOUS 09	PL B681 400	K. Belous <i>et al.</i>	(BELLE Collab.)
SOKOLOV 09	PR D79 051103R	A. Sokolov <i>et al.</i>	(BELLE Collab.)
AUBERT 08BO	PR D78 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG 08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER 07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
HUANG 07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)
SOKOLOV 07	PR D75 071103R	A. Sokolov <i>et al.</i>	(BELLE Collab.)
TAJIMA 07A	PRL 99 211601	O. Tajima <i>et al.</i>	(BELLE Collab.)
AUBERT 06R	PRL 96 232001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE 06F	PR D74 111103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
ARTUSO 05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT 05Q	PR D72 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B 05H	PRL 95 042001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 04F	PR D69 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)
HASTINGS 03	PR D67 052004	N.C. Hastings <i>et al.</i>	(BELLE Collab.)
ABE 02D	PRL 88 052001	K. Abe <i>et al.</i>	(BELLE Collab.)
ATHAR 02	PR D66 052003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT 02	PR D65 032001	B. Aubert <i>et al.</i>	(BaBar Collab.)
ALEXANDER 01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AUBERT 01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BaBar Collab.)
GLENN 99	PR D59 052003	S. Glenn <i>et al.</i>	(CLEO Collab.)
BARISH 96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
ALBRECHT 95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARISH 95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
ALEXANDER 90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)
BEBEK 87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
BESSON 85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK 85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)
LEYAOUANC 77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)