

$$J^G(J^{PC}) = 0^-(1^{--})$$

J/ψ(1S) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.916±0.011 OUR AVERAGE				
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	¹ ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3097.5 ±0.3		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^-A$
3096.93 ±0.09	502	³ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ±1		⁴ BRANDELIK 79C	DASP	e^+e^-

¹ Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

² Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

³ Superseded by ARTAMONOV 00.

⁴ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

J/ψ(1S) WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.9± 2.8 OUR AVERAGE Error includes scale factor of 1.1.				
96.1± 3.2	13k	⁵ ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4± 8.9		BAI 95B	BES	e^+e^-
91 ±11 ±6		⁶ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 ⁺ ₋ 6.1 5.8		⁷ HSUEH 92	RVUE	See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
94.1± 2.7		⁸ ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7± 3.5	7.8k	⁵ AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

⁷ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

⁸ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 \pm 0.30) %	
Γ_3 $g g g$	(64.1 \pm 1.0) %	
Γ_4 $\gamma g g$	(8.8 \pm 0.5) %	
Γ_5 $e^+ e^-$	(5.94 \pm 0.06) %	
Γ_6 $e^+ e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+ \mu^-$	(5.93 \pm 0.06) %	
Decays involving hadronic resonances		
Γ_8 $\rho \pi$	(1.69 \pm 0.15) %	S=2.4
Γ_9 $\rho^0 \pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$	
Γ_{10} $a_2(1320) \rho$	(1.09 \pm 0.22) %	
Γ_{11} $\omega \pi^+ \pi^+ \pi^- \pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$	
Γ_{12} $\omega \pi^+ \pi^- \pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$	
Γ_{13} $\omega \pi^+ \pi^-$	(8.6 \pm 0.7) $\times 10^{-3}$	S=1.1
Γ_{14} $\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$	
Γ_{15} $K^*(892)^0 \bar{K}^*(892)^0$	(2.3 \pm 0.7) $\times 10^{-4}$	
Γ_{16} $K^*(892)^\pm \bar{K}^*(892)^\mp$	(1.00 $^{+0.22}_{-0.40}$) $\times 10^{-3}$	
Γ_{17} $K^*(892)^\pm \bar{K}^*(800)^\mp$	(1.1 $^{+1.0}_{-0.6}$) $\times 10^{-3}$	
Γ_{18} $\eta K^*(892)^0 \bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	
Γ_{19} $K^*(892)^0 \bar{K}_2^*(1430)^0 + c.c.$	(6.0 \pm 0.6) $\times 10^{-3}$	
Γ_{20} $K^*(892)^0 \bar{K}_2^*(1770)^0 + c.c. \rightarrow$ $K^*(892)^0 K^- \pi^+ + c.c.$	(6.9 \pm 0.9) $\times 10^{-4}$	
Γ_{21} $\omega K^*(892) \bar{K} + c.c.$	(6.1 \pm 0.9) $\times 10^{-3}$	
Γ_{22} $K^+ \bar{K}^*(892)^- + c.c.$	(5.12 \pm 0.30) $\times 10^{-3}$	
Γ_{23} $K^+ \bar{K}^*(892)^- + c.c. \rightarrow$ $K^+ K^- \pi^0$	(1.97 \pm 0.20) $\times 10^{-3}$	
Γ_{24} $K^+ \bar{K}^*(892)^- + c.c. \rightarrow$ $K^0 K^\pm \pi^\mp$	(3.0 \pm 0.4) $\times 10^{-3}$	
Γ_{25} $K^0 \bar{K}^*(892)^0 + c.c.$	(4.39 \pm 0.31) $\times 10^{-3}$	
Γ_{26} $K^0 \bar{K}^*(892)^0 + c.c. \rightarrow$ $K^0 K^\pm \pi^\mp$	(3.2 \pm 0.4) $\times 10^{-3}$	
Γ_{27} $K_1(1400)^\pm K^\mp$	(3.8 \pm 1.4) $\times 10^{-3}$	
Γ_{28} $\bar{K}^*(892)^0 K^+ \pi^- + c.c.$	seen	
Γ_{29} $\omega \pi^0 \pi^0$	(3.4 \pm 0.8) $\times 10^{-3}$	
Γ_{30} $b_1(1235)^\pm \pi^\mp$	[b] (3.0 \pm 0.5) $\times 10^{-3}$	
Γ_{31} $\omega K^\pm K_S^0 \pi^\mp$	[b] (3.4 \pm 0.5) $\times 10^{-3}$	
Γ_{32} $b_1(1235)^0 \pi^0$	(2.3 \pm 0.6) $\times 10^{-3}$	
Γ_{33} $\eta K^\pm K_S^0 \pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$	
Γ_{34} $\phi K^*(892) \bar{K} + c.c.$	(2.18 \pm 0.23) $\times 10^{-3}$	

Γ ₃₅	$\omega K\bar{K}$		$(1.6 \pm 0.5) \times 10^{-4}$	
Γ ₃₆	$\omega f_0(1710) \rightarrow \omega K\bar{K}$		$(4.8 \pm 1.1) \times 10^{-4}$	
Γ ₃₇	$\phi 2(\pi^+\pi^-)$		$(1.66 \pm 0.23) \times 10^{-3}$	
Γ ₃₈	$\Delta(1232)^{++} \bar{p}\pi^-$		$(1.6 \pm 0.5) \times 10^{-3}$	
Γ ₃₉	$\omega\eta$		$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
Γ ₄₀	$\phi K\bar{K}$		$(1.83 \pm 0.24) \times 10^{-3}$	S=1.5
Γ ₄₁	$\phi f_0(1710) \rightarrow \phi K\bar{K}$		$(3.6 \pm 0.6) \times 10^{-4}$	
Γ ₄₂	$\phi f_2(1270)$		$(7.2 \pm 1.3) \times 10^{-4}$	
Γ ₄₃	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
Γ ₄₄	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)}$	[b]	$(1.03 \pm 0.13) \times 10^{-3}$	
Γ ₄₅	$\phi f_2'(1525)$		$(8 \pm 4) \times 10^{-4}$	S=2.7
Γ ₄₆	$\phi\pi^+\pi^-$		$(8.7 \pm 0.8) \times 10^{-4}$	
Γ ₄₇	$\phi\pi^0\pi^0$		$(5.6 \pm 1.6) \times 10^{-4}$	
Γ ₄₈	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ ₄₉	$\omega f_1(1420)$		$(6.8 \pm 2.4) \times 10^{-4}$	
Γ ₅₀	$\phi\eta$		$(7.5 \pm 0.8) \times 10^{-4}$	S=1.5
Γ ₅₁	$\Xi^0 \Xi^0$		$(1.20 \pm 0.24) \times 10^{-3}$	
Γ ₅₂	$\Xi(1530)^- \Xi^+$		$(5.9 \pm 1.5) \times 10^{-4}$	
Γ ₅₃	$\rho K^- \bar{\Sigma}(1385)^0$		$(5.1 \pm 3.2) \times 10^{-4}$	
Γ ₅₄	$\omega\pi^0$		$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
Γ ₅₅	$\phi\eta'(958)$		$(4.0 \pm 0.7) \times 10^{-4}$	S=2.1
Γ ₅₆	$\phi f_0(980)$		$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
Γ ₅₇	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$		$(2.2 \pm 0.4) \times 10^{-4}$	
Γ ₅₈	$\phi f_0(980) \rightarrow \phi\pi^0\pi^0$		$(1.7 \pm 0.7) \times 10^{-4}$	
Γ ₅₉	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$		$(3.2 \pm 1.0) \times 10^{-4}$	
Γ ₆₀	$\Xi(1530)^0 \Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$	
Γ ₆₁	$\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)}$	[b]	$(3.1 \pm 0.5) \times 10^{-4}$	
Γ ₆₂	$\phi f_1(1285)$		$(2.6 \pm 0.5) \times 10^{-4}$	S=1.1
Γ ₆₃	$\eta\pi^+\pi^-$		$(4.0 \pm 1.7) \times 10^{-4}$	
Γ ₆₄	$\rho\eta$		$(1.93 \pm 0.23) \times 10^{-4}$	
Γ ₆₅	$\omega\eta'(958)$		$(1.82 \pm 0.21) \times 10^{-4}$	
Γ ₆₆	$\omega f_0(980)$		$(1.4 \pm 0.5) \times 10^{-4}$	
Γ ₆₇	$\rho\eta'(958)$		$(1.05 \pm 0.18) \times 10^{-4}$	
Γ ₆₈	$a_2(1320)^\pm \pi^\mp$	[b]	$< 4.3 \times 10^{-3}$	CL=90%
Γ ₆₉	$K\bar{K}_2^*(1430) + \text{c.c.}$		$< 4.0 \times 10^{-3}$	CL=90%
Γ ₇₀	$K_1(1270)^\pm K^\mp$		$< 3.0 \times 10^{-3}$	CL=90%
Γ ₇₁	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$		$< 2.9 \times 10^{-3}$	CL=90%
Γ ₇₂	$\phi\pi^0$		$< 6.4 \times 10^{-6}$	CL=90%
Γ ₇₃	$\phi\eta(1405) \rightarrow \phi\eta\pi\pi$		$< 2.5 \times 10^{-4}$	CL=90%
Γ ₇₄	$\omega f_2'(1525)$		$< 2.2 \times 10^{-4}$	CL=90%
Γ ₇₅	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$		$< 2.52 \times 10^{-4}$	CL=90%
Γ ₇₆	$\Sigma(1385)^0 \bar{\Lambda}$		$< 2 \times 10^{-4}$	CL=90%
Γ ₇₇	$\Delta(1232)^+ \bar{p}$		$< 1 \times 10^{-4}$	CL=90%

Γ_{78}	$\Theta(1540)\bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	< 1.1	$\times 10^{-5}$	CL=90%
Γ_{79}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1	$\times 10^{-5}$	CL=90%
Γ_{80}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6	$\times 10^{-5}$	CL=90%
Γ_{81}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6	$\times 10^{-5}$	CL=90%
Γ_{82}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1	$\times 10^{-5}$	CL=90%
Γ_{83}	$\Sigma^0 \bar{\Lambda}$	< 9	$\times 10^{-5}$	CL=90%

Decays into stable hadrons

Γ_{84}	$2(\pi^+ \pi^-) \pi^0$	$(5.5 \pm 0.4) \%$		
Γ_{85}	$3(\pi^+ \pi^-) \pi^0$	$(2.9 \pm 0.6) \%$		
Γ_{86}	$\pi^+ \pi^- \pi^0$	$(2.07 \pm 0.12) \%$		S=1.6
Γ_{87}	$\pi^+ \pi^- \pi^0 K^+ K^-$	$(1.94 \pm 0.15) \%$		
Γ_{88}	$4(\pi^+ \pi^-) \pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$		
Γ_{89}	$\pi^+ \pi^- K^+ K^-$	$(6.6 \pm 0.5) \times 10^{-3}$		
Γ_{90}	$\pi^+ \pi^- K^+ K^- \eta$	$(1.84 \pm 0.28) \times 10^{-3}$		
Γ_{91}	$\pi^0 \pi^0 K^+ K^-$	$(2.45 \pm 0.31) \times 10^{-3}$		
Γ_{92}	$K \bar{K} \pi$	$(6.1 \pm 1.0) \times 10^{-3}$		
Γ_{93}	$2(\pi^+ \pi^-)$	$(3.55 \pm 0.23) \times 10^{-3}$		
Γ_{94}	$3(\pi^+ \pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$		
Γ_{95}	$2(\pi^+ \pi^- \pi^0)$	$(1.61 \pm 0.21) \%$		
Γ_{96}	$2(\pi^+ \pi^-) \eta$	$(2.29 \pm 0.24) \times 10^{-3}$		
Γ_{97}	$3(\pi^+ \pi^-) \eta$	$(7.2 \pm 1.5) \times 10^{-4}$		
Γ_{98}	$\rho \bar{\rho}$	$(2.17 \pm 0.07) \times 10^{-3}$		
Γ_{99}	$\rho \bar{\rho} \pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$		S=1.1
Γ_{100}	$\rho \bar{\rho} \pi^+ \pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$		S=1.3
Γ_{101}	$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	[c] $(2.3 \pm 0.9) \times 10^{-3}$		S=1.9
Γ_{102}	$\rho \bar{\rho} \eta$	$(2.00 \pm 0.12) \times 10^{-3}$		
Γ_{103}	$\rho \bar{\rho} \rho$	< 3.1	$\times 10^{-4}$	CL=90%
Γ_{104}	$\rho \bar{\rho} \omega$	$(1.10 \pm 0.15) \times 10^{-3}$		S=1.3
Γ_{105}	$\rho \bar{\rho} \eta'(958)$	$(2.1 \pm 0.4) \times 10^{-4}$		
Γ_{106}	$\rho \bar{\rho} \phi$	$(4.5 \pm 1.5) \times 10^{-5}$		
Γ_{107}	$n \bar{n}$	$(2.2 \pm 0.4) \times 10^{-3}$		
Γ_{108}	$n \bar{n} \pi^+ \pi^-$	$(4 \pm 4) \times 10^{-3}$		
Γ_{109}	$\Sigma^+ \bar{\Sigma}^-$	$(1.50 \pm 0.24) \times 10^{-3}$		
Γ_{110}	$\Sigma^0 \bar{\Sigma}^0$	$(1.29 \pm 0.09) \times 10^{-3}$		
Γ_{111}	$2(\pi^+ \pi^-) K^+ K^-$	$(5.0 \pm 0.5) \times 10^{-3}$		
Γ_{112}	$\rho \bar{n} \pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$		
Γ_{113}	$n N(1440)$	seen		
Γ_{114}	$n N(1520)$	seen		
Γ_{115}	$n N(1535)$	seen		
Γ_{116}	$\Xi^- \bar{\Xi}^+$	$(8.5 \pm 1.6) \times 10^{-4}$		S=1.5
Γ_{117}	$\Lambda \bar{\Lambda}$	$(1.61 \pm 0.15) \times 10^{-3}$		S=2.0
Γ_{118}	$\Lambda \bar{\Sigma}^- \pi^+ (\text{or c.c.})$	[b] $(8.3 \pm 0.7) \times 10^{-4}$		S=1.2

Γ_{119}	$pK^-\bar{\Lambda}$	$(8.9 \pm 1.6) \times 10^{-4}$	
Γ_{120}	$2(K^+K^-)$	$(7.6 \pm 0.9) \times 10^{-4}$	
Γ_{121}	$pK^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{122}	K^+K^-	$(2.37 \pm 0.31) \times 10^{-4}$	
Γ_{123}	$K_S^0 K_L^0$	$(1.46 \pm 0.26) \times 10^{-4}$	S=2.7
Γ_{124}	$\Lambda\bar{\Lambda}\eta$	$(2.6 \pm 0.7) \times 10^{-4}$	
Γ_{125}	$\Lambda\bar{\Lambda}\pi^0$	$< 6.4 \times 10^{-5}$	CL=90%
Γ_{126}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{127}	$\pi^+\pi^-$	$(1.47 \pm 0.23) \times 10^{-4}$	
Γ_{128}	$\Lambda\bar{\Sigma} + \text{c.c.}$	$< 1.5 \times 10^{-4}$	CL=90%
Γ_{129}	$K_S^0 K_S^0$	$< 1 \times 10^{-6}$	CL=95%

Radiative decays

Γ_{130}	3γ	$(1.2 \pm 0.4) \times 10^{-5}$	
Γ_{131}	4γ	$< 9 \times 10^{-6}$	CL=90%
Γ_{132}	5γ	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{133}	$\gamma\eta_c(1S)$	$(1.7 \pm 0.4) \%$	S=1.6
Γ_{134}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	$(1.2 \begin{smallmatrix} +2.7 \\ -1.1 \end{smallmatrix}) \times 10^{-6}$	
Γ_{135}	$\gamma\pi^+\pi^-2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{136}	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{137}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	
Γ_{138}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	[d] $(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
Γ_{139}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
Γ_{140}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{141}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%
Γ_{142}	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
Γ_{143}	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90%
Γ_{144}	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{145}	$\gamma\eta'(958)$	$(5.16 \pm 0.15) \times 10^{-3}$	S=1.1
Γ_{146}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{147}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{148}	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$	
Γ_{149}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{150}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{151}	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
Γ_{152}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3
Γ_{153}	$\gamma f_2(1270)$	$(1.43 \pm 0.11) \times 10^{-3}$	
Γ_{154}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(8.5 \begin{smallmatrix} +1.2 \\ -0.9 \end{smallmatrix}) \times 10^{-4}$	S=1.2
Γ_{155}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(4.0 \pm 1.0) \times 10^{-4}$	
Γ_{156}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$	
Γ_{157}	$\gamma\eta$	$(1.104 \pm 0.034) \times 10^{-3}$	
Γ_{158}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$	

Γ_{159}	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	
Γ_{160}	$\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	
Γ_{161}	$\gamma f_2'(1525)$	$(4.5 \begin{smallmatrix} +0.7 \\ -0.4 \end{smallmatrix}) \times 10^{-4}$	
Γ_{162}	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$	
Γ_{163}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$	
Γ_{164}	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$	
Γ_{165}	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
Γ_{166}	$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{167}	$\gamma \rho \bar{\rho}$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{168}	$\gamma \eta(2225)$	$(3.3 \pm 0.5) \times 10^{-4}$	
Γ_{169}	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
Γ_{170}	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
Γ_{171}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.6 \pm 0.4) \times 10^{-4}$	
Γ_{172}	$\gamma X(1835) \rightarrow \gamma \rho \bar{\rho}$	$(7.5 \begin{smallmatrix} +1.9 \\ -0.9 \end{smallmatrix}) \times 10^{-5}$	
Γ_{173}	$\gamma (K \bar{K} \pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ_{174}	$\gamma \pi^0$	$(3.49 \begin{smallmatrix} +0.33 \\ -0.30 \end{smallmatrix}) \times 10^{-5}$	
Γ_{175}	$\gamma \rho \bar{\rho} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
Γ_{176}	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{177}	$\gamma f_0(2200)$		
Γ_{178}	$\gamma f_J(2220)$	$> 2.50 \times 10^{-3}$	CL=99.9%
Γ_{179}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$(8 \pm 4) \times 10^{-5}$	
Γ_{180}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 3.6 \times 10^{-5}$	
Γ_{181}	$\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho}$	$(1.5 \pm 0.8) \times 10^{-5}$	
Γ_{182}	$\gamma f_0(1500)$	$(1.01 \pm 0.32) \times 10^{-4}$	
Γ_{183}	$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%

Weak decays

Γ_{184}	$D^- e^+ \nu_e + \text{c.c.}$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{185}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{186}	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 3.6 \times 10^{-5}$	CL=90%
Γ_{187}	$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90%
Γ_{188}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{189}	$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{190}	$\gamma \gamma$	C	$< 5 \times 10^{-6}$	CL=90%
Γ_{191}	$e^\pm \mu^\mp$	LF	$< 1.1 \times 10^{-6}$	CL=90%
Γ_{192}	$e^\pm \tau^\mp$	LF	$< 8.3 \times 10^{-6}$	CL=90%
Γ_{193}	$\mu^\pm \tau^\mp$	LF	$< 2.0 \times 10^{-6}$	CL=90%

Other decays

Γ_{194} invisible $< 7 \times 10^{-4}$ CL=90%

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes $\rho\bar{\rho}\pi^+\pi^-\gamma$ and excludes $\rho\bar{\rho}\eta$, $\rho\bar{\rho}\omega$, $\rho\bar{\rho}\eta'$.

[d] See the "Note on the $\eta(1405)$ " in the $\eta(1405)$ Particle Listings.

[e] For a narrow state A with mass less than 960 MeV.

$J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
74.1 ± 8.1	BAI	95B	BES e^+e^-
59 ± 24	BALDINI-...	75	FRAG e^+e^-
59 ± 14	BOYARSKI	75	MRK1 e^+e^-
50 ± 25	ESPOSITO	75B	FRAM e^+e^-

$\Gamma(e^+e^-)$

Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.55 ± 0.14 ± 0.02 OUR EVALUATION				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.71 ± 0.16	13k	⁹ ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ± 0.19	7.8k	⁹ AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ± 0.39		BAI	95B	BES e^+e^-
$5.36^{+0.29}_{-0.28}$		¹⁰ HSUEH	92	RVUE See Υ mini-review
4.72 ± 0.35		ALEXANDER	89	RVUE See Υ mini-review
4.4 ± 0.6		¹⁰ BRANDELIK	79C	DASP e^+e^-
4.6 ± 0.8		¹¹ BALDINI-...	75	FRAG e^+e^-
4.8 ± 0.6		BOYARSKI	75	MRK1 e^+e^-
4.6 ± 1.0		ESPOSITO	75B	FRAM e^+e^-

⁹ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

¹⁰ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

¹¹ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+\mu^-)$

Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5.13 ± 0.52	BAI	95B	BES e^+e^-
4.8 ± 0.6	BOYARSKI	75	MRK1 e^+e^-
5 ± 1	ESPOSITO	75B	FRAM e^+e^-

$\Gamma(\gamma\gamma)$					Γ_{190}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<5.4	90	BRANDELIK	79C	DASP	e^+e^-

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel i in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (keV)		DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 0.8		¹² BALDINI-...	75	FRAG	e^+e^-
3.9 ± 0.8		¹² ESPOSITO	75B	FRAM	e^+e^-

¹²Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_5\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	

332.3 ± 6.4 ± 4.8 ANASHIN 10 KEDR 3.097 $e^+e^- \rightarrow e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

350 ± 20		BRANDELIK	79C	DASP	e^+e^-
320 ± 70		¹³ BALDINI-...	75	FRAG	e^+e^-
340 ± 90		¹³ ESPOSITO	75B	FRAM	e^+e^-
360 ± 100		¹³ FORD	75	SPEC	e^+e^-

¹³Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	

334 ± 5 OUR AVERAGE

331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR	3.097 $e^+e^- \rightarrow \mu^+\mu^-$
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90		DASP	75	DASP	e^+e^-
380 ± 50		¹⁴ ESPOSITO	75B	FRAM	e^+e^-

¹⁴Data redundant with branching ratios or partial widths above.

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{12}\Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	

2.2 ± 0.3 ± 0.2 170 AUBERT 06D BABR 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

53.6±5.0±0.4	788	¹⁵ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ
---------------------	-----	----------------------	-----------	---

¹⁵ AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 47.8 ± 3.1 ± 3.2 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

33±4±1	317 ± 23	^{16,17} AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
---------------	----------	-------------------------	-----------	--

¹⁶ Dividing by 2/3 to take into account that B(K^{*0} → K⁺π⁻) = 2/3.

¹⁷ AUBERT 07AK reports [$\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(K₂^{*}(1430) → Kπ)] = 16.4 ± 1.1 ± 1.4 eV which we divide by our best value B(K₂^{*}(1430) → Kπ) = (49.9 ± 1.2) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.8±0.4±0.3	110 ± 14	¹⁸ AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
--------------------	----------	----------------------	-----------	--

¹⁸ Dividing by 2/3 to take into account that B(K^{*0} → K⁺π⁻) = 2/3.

$\Gamma(K^+\bar{K}^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

29.0±1.7±1.3	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K [*] (892) ⁻ γ
---------------------	--------	----------	---

$\Gamma(K^+\bar{K}^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

10.96±0.85±0.70	155	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ γ
------------------------	-----	--------	----------	---

$\Gamma(K^+\bar{K}^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

16.76±1.70±1.00	89	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ
------------------------	----	--------	----------	--

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

26.6±2.5±1.5	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁰ $\bar{K}^*(892)^0$ γ
---------------------	--------	----------	--

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

17.70±1.70±1.00	94	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ
------------------------	----	--------	----------	--

$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{35}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.70±1.98±0.03	24	¹⁹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ω K ⁺ K ⁻ γ
-----------------------	----	----------------------	-----------	--

¹⁹ AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 3.3 ± 1.3 ± 1.2 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{37}\Gamma_5/\Gamma$**

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------------	------	-------------	------	---------

0.96±0.19±0.01	35	²⁰ AUBERT	06D BABR	10.6 e ⁺ e ⁻ → φ 2(π ⁺ π ⁻) γ
-----------------------	----	----------------------	----------	--

²⁰ AUBERT 06D reports [$\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = (0.47 ± 0.09 ± 0.03) × 10⁻² keV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{46}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.8 ± 0.4 OUR AVERAGE

4.52±0.48±0.04	254 ± 23	²¹ SHEN	09 BELL	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
----------------	----------	--------------------	---------	--

5.33±0.71±0.05	103	²² AUBERT, BE	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
----------------	-----	--------------------------	----------	--

²¹ SHEN 09 reports 4.50 ± 0.41 ± 0.26 eV from a measurement of [$\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] assuming B(φ(1020) → K⁺K⁻) = (49.2 ± 0.6) × 10⁻², which we rescale to our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

²² AUBERT, BE 06D reports [$\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 2.61 ± 0.30 ± 0.18 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{47}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.15±0.88±0.03	23	²³ AUBERT, BE	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ π ⁰ γ
-----------------------	----	--------------------------	----------	--

²³ AUBERT, BE 06D reports [$\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 1.54 ± 0.40 ± 0.16 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{50}\Gamma_5/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

6.1±2.7±0.4	6	²⁴ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → φ η γ
--------------------	---	----------------------	-----------	--

²⁴ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{57} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.21 ± 0.23 OUR AVERAGE Error includes scale factor of 1.2.

1.48 ± 0.27 ± 0.09 60 ± 11 ²⁵ SHEN 09 BELL 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

1.02 ± 0.24 ± 0.01 20 ± 5 ²⁶ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

²⁵ Multiplied by 2/3 to take into account the $\phi \pi^+ \pi^-$ mode only. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

²⁶ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{58} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.96 ± 0.40 ± 0.01 7.0 ± 2.8 ²⁷ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

²⁷ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{63} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.24 ± 0.98 ± 0.03 9 ²⁸ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

²⁸ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.74 \pm 0.28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{15} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.28 ± 0.40 ± 0.11 25 ± 8 ²⁹ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

²⁹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{42} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.0 ± 0.7 ± 0.1 44 ± 7 ^{30,31} AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

³⁰ Using $B(\phi \rightarrow (K + K)^-) = (49.3 \pm 0.6)\%$.

³¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 3.41 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{84} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

303 ± 5 ± 18 4990 AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{86}\Gamma_5/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.122±0.005±0.008		AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ

$$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{87}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0±4.3±6.4	768	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{89}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
36.3±1.3±2.1	1586 ± 58	AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

33.6±2.7±2.7	233	³² AUBERT	05D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
³² Superseded by AUBERT 07AK.				

$$\Gamma(\pi^+\pi^-K^+K^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{90}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	³³ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ

³³ AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 10.2 ± 1.3 ± 0.8 eV which we divide by our best value B(η → 2γ) = (39.31 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{91}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.6±1.1±1.3	203 ± 16	AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{93}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
19.5±1.4±1.3	270	AUBERT	05D BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)γ

$$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{94}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → 3(π ⁺ π ⁻)γ

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{95}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±0.5±1.0	761	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰)γ

$$\Gamma(2(\pi^+\pi^-\eta)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{96}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	³⁴ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ

³⁴ AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 5.16 ± 0.85 ± 0.39 eV which we divide by our best value B(η → 2γ) = (39.31 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{98}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.6±0.9 OUR AVERAGE		Error includes scale factor of 1.2.			
12.0±0.6±0.5	438	AUBERT	06B	$e^+e^- \rightarrow p\bar{p}\gamma$	
9.7±1.7		³⁵ ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$	
³⁵ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.					

$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{110}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
6.4±1.2±0.6		AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$	

$\Gamma(2(\pi^+\pi^-)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{111}\Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.75±0.23±0.17	205	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$	

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{117}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.7±0.9±0.7		AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$	

$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{120}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.11±0.39±0.30	156 ± 15	AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.0 ± 0.7 ± 0.6	38	³⁶ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$	
³⁶ Superseded by AUBERT 07AK.					

J/ψ(1S) BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) × Γ(e⁺e⁻)/Γ_{total} above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.877±0.005 OUR AVERAGE					
0.878±0.005		BAI	95B BES	e^+e^-	
0.86 ± 0.02		BOYARSKI	75 MRK1	e^+e^-	

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.135±0.003	37,38	SETH	04 RVUE	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.17 ± 0.02	37	BOYARSKI	75 MRK1	e^+e^-	
³⁷ Included in Γ(hadrons)/Γ _{total} .					
³⁸ Using B(J/ψ → ℓ ⁺ ℓ ⁻) = (5.90 ± 0.09)% from RPP-2002 and R = 2.28 ± 0.04 determined by a fit to data from BAI 00 and BAI 02C.					

$\Gamma(ggg)/\Gamma_{\text{total}}$ **Γ_3/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
64.1 ± 1.0	6 M	³⁹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- + \text{hadrons}$

³⁹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+ \ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma \eta_C)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$ **Γ_4/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.79 ± 1.05	200 k	⁴⁰ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

⁴⁰ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma gg)/\Gamma(ggg)$ **Γ_4/Γ_3**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13.7 ± 0.1 ± 0.7	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ **Γ_5/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.94 ± 0.06 OUR AVERAGE				
5.945 ± 0.067 ± 0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B	BES $e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ **Γ_6/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8 ± 1.3 ± 0.4	⁴¹ ARMSTRONG	96	E760 $\bar{p} p \rightarrow e^+ e^- \gamma$

⁴¹ For $E_\gamma > 100$ MeV.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_7/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.93 ± 0.06 OUR AVERAGE				
5.960 ± 0.065 ± 0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B	BES $e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$ Γ_5/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.998±0.012 OUR AVERAGE			
1.002±0.021±0.013	⁴² ANASHIN	10 KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
0.997±0.012±0.006	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.00 ±0.07	BAI	95B BES	e^+e^-
1.00 ±0.05	BOYARSKI	75 MRK1	e^+e^-
0.91 ±0.15	ESPOSITO	75B FRAM	e^+e^-
0.93 ±0.10	FORD	75 SPEC	e^+e^-

⁴² Not independent of the corresponding measurements of $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$.

———— HADRONIC DECAYS ————

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_8/Γ


<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.69 ±0.15 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.				
2.18 ±0.19	43,44	AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
2.184±0.005±0.201	220k	44,45 BAI	04H BES	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
2.091±0.021±0.116	44,46	BAI	04H BES	$\psi(2S) \rightarrow \pi^+\pi^-J/\psi$
1.21 ±0.20		BAI	96D BES	$e^+e^- \rightarrow \rho\pi$
1.42 ±0.01 ±0.19		COFFMAN	88 MRK3	e^+e^-
1.3 ±0.3	150	FRANKLIN	83 MRK2	e^+e^-
1.6 ±0.4	183	ALEXANDER	78 PLUT	e^+e^-
1.33 ±0.21		BRANDELIK	78B DASP	e^+e^-
1.0 ±0.2	543	BARTEL	76 CNTR	e^+e^-
1.3 ±0.3	153	JEAN-MARIE	76 MRK1	e^+e^-

⁴³ From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).

⁴⁴ Not independent of their $B(\pi^+\pi^-\pi^0)$.

⁴⁵ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

⁴⁶ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
$0.328 \pm 0.005 \pm 0.027$	COFFMAN 88	MRK3	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.35 ± 0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL 76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE 76	MRK1	e^+e^-

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.2	OUR AVERAGE			
$11.7 \pm 0.7 \pm 2.5$	7584	AUGUSTIN 89	DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4 ± 4.5	36	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85 ± 34	140	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.40 \pm 0.06 \pm 0.04$	170	⁴⁷ AUBERT 06D	BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

⁴⁷ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6±0.7 OUR AVERAGE		Error includes scale factor of 1.1.		
9.7±0.6±0.6	788	⁴⁸ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8±1.6	215	BURMESTER	77D PLUT	e^+e^-
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
⁴⁸ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 47.8 \pm 3.1 \pm 3.2$ eV.				

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±0.6 OUR AVERAGE				
4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	e^+e^-
4.0±1.6	70	BURMESTER	77D PLUT	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.7±0.1		25 ± 8	⁴⁹ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5	90		VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$
⁴⁹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (1.28 \pm 0.40 \pm 0.11) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(K^*(892)^\pm\bar{K}^*(892)^\mp)/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.19^{+0.11}_{-0.32}	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0\pi^\mp\pi^0$

$\Gamma(K^*(892)^\pm\bar{K}^*(800)^\mp)/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.18^{+0.94}_{-0.54}	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0\pi^\mp\pi^0$

$\Gamma(\eta K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.15±0.13±0.22	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^-K^-\pi^+$

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{19}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±0.6 OUR AVERAGE				
5.9±0.6±0.2	317 ± 23	^{50,51} AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
6.7±2.6	40	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

⁵⁰ Using $B(K_2^*(1430)^0 \rightarrow K\pi) = (49.9 \pm 1.2)\%$.

⁵¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (32.9 \pm 2.3 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.12 ± 0.30 OUR AVERAGE				
5.2 ± 0.4 ± 0.1		⁵² AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K^+ K^*(892)^- \gamma$
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

⁵² AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ \bar{K}^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (29.0 \pm 1.7 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.97 ± 0.20 ± 0.05	155	⁵³ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K^+ K^- \pi^0 \gamma$

⁵³ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ \bar{K}^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.96 \pm 0.85 \pm 0.70) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.4 ± 0.1	89	⁵⁴ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

⁵⁴ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ \bar{K}^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (16.76 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

4.39±0.31 OUR AVERAGE

4.8 ±0.5 ±0.1		55 AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$
3.96±0.15±0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33±0.12±0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ±0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
----------	----	----------	---------	--

⁵⁵ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (26.6 \pm 2.5 \pm 1.5) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.})$ Γ_{25}/Γ_{22}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.82±0.05±0.09 COFFMAN 88 MRK3 $J/\psi \rightarrow K \bar{K}^*(892) + \text{c.c.}$

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

3.2±0.4±0.1 94 ⁵⁶ AUBERT 08S BABR 10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

⁵⁶ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (17.70 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

3.8±0.8±1.2 ⁵⁷ BAI 99C BES e^+e^-

⁵⁷ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen ⁵⁸ ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

⁵⁸ A $K_0^*(800)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(\omega \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

3.4±0.3±0.7 509 AUGUSTIN 89 DM2 $J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

30±5 OUR AVERAGE

31±6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29±7	87	BURMESTER	77D PLUT	e^+e^-

$\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
23 ± 3 ± 5	229	AUGUSTIN	89 DM2	$e^+ e^-$

$\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.2 ± 3.4	232 ± 23	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\phi K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.3 OUR AVERAGE				
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(1.6 ± 0.5) OUR AVERAGE				
1.36 ± 0.50 ± 0.10	24	⁵⁹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega K^+ K^- \gamma$
19.8 ± 2.1 ± 3.9		⁶⁰ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

⁵⁹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega K^+ K^-) \cdot B(\eta \rightarrow 3\pi) = 3.3 \pm 1.3 \pm 0.2 \text{ eV}$.

⁶⁰ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

$\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.8 ± 1.1 ± 0.3	^{61,62} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

⁶¹ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

⁶² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
16.6 ± 2.3 OUR AVERAGE				
17.3 ± 3.3 ± 1.2	35	⁶³ AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
16.0 ± 1.0 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

⁶³ Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$.

$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.23 ± 0.40	332	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$


Γ_{39}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
2.352 ± 0.273	5k	⁶⁴ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta$
1.44 ± 0.40 ± 0.14	13	⁶⁵ AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta$

⁶⁴ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

⁶⁵ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

WEIGHTED AVERAGE
1.74±0.20 (Error scaled by 1.6)



$\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$


Γ_{40}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
18.3 ± 2.4 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
48 $^{+20}_{-16}$ ± 6	9.0 $^{+3.7}_{-3.0}$	^{66,67} HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		⁶⁸ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	e^+e^-

⁶⁶ We have multiplied K^+K^- measurement by 2 to obtain $K\bar{K}$.

⁶⁷ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

⁶⁸ Addition of ϕK^+K^- and $\phi K^0\bar{K}^0$ branching ratios.



$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6±0.2±0.6	69,70 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

⁶⁹ Including interference with $f_2'(1525)$.

⁷⁰ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.72±0.13±0.02	44 ± 7	71,72	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.45 90 FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

< 0.37 90 VANNUCCI 77 MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

⁷¹ Using $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2})\%$

⁷² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.02 \pm 0.65 \pm 0.33) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.09±0.28	233	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03±0.13 OUR AVERAGE				
1.00±0.04±0.21	631 ± 25	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.19±0.04±0.25	754 ± 27	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*+}$
0.86±0.18±0.22	56	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.03±0.24±0.25	68	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(8+4) OUR AVERAGE Error includes scale factor of 2.7.				
12.3±0.6±2.0	^{73,74}	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
4.8±1.8	46 ⁷³	GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$
⁷³ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.				
⁷⁴ Including interference with $f_0(1710)$.				

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94±0.09 OUR AVERAGE Error includes scale factor of 1.2.				
0.96±0.13	103 ⁷⁵	AUBERT,BE 06D	BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
1.09±0.02±0.13		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ±0.9	23	FELDMAN 77	MRK1	$e^+ e^-$
⁷⁵ Derived by us. AUBERT,BE 06D measures $\Gamma(J/\psi \rightarrow e^+ e^-) \times B(J/\psi \rightarrow \phi\pi^+\pi^-) \times B(\phi \rightarrow K^+ K^-) = (2.61 \pm 0.30 \pm 0.18) \text{ eV}$				

$\Gamma(\phi\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.56±0.16				
	23	⁷⁶ AUBERT,BE 06D	BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
⁷⁶ Derived by us. AUBERT,BE 06D measures $\Gamma(J/\psi \rightarrow e^+ e^-) \times B(J/\psi \rightarrow \phi\pi^0\pi^0) \times B(\phi \rightarrow K^+ K^-) = (1.54 \pm 0.40 \pm 0.16) \text{ eV}$				

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(7.2±0.8) OUR AVERAGE				
7.4±0.6±1.4	227 ± 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4±0.9±1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ±0.6±1.0	163 ± 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{49}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8^{+1.9}_{-1.6}±1.7				
	111 ⁺³¹ ₋₂₆	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$

Γ_{50}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.08	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
1.4 ± 0.6 ± 0.1	6	⁷⁷ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \phi\eta\gamma$
0.898 ± 0.024 ± 0.089		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.64 ± 0.04 ± 0.11	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.661 ± 0.045 ± 0.078		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$
⁷⁷ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+ K^-) \cdot B(\eta \rightarrow \gamma\gamma) = 0.84 \pm 0.37 \pm 0.05 \text{ eV}$.				

WEIGHTED AVERAGE
0.75 ± 0.08 (Error scaled by 1.5)



$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

Γ_{51}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.20 ± 0.12 ± 0.21	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$

Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.59 ± 0.09 ± 0.12	75 ± 11	HENRARD	87 DM2	$e^+ e^-$

$\Gamma(\rho K^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

Γ_{53}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.26 ± 0.18	89	EATON	84 MRK2	$e^+ e^-$


$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

Γ_{54}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
0.538 ± 0.012 ± 0.065	2090	⁷⁸ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

⁷⁸ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

WEIGHTED AVERAGE
0.45±0.05 (Error scaled by 1.4)




$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$

Γ_{55}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.40 ± 0.07 OUR AVERAGE			Error includes scale factor of 2.1. See the ideogram below.		
0.546 ± 0.031 ± 0.056			ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.41 ± 0.03 ± 0.08		167	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.308 ± 0.034 ± 0.036			COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3	90	VANNUCCI	77 MRK1	e^+e^-
-------	----	----------	---------	----------



$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$

Γ_{56}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(3.2±0.9) OUR AVERAGE		Error includes scale factor of 1.9.		
4.6±0.4±0.8		⁷⁹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6±0.6	50	⁷⁹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

⁷⁹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{57}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.182±0.042±0.005	19.5 ± 4.5	^{80,81} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

⁸⁰ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

⁸¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (1.01 \pm 0.22 \pm 0.08) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0)/\Gamma_{\text{total}}$

Γ_{58}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.171±0.073±0.004	7.0 ± 2.8	^{82,83} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

⁸² Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

⁸³ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (0.95 \pm 0.39 \pm 0.10) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.23±0.75±0.73	52	ABLIKIM	08F	BES $J/\psi \rightarrow \eta\phi f_0(980)$

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.32±0.12±0.07	24 ± 9	HENRARD	87	DM2 e^+e^-

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.31±0.05 OUR AVERAGE				
0.30±0.03±0.07	74 ± 8	HENRARD	87	DM2 $e^+e^- \rightarrow \Sigma^{*-}$
0.34±0.04±0.07	77 ± 9	HENRARD	87	DM2 $e^+e^- \rightarrow \Sigma^{*+}$
0.29±0.11±0.10	26	EATON	84	MRK2 $e^+e^- \rightarrow \Sigma^{*-}$
0.31±0.11±0.11	28	EATON	84	MRK2 $e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{62}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(2.6±0.5) OUR AVERAGE				Error includes scale factor of 1.1.
3.2±0.6±0.4		JOUSSET	90	DM2 $J/\psi \rightarrow \phi 2(\pi^+\pi^-)$
2.1±0.5±0.4	25	⁸⁴ JOUSSET	90	DM2 $J/\psi \rightarrow \phi\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.6±0.2±0.1	16 ± 6	BECKER	87	MRK3 $J/\psi \rightarrow \phi K \bar{K} \pi$

⁸⁴ We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.40±0.17±0.03	9	⁸⁵ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$
⁸⁵ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \eta\pi^+\pi^-) \cdot B(\eta \rightarrow 3\pi) = 0.51 \pm 0.22 \pm 0.03$ eV.				

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$ Γ_{64}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.182±0.021 OUR AVERAGE				
0.226±0.043	218	⁸⁶ ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\eta'$
0.18 $^{+0.10}_{-0.08}$ ±0.03	6	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.166±0.017±0.019		COFFMAN	88	MRK3 $e^+e^- \rightarrow 3\pi\eta'$

⁸⁶ Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{66}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.41 \pm 0.27 \pm 0.47$		⁸⁷ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
⁸⁷ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.				

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ Γ_{67}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.105 ± 0.018 OUR AVERAGE				
$0.083 \pm 0.030 \pm 0.012$	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.114 \pm 0.014 \pm 0.016$		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

$\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<43	90	BRAUNSCH...	76 DASP	$e^+ e^-$

$\Gamma(K \bar{K}_2^*(1430) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<40	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	BRAUNSCH...	76 DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
-----	----	-------------	---------	--

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{70}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.0	90	⁸⁸ BAI	99C BES	$e^+ e^-$

⁸⁸ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{71}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<29	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.4	90	ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$
------	----	---------	---------	-------------------------------------

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.5	90	⁸⁹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

⁸⁹ Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\omega f_2'(1525))/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	⁹⁰ VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<2.8	90	⁹⁰ FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
⁹⁰ Re-evaluated assuming $B(f_2'(1525) \rightarrow K\bar{K}) = 0.713$.				

$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.52	90	ABLIKIM 10c	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

$\Gamma(\Sigma(1385)^0 \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.2	90	HENRARD 87	DM2	e^+e^-

$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	HENRARD 87	DM2	e^+e^-

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI 04G	BES2	e^+e^-

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	BAI 04G	BES2	e^+e^-

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BAI 04G	BES2	e^+e^-

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.6	90	BAI 04G	BES2	e^+e^-

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI 04G	BES2	e^+e^-

$\Gamma(\Sigma^0 \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	HENRARD 87	DM2	e^+e^-


STABLE HADRONS

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.1 ± 0.5 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
5.46 ± 0.34 ± 0.14	4990	⁹¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+e^- \rightarrow$ hadrons
3.64 ± 0.52	1500	BURMESTER	77D PLUT	e^+e^-
4 ± 1	675	JEAN-MARIE	76 MRK1	e^+e^-

⁹¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = 0.303 \pm 0.005 \pm 0.018$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
4.1 ± 0.5 (Error scaled by 2.4)



$\Gamma(\omega\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-)\pi^0)$ Γ_{13}/Γ_{84}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3	⁹² JEAN-MARIE	76	MRK1 e^+e^-
-----	--------------------------	----	---------------

⁹² Final state $(\pi^+\pi^-)\pi^0$ under the assumption that $\pi\pi$ is isospin 0.

$\Gamma(3\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{85}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.029±0.006 OUR AVERAGE				
0.028±0.009	11	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$
0.029±0.007	181	JEAN-MARIE 76	MRK1	e^+e^-

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{86}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
20.7 ±1.2 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
23.6 ±2.1 ±0.5	256	⁹³ AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
21.8 ±1.9		^{94,95} AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
21.84±0.05±2.01	220k	^{95,96} BAI 04H	BES	e^+e^-
20.91±0.21±1.16		^{95,97} BAI 04H	BES	e^+e^-
15 ±2	168	FRANKLIN 83	MRK2	e^+e^-


⁹³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.789 \pm 0.015$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹⁴ From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).

⁹⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁹⁶ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

⁹⁷ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+\pi^-\pi^0K^+K^-)/\Gamma_{\text{total}}$ **Γ_{87}/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

1.79±0.29 OUR AVERAGE Error includes scale factor of 2.2.

1.93±0.14±0.05	768	⁹⁸ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$
----------------	-----	----------------------	-----------	---

1.2 ±0.3	309	VANNUCCI	77 MRK1	e^+e^-
----------	-----	----------	---------	----------

⁹⁸ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = 0.1070 \pm 0.0043 \pm 0.0064$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{88}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

90±30	13	JEAN-MARIE	76 MRK1	e^+e^-
--------------	----	------------	---------	----------

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ **Γ_{89}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

6.6±0.5 OUR AVERAGE

6.5±0.4±0.2	1.6k	⁹⁹ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
-------------	------	----------------------	-----------	--

7.2±2.3	205	VANNUCCI	77 MRK1	e^+e^-
---------	-----	----------	---------	----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±0.7±0.2	233	¹⁰⁰ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
-------------	-----	-----------------------	----------	--

⁹⁹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (36.3 \pm 1.3 \pm 2.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰⁰ Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (33.6 \pm 2.7 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-K^+K^-\eta)/\Gamma_{\text{total}}$ **Γ_{90}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

1.84±0.28±0.05	73	¹⁰¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$
-----------------------	----	-----------------------	-----------	--

¹⁰¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.2 \pm 1.3 \pm 0.8) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\pi^0K^+K^-)/\Gamma_{\text{total}}$ **Γ_{91}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.45±0.31±0.06	203 ± 16	¹⁰² AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
-----------------------	----------	-----------------------	-----------	--

¹⁰² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^0\pi^0K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (13.6 \pm 1.1 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ **Γ_{92}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
61 ± 10 OUR AVERAGE				
55.2 ± 12.0	25	FRANKLIN	83	MRK2 $e^+e^- \rightarrow K^+K^-\pi^0$
78.0 ± 21.0	126	VANNUCCI	77	MRK1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ **Γ_{93}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.55 ± 0.23 OUR AVERAGE				
3.53 ± 0.12 ± 0.29	1107	¹⁰³ ABLIKIM	05H BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $J/\psi \pi^+ \pi^-, J/\psi \rightarrow$ $2(\pi^+ \pi^-)$
3.51 ± 0.34 ± 0.09	270	¹⁰⁴ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+ \pi^-)\gamma$
4.0 ± 1.0	76	JEAN-MARIE	76 MRK1	e^+e^-

¹⁰³ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

¹⁰⁴ AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (19.5 \pm 1.4 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ **Γ_{94}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
43 ± 4 OUR AVERAGE				
43.0 ± 2.9 ± 2.8	496	¹⁰⁵ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+ \pi^-)\gamma$
40 ± 20	32	JEAN-MARIE	76 MRK1	e^+e^-

¹⁰⁵ Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ **Γ_{95}/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.62 ± 0.09 ± 0.19				
	761	¹⁰⁶ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+ \pi^- \pi^0)\gamma$

¹⁰⁶ Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ **Γ_{96}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.29 ± 0.24 OUR AVERAGE				
2.35 ± 0.39 ± 0.20	85	¹⁰⁷ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+ \pi^-)\eta\gamma$
2.26 ± 0.08 ± 0.27	4839	ABLIKIM	05C BES2	$e^+e^- \rightarrow 2(\pi^+ \pi^-)\eta$

¹⁰⁷ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow 2(\pi^+ \pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 5.16 \pm 0.85 \pm 0.39$ eV.

$\Gamma(3(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ **Γ_{97}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.24 ± 0.96 ± 1.11				
	616	ABLIKIM	05C BES2	$e^+e^- \rightarrow 3(\pi^+ \pi^-)\eta$

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ **Γ_{98}/Γ**

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

2.17±0.07 OUR AVERAGE

2.18±0.16±0.07	317	¹⁰⁸ WU	06	BELL	$B^+ \rightarrow \rho\bar{\rho}K^+$
2.26±0.01±0.14	63316	BAI	04E	BES2	$e^+e^- \rightarrow J/\psi$
1.97±0.22	99	BALDINI	98	FENI	e^+e^-
1.91±0.04±0.30		PALLIN	87	DM2	e^+e^-
2.16±0.07±0.15	1420	EATON	84	MRK2	e^+e^-
2.5 ±0.4	133	BRANDELIK	79C	DASP	e^+e^-
2.0 ±0.5		BESCH	78	BONA	e^+e^-
2.2 ±0.2	331	¹⁰⁹ PERUZZI	78	MRK1	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ±0.3	48	ANTONELLI	93	SPEC	e^+e^-
----------	----	-----------	----	------	----------

¹⁰⁸WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.013 \pm 0.034) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰⁹Assuming angular distribution $(1+\cos^2\theta)$.

$\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$ **Γ_{99}/Γ**

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

1.19±0.08 OUR AVERAGE Error includes scale factor of 1.1.

1.33±0.02±0.11	11k	ABLIKIM	09B	BES2	e^+e^-
1.13±0.09±0.09	685	EATON	84	MRK2	e^+e^-
1.4 ±0.4		BRANDELIK	79C	DASP	e^+e^-
1.00±0.15	109	PERUZZI	78	MRK1	e^+e^-


$\Gamma(\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{100}/Γ**

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

6.0 ±0.5 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

6.46±0.17±0.43	1435	EATON	84	MRK2	e^+e^-
3.8 ±1.6	48	BESCH	81	BONA	e^+e^-
5.5 ±0.6	533	PERUZZI	78	MRK1	e^+e^-

WEIGHTED AVERAGE
 6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(\rho^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{101}/Γ

Including $\rho^+\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho^+\eta)/\Gamma_{\text{total}}$

Γ_{102}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
$1.91 \pm 0.02 \pm 0.17$	13k	¹¹⁰ ABLIKIM	09	BES2 e^+e^-
$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79C	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-

¹¹⁰ From the combination of $\rho^+\eta \rightarrow \rho^+\gamma\gamma$ and $\rho^+\eta \rightarrow \rho^+\pi^+\pi^-\pi^0$ channels.

$\Gamma(\rho^+\rho)/\Gamma_{\text{total}}$

Γ_{103}/Γ


VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.31	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

Γ_{104}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.10±0.15 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
0.98±0.03±0.14	2449	ABLIKIM	08	BES2 e^+e^-
1.10±0.17±0.18	486	EATON	84	MRK2 e^+e^-
1.6 ±0.3	77	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
1.10±0.15 (Error scaled by 1.3)



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$

Γ_{105}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21 ±0.04 OUR AVERAGE				
0.200±0.023±0.028	265 ± 31	¹¹¹ ABLIKIM	09	BES2 e^+e^-
0.68 ±0.23 ±0.17	19	EATON	84	MRK2 e^+e^-
1.8 ±0.6	19	PERUZZI	78	MRK1 e^+e^-

¹¹¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$

Γ_{106}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.45±0.13±0.07	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$ **Γ_{107}/Γ**
VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
0.22 ± 0.04 OUR AVERAGE

0.231 ± 0.049	79	BALDINI	98	FENI	e^+e^-
0.18 ± 0.09		BESCH	78	BONA	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.190 ± 0.055	40	ANTONELLI	93	SPEC	e^+e^-

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{108}/Γ**
VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
3.8 ± 3.6

3.8 ± 3.6	5	BESCH	81	BONA	e^+e^-
-----------	---	-------	----	------	----------

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ **Γ_{109}/Γ**
VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
1.50 ± 0.10 ± 0.22

1.50 ± 0.10 ± 0.22	399	ABLIKIM	080	BES2	$e^+e^- \rightarrow J/\psi$
--------------------	-----	---------	-----	------	-----------------------------

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ **Γ_{110}/Γ**
VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
1.29 ± 0.09 OUR AVERAGE

1.15 ± 0.24 ± 0.03		¹¹² AUBERT	07BD	BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$
1.33 ± 0.04 ± 0.11	1779	ABLIKIM	06	BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884 ± 30	PALLIN	87	DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84	MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78	MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.4 ± 2.6	3	BESCH	81	BONA	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$
¹¹² AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (6.4 \pm 1.2 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$ **Γ_{111}/Γ**
VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
47 ± 7 OUR AVERAGE Error includes scale factor of 1.3.

49.8 ± 4.2 ± 3.4	205	¹¹³ AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow \omega K^+K^- 2(\pi^+\pi^-)\gamma$
31 ± 13	30	VANNUCCI	77	MRK1	e^+e^-

¹¹³ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ **Γ_{112}/Γ**
VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
2.12 ± 0.09 OUR AVERAGE


2.36 ± 0.02 ± 0.21	59k	ABLIKIM	06k	BES2	$J/\psi \rightarrow p\pi^-\bar{n}$
2.47 ± 0.02 ± 0.24	55k	ABLIKIM	06k	BES2	$J/\psi \rightarrow \bar{p}\pi^+n$
2.02 ± 0.07 ± 0.16	1288	EATON	84	MRK2	$e^+e^- \rightarrow p\pi^-$
1.93 ± 0.07 ± 0.16	1191	EATON	84	MRK2	$e^+e^- \rightarrow \bar{p}\pi^+$
1.7 ± 0.7	32	BESCH	81	BONA	$e^+e^- \rightarrow p\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA	$e^+e^- \rightarrow \bar{p}\pi^+$
2.16 ± 0.29	194	PERUZZI	78	MRK1	$e^+e^- \rightarrow p\pi^-$
2.04 ± 0.27	204	PERUZZI	78	MRK1	$e^+e^- \rightarrow \bar{p}\pi^+$

$\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$

Γ_{116}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.85±0.16 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
0.70±0.06±0.12	132 ± 11	HENRARD	87 DM2	$e^+ e^- \rightarrow \Xi^- \Xi^+$
1.14±0.08±0.20	194	EATON	84 MRK2	$e^+ e^- \rightarrow \Xi^- \Xi^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Xi^- \Xi^+$

WEIGHTED AVERAGE
0.85±0.16 (Error scaled by 1.5)



$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$


Γ_{117}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61±0.15 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
1.93±0.21±0.05	114	AUBERT	07BD BABR	10.6 $e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
2.03±0.03±0.15	8887	ABLIKIM	06 BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
2.0 $^{+0.5}_{-0.4}$ ± 0.1	46	115 WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08±0.06±0.24	631	BAI	98G BES	$e^+ e^-$
1.38±0.05±0.20	1847	PALLIN	87 DM2	$e^+ e^-$
1.58±0.08±0.19	365	EATON	84 MRK2	$e^+ e^-$
2.6 ± 1.6	5	BESCH	81 BONA	$e^+ e^-$
1.1 ± 0.2	196	PERUZZI	78 MRK1	$e^+ e^-$

114 AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.7 \pm 0.9 \pm 0.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

115 WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+)$

$= (1.013 \pm 0.034) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$				Γ_{117}/Γ_{98}
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.90^{+0.15}_{-0.14} \pm 0.10$	¹¹⁶ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$	

¹¹⁶ Not independent of other $J/\psi \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$ branching ratios reported by WU 06.

$\Gamma(\Lambda\bar{\Sigma}^- \pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$				Γ_{118}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.2.
0.770 ± 0.051 ± 0.083	335	¹¹⁷ ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
0.747 ± 0.056 ± 0.076	254	¹¹⁷ ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
0.90 ± 0.06 ± 0.16	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
1.11 ± 0.06 ± 0.20	342 ± 18	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
1.53 ± 0.17 ± 0.38	135	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+ \pi^-$
1.38 ± 0.21 ± 0.35	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$

¹¹⁷ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(pK^-\bar{\Lambda})/\Gamma_{\text{total}}$				Γ_{119}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.89 ± 0.07 ± 0.14	307	EATON	84 MRK2	e^+e^-

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ **Γ_{120}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

0.76±0.09 OUR AVERAGE

0.74±0.09±0.02	156 ± 15	¹¹⁸ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
----------------	----------	-----------------------	-----------	---

1.4 $^{+0.5}_{-0.4} \pm 0.2$	11.0 $^{+4.3}_{-3.5}$	¹¹⁹ HUANG	03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$
------------------------------	-----------------------	----------------------	---------	----------------------------------

0.7 ± 0.3		VANNUCCI	77 MRK1	$e^+ e^-$
-----------	--	----------	---------	-----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72±0.17±0.02	38	¹²⁰ AUBERT	05D BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
----------------	----	-----------------------	----------	---

¹¹⁸ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.11 \pm 0.39 \pm 0.30) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹¹⁹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

¹²⁰ Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.0 \pm 0.7 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho K^- \bar{\Sigma}^0)/\Gamma_{\text{total}}$ **Γ_{121}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

0.29±0.06±0.05	90	EATON	84 MRK2	$e^+ e^-$
----------------	----	-------	---------	-----------

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ **Γ_{122}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

(2.37±0.31) OUR AVERAGE

2.39±0.24±0.22	107	BALTRUSAIT..85D	MRK3	$e^+ e^-$
----------------	-----	-----------------	------	-----------

2.2 ± 0.9	6	BRANDELIK	79C DASP	$e^+ e^-$
-----------	---	-----------	----------	-----------

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ **Γ_{123}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------


(1.46±0.26) OUR AVERAGE Error includes scale factor of 2.7. See the ideogram below.

1.82±0.04±0.13	2155 ± 45	¹²¹ BAI	04A BES2	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
----------------	-----------	--------------------	----------	--

1.18±0.12±0.18		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
----------------	--	---------	--------	-------------------------------------

1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	$e^+ e^-$
----------------	----	-----------------	------	-----------

¹²¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.62±0.60±0.44	44	122 ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

¹²² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.64	90	123	ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ±0.7±0.8	11	BAI	98G BES	e^+e^-
2.2 ±0.5±0.5	19 ± 4	HENRARD	87 DM2	e^+e^-

¹²³ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	124 ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

¹²⁴ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(1.47±0.23) OUR AVERAGE				
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ±0.5	5	BRANDELIK	78B DASP	e^+e^-
1.6 ±1.6	1	VANNUCCI	77 MRK1	e^+e^-

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.15	90	PERUZZI 78	MRK1	$e^+e^- \rightarrow \Lambda X$

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	¹²⁵ BAI 04D	BES	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.052	90	¹²⁵ BALTRUSAIT..85C	MRK3	e^+e^-
--------	----	--------------------------------	------	----------

¹²⁵Forbidden by CP.

————— RADIATIVE DECAYS —————

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$12 \pm 3 \pm 2$		$24.2^{+7.2}_{-6.0}$	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	PARTRIDGE 80	CBAL	e^+e^-
-----	----	--------------	------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.7 ± 0.4 OUR AVERAGE Error includes scale factor of 1.6.

$2.06 \pm 0.32 \pm 0.03$	¹²⁶	MITCHELL 09	CLEO	$e^+e^- \rightarrow \gamma X$
--------------------------	----------------	-------------	------	-------------------------------

1.27 ± 0.36		GAISER 86	CBAL	$J/\psi \rightarrow \gamma X$
-----------------	--	-----------	------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.79 ± 0.20	273 ± 43	¹²⁷ AUBERT 06E	BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
-----------------	--------------	---------------------------	------	--

seen	16	BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$
------	----	----------------	------	----------------------------------

¹²⁶MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33.6 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹²⁷Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.2^{+2.7}_{-1.1} \pm 0.3$	$1.2^{+2.8}_{-1.1}$	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(\gamma\pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.2 \pm 3.1$	¹²⁸ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

¹²⁸ 4π mass less than 2.0 GeV.

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 ± 1.0 OUR AVERAGE			
$5.85 \pm 0.3 \pm 1.05$	¹²⁹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+ \pi^-$
$7.8 \pm 1.2 \pm 2.4$	¹²⁹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$

¹²⁹ Broad enhancement at 1700 MeV.

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 2.2 \pm 0.9$	BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+ \pi^-$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
$1.66 \pm 0.1 \pm 0.58$	^{130,131} BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$3.8 \pm 0.3 \pm 0.6$	¹³² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$4.0 \pm 0.7 \pm 1.0$	¹³² EDWARDS	82E	CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{132,133} SCHARRE	80	MRK2 $e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1.78 \pm 0.21 \pm 0.33$	^{132,134,135} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.83 \pm 0.13 \pm 0.18$	^{132,136,137} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.66^{+0.17+0.24}_{-0.16-0.15}$	^{132,135,138} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
$1.03^{+0.21+0.26}_{-0.18-0.19}$	^{132,137,139} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹³⁰ Interference with the $J/\psi(1S)$ radiative transition to the broad $K \bar{K} \pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

¹³¹ Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

¹³² Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

¹³³ Corrected for spin-zero hypothesis for $\eta(1405)$.

¹³⁴ From fit to the $a_0(980)\pi 0^-+$ partial wave.


¹³⁵ $a_0(980)\pi$ mode.

¹³⁶ From fit to the $K^*(892)K 0^-+$ partial wave.

¹³⁷ $K^* K$ mode.

¹³⁸ From $a_0(980)\pi$ final state.

¹³⁹ From $K^*(890)K$ final state.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$ **Γ_{139} / Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
$1.07 \pm 0.17 \pm 0.11$	¹⁴⁰ BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$0.64 \pm 0.12 \pm 0.07$	¹⁴⁰ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹⁴⁰ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ **Γ_{140} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(3.0 ± 0.5) OUR AVERAGE				
$2.6 \pm 0.7 \pm 0.4$		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$3.38 \pm 0.33 \pm 0.64$		¹⁴¹ BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$7.0 \pm 0.6 \pm 1.1$	261	¹⁴² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹⁴¹ Via $a_0(980)\pi$.
¹⁴² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi) / \Gamma_{\text{total}}$ **Γ_{141} / Γ**

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.82	95	BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$ **Γ_{142}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.5 ± 0.8		OUR AVERAGE		
4.7 ± 0.3 ± 0.9		143 BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75 ± 1.05 ± 1.20		144 BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.09	90	145 BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$

¹⁴³ 4π mass less than 2.0 GeV.

¹⁴⁴ 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

¹⁴⁵ 4π mass in the range 2.0–25 GeV.

$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$ **Γ_{143}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.4	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$ **Γ_{144}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.8	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ **Γ_{145}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.16 ± 0.15		OUR AVERAGE		
Error includes scale factor of 1.1.				

4.86 ± 0.23 ± 0.08		¹⁴⁶ ABLIKIM	11	BES3 $J/\psi \rightarrow \eta'\gamma$
5.24 ± 0.12 ± 0.11		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
5.55 ± 0.44	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta'\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●


4.50 ± 0.14 ± 0.53		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
4.30 ± 0.31 ± 0.71		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
4.04 ± 0.16 ± 0.85	622	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
4.39 ± 0.09 ± 0.66	2420	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
4.1 ± 0.3 ± 0.6		BLOOM	83	CBAL $e^+e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ± 1.1	6	BRANDELIK	79C	DASP $e^+e^- \rightarrow 3\gamma$
2.4 ± 0.7	57	BARTEL	76	CNTR $e^+e^- \rightarrow 2\gamma\rho$

¹⁴⁶ ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.4 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ **Γ_{146}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8 ± 0.5	OUR AVERAGE		
Error includes scale factor of 1.9. See the ideogram below.			
4.32 ± 0.14 ± 0.73	¹⁴⁷ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	¹⁴⁸ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	¹⁴⁸ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	¹⁴⁹ BURKE	82	MRK2 e^+e^-

- 147 4π mass less than 3.0 GeV.
- 148 4π mass less than 2.0 GeV.
- 149 4π mass less than 2.5 GeV.



$\Gamma(\gamma f_2(1270) f_2(1270)) / \Gamma_{\text{total}}$ Γ_{147} / Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.5 \pm 0.7 \pm 1.6$	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant})) / \Gamma_{\text{total}}$ Γ_{148} / Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.2 \pm 0.8 \pm 1.7$	¹⁵⁰ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹⁵⁰ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{149} / Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.1 \pm 0.1 \pm 0.6$	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{150} / Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 0.5 \pm 0.5$	¹⁵¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹⁵¹ Assuming branching fraction $f_4(2050) \rightarrow \pi\pi / \text{total} = 0.167$.

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$					Γ_{151}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.61±0.33 OUR AVERAGE					
6.0 ±4.8 ±1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$	
1.41±0.2 ±0.42	120 ± 17	BISELLO	87 SPEC	e^+e^- , hadrons γ	
1.76±0.09±0.45		BALTRUSAIT..85C	MRK3	$e^+e^- \rightarrow$ hadrons γ	

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$					Γ_{152}/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
1.7 ±0.4 OUR AVERAGE Error includes scale factor of 1.3.					
2.1 ±0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$		
1.36±0.38	152,153 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$		

¹⁵² Estimated by us from various fits.

¹⁵³ Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$					Γ_{153}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.43±0.11 OUR AVERAGE					
1.62±0.26 ^{+0.02} _{-0.05}	154	ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
1.42±0.21 ^{+0.02} _{-0.04}	155	ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$	
1.33±0.05±0.20	156	AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1.36±0.09±0.23	156	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1.48±0.25±0.30	178	EDWARDS	82B CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$	
2.0 ±0.7	35	ALEXANDER	78 PLUT	e^+e^-	
1.2 ±0.6	30	157 BRANDELIK	78B DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	

¹⁵⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁵⁵ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁵⁶ Estimated using $B(f_2(1270) \rightarrow \pi\pi)=0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

¹⁵⁷ Restated by us to take account of spread of E1, M2, E3 transitions.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$					Γ_{154}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
(8.5±1.2-0.9) OUR AVERAGE Error includes scale factor of 1.2.					
9.62±0.29 ^{+3.51} _{-1.86}		158 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$	
5.0 ± 0.8 ^{+1.8} _{-0.4}		159,160 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$	
9.2 ± 1.4±1.4		160 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$	
10.4 ± 1.2±1.6		160 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	
9.6 ± 1.2±1.8		160 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.6 \pm 0.2^{+0.6}_{-0.2}$	160,161	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90	162 BISELLO	89B		$J/\psi \rightarrow 4\pi\gamma$
$1.6 \pm 0.4 \pm 0.3$		163 BALTRUSAIT..	87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.8 ± 1.6		164 EDWARDS	82D	CBAL	$e^+ e^- \rightarrow \eta \eta \gamma$

158 Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

159 Assuming $J^P = 2^+$ for $f_0(1710)$.

160 Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

161 Assuming $J^P = 0^+$ for $f_0(1710)$.

162 Includes unknown branching fraction to $\rho^0 \rho^0$.

163 Includes unknown branching fraction to $\pi^+ \pi^-$.

164 Includes unknown branching fraction to $\eta \eta$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
(4.0±1.0) OUR AVERAGE			
$3.96 \pm 0.06 \pm 1.12$	165 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$3.99 \pm 0.15 \pm 2.64$	165 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.5 \pm 1.6 \pm 0.8$	BAI	98H	BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
-----------------------	-----	-----	-----	---

165 Including unknown branching fraction to $\pi \pi$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma \eta) / \Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.104 ± 0.034 OUR AVERAGE				
$1.101 \pm 0.029 \pm 0.022$		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.88 \pm 0.08 \pm 0.11$	BLOOM	83	CBAL	$e^+ e^-$
0.82 ± 0.10	BRANDELIK	79C	DASP	$e^+ e^-$
1.3 ± 0.4	21 BARTEL	77	CNTR	$e^+ e^-$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.13 OUR AVERAGE			
$0.68 \pm 0.04 \pm 0.24$	BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.76 \pm 0.15 \pm 0.21$	166,167 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.87 \pm 0.14^{+0.14}_{-0.11}$	166 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

166 Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

167 From fit to the $K^*(892) K 1^{++}$ partial wave.

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$ **Γ_{159}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.61 ± 0.08 OUR AVERAGE			
0.69 ± 0.16 ± 0.20	168 BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\rho^0$
0.61 ± 0.04 ± 0.21	169 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 ± 0.09 ± 0.17	170 BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.625 ± 0.063 ± 0.103	171 BOLTON	92 MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.08 ± 0.16	172 BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

168 Assuming $B(f_1(1285) \rightarrow \rho^0\gamma) = 0.055 \pm 0.013$.

169 Assuming $\Gamma(f_1(1285) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.

170 Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.

171 Obtained summing the sequential decay channels

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi\pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}.$$

172 Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{160}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.5 ± 1.0 ± 0.7	BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ **Γ_{161}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(4.5+0.7-0.4) OUR AVERAGE					
3.85 ± 0.17 ^{+1.91} / _{-0.73}			173 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
3.6 ± 0.4 ^{+1.4} / _{-0.4}			173 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
5.6 ± 1.4 ± 0.9			173 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
4.5 ± 0.4 ± 0.9			173 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.8 ± 1.6 ± 1.4			173 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4 90 4 174 BRANDELIK 79C DASP $e^+e^- \rightarrow \pi^+\pi^-\gamma$

<2.3 90 3 ALEXANDER 78 PLUT $e^+e^- \rightarrow K^+K^-\gamma$

173 Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

174 Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ **Γ_{162}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.28 ± 0.05 ± 0.17	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ **Γ_{163}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.20 ± 0.04 ± 0.13	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.1 ± 0.2	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{165}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.3 ± 1.3	320	175 BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

175 Summed over all charges.

$\Gamma(\gamma \phi \phi)/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(4.0 ± 1.2) OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
7.5 ± 0.6 ± 1.2	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
3.4 ± 0.8 ± 0.6	33 ± 7	176 BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
3.1 ± 0.7 ± 0.4		176 BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

176 $\phi\phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.07 ± 0.07		49	EATON	84 MRK2	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	90	PERUZZI	78 MRK1	$e^+ e^-$
-------	----	---------	---------	-----------

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{168}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.33±0.05 OUR AVERAGE				
0.44±0.04±0.08	196 ± 19	177 ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
0.33±0.08±0.05		177 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
0.27±0.06±0.06		177 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
0.24 ^{+0.15} _{-0.10}	178,179	BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

177 Includes unknown branching fraction to $\phi\phi$.
 178 Estimated by us from various fits.
 179 Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{169}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13±0.09	180,181 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

180 Estimated by us from various fits.
 181 Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{170}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.98±0.08±0.32	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{171}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(2.6±0.4) OUR AVERAGE				
2.87±0.09 ^{+0.49} _{-0.52}	4265	182 ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
2.2 ± 0.4 ± 0.4	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

182 From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75^{+0.19}_{-0.09} OUR AVERAGE				
1.14 ^{+0.43+0.42} _{-0.30-0.26}	231	183 ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
0.70±0.04 ^{+0.19} _{-0.08}		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

183 From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

$\Gamma(\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}])/\Gamma_{\text{total}}$ Γ_{173}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58±0.03±0.20	184 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	185 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

184 For a broad structure around 1800 MeV.
 185 For a broad structure around 2040 MeV.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{174}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

(3.49+0.33-0.30) OUR AVERAGE

$3.63 \pm 0.36 \pm 0.13$		PEDLAR	09	CLE3 $J/\psi \rightarrow \pi^0 \gamma$
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E	BES2 $J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 1.1 \pm 0.7$		BLOOM	83	CBAL $e^+ e^-$
7.3 ± 4.7	10	BRANDELIK	79C	DASP $e^+ e^-$

$\Gamma(\gamma\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.79	90	EATON	84	MRK2 $e^+ e^-$
-----------------	----	-------	----	----------------

$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{176}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.13	90	HENRARD	87	DM2 $e^+ e^-$
-----------------	----	---------	----	---------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16	90	BAI	98G	BES $e^+ e^-$
-------	----	-----	-----	---------------

$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	¹⁸⁶ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-------------------------	----	---

¹⁸⁶ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

>250	99.9		¹⁸⁷ HASAN	96	SPEC $\bar{p}p \rightarrow \pi^+ \pi^-$
----------------	------	--	----------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			¹⁸⁸ BAI	96B	BES $e^+ e^- \rightarrow \gamma \bar{p}p, K\bar{K}$
------	--	--	--------------------	-----	---

< 2.3	95		¹⁸⁹ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-$
-------	----	--	-------------------------	----	---

< 1.6	95		¹⁸⁹ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-------	----	--	-------------------------	----	---

$12.4^{+6.4}_{-5.2} \pm 2.8$	23	¹⁸⁹ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
------------------------------	----	--------------------------------	------	---

$8.4^{+3.4}_{-2.8} \pm 1.6$	93	¹⁸⁹ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------------	----	--------------------------------	------	-------------------------------------

¹⁸⁷ Using BAI 96B.

¹⁸⁸ Using BARNES 93.

¹⁸⁹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{179}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.84 ± 0.26 ± 0.30	BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
---------------------------	-----	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4 \pm 0.8 \pm 0.4$	BAI	98H	BES $J/\psi \rightarrow \gamma\pi^0\pi^0$
-----------------------	-----	-----	---

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{180}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
< 3.6	¹⁹⁰ DEL-AMO-SA...100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 2.9	¹⁹⁰ DEL-AMO-SA...100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.6 ± 2.9 ± 2.4	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
10.8 ± 4.0 ± 3.2	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹⁹⁰ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{181}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.6 ± 0.5	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma \rho \bar{\rho}$

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{182}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
(1.01 ± 0.32) OUR AVERAGE			
1.00 ± 0.03 ± 0.45	¹⁹¹ ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.02 ± 0.09 ± 0.45	¹⁹¹ ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 5.7 ± 0.8 ^{192,193} BUGG 95 MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$

¹⁹¹ Including unknown branching fraction to $\pi\pi$.

¹⁹² Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

¹⁹³ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

$\Gamma(\gamma A \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{183}/Γ
(narrow state A with $m_A < 960$ MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.3	90	¹⁹⁴ INSLER	10	CLEO $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$

¹⁹⁴ The limit varies with mass m_A of a narrow state A and is 4.3×10^{-6} for $m_A = 0$ MeV, reaches its largest value of 6.3×10^{-6} at $m_A = 500$ MeV, and is 3.6×10^{-6} at $m_A = 960$ MeV.

WEAK DECAYS

$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{184}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	ABLIKIM	06M	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.1	90	ABLIKIM	06M	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6	90	¹⁹⁵ ABLIKIM	06M	BES2 $e^+e^- \rightarrow J/\psi$

¹⁹⁵ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$.

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{187}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-5}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{188}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{189}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				Γ_{190}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.5	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	¹⁹⁶ WICHT 08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
< 2.2	90	ABLIKIM 07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
<50	90	BARTEL 77	CNTR	$e^+ e^-$

¹⁹⁶WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.013 \times 10^{-3}$.

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$				Γ_{191}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI 03D	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{192}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{193}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.0	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$				Γ_{194}/Γ_7
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-2}$	90	ABLIKIM 08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$J/\psi(1S)$ REFERENCES

ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101R	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101R	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103R	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)

BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 R2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAITIS...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAITIS...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAITIS...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAITIS...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAITIS...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
BALTRUSAITIS...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34	1471.	
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI-...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)
