

CHARMED MESONS

($C = \pm 1$)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \text{ similarly for } D^{*'}s$$

$D^\pm$

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1869.62 \pm 0.15 \text{ MeV} \quad (S = 1.1)$$

$$\text{Mean life } \tau = (1040 \pm 7) \times 10^{-15} \text{ s}$$

$$c\tau = 311.8 \text{ } \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.096 \pm 0.004 \text{ } [a]$$

$$\Gamma(c \rightarrow D^{*}(2010)^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.1 \pm 1.0)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (1.0 \pm 1.3)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (0.3 \pm 0.9)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.1 \pm 1.3)\%$$

$$A_{CP}(\pi^\pm \pi^0) = (2.9 \pm 2.9)\%$$

$$A_{CP}(\pi^\pm \eta) = (1.0 \pm 1.5)\% \quad (S = 1.4)$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.5 \pm 1.2)\% \quad (S = 1.1)$$

$$A_{CP}(K_S^0 K^\pm) = (-0.23 \pm 0.31)\%$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.3 \pm 0.6)\%$$

$$A_{CP}(K^\pm K^{*0}) = (0.1 \pm 1.3)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.42 \pm 0.28)\%$$

$$A_{CP}(K^\pm K_0^{*}(1430)^0) = (8_{-6}^{+7})\%$$

$$A_{CP}(K^\pm K_2^{*}(1430)^0) = (43_{-26}^{+20})\%$$

$$A_{CP}(K^\pm K_0^{*}(800)) = (-12_{-13}^{+18})\%$$

$$A_{CP}(a_0(1450)^0 \pi^\pm) = (-19_{-16}^{+14})\%$$

$$A_{CP}(\phi(1680) \pi^\pm) = (-9 \pm 26)\%$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-2 \pm 4)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) = (-4 \pm 7)\%$$

$$A_{CP}(K^\pm \pi^0) = (-4 \pm 11)\%$$

T-violation decay-rate asymmetry

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-12 \pm 11) \times 10^{-3} \text{ } [b]$$

D^+ form factors

$$\begin{aligned}
f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= 0.707 \pm 0.013 \\
r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -1.7 \pm 0.5 \\
r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -14 \pm 11 \\
f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell &= 0.146 \pm 0.007 \\
r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -1.4 \pm 0.9 \\
r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -4 \pm 5 \\
f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= 0.086 \pm 0.006 \\
r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= -1.8 \pm 2.2 \\
r_v \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.51 \pm 0.07 \quad (S = 2.2) \\
r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.807 \pm 0.025 \\
r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.0 \pm 0.4 \\
\Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.13 \pm 0.08 \\
\Gamma_+/\Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.22 \pm 0.06 \quad (S = 1.6)
\end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as $\bar{K}^0$ modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 $\pm$ 0.30) %		—
μ^+ anything	(17.6 $\pm$ 3.2) %		—
K^- anything	(25.7 $\pm$ 1.4) %		—
$\bar{K}^0$ anything + K^0 anything	(61 $\pm$ 5) %		—
K^+ anything	(5.9 $\pm$ 0.8) %		—
$K^*(892)^-$ anything	(6 $\pm$ 5) %		—
$\bar{K}^*(892)^0$ anything	(23 $\pm$ 5) %		—
$K^*(892)^0$ anything	< 6.6 %	CL=90%	—
η anything	(6.3 $\pm$ 0.7) %		—
η' anything	(1.04 $\pm$ 0.18) %		—
ϕ anything	(1.03 $\pm$ 0.12) %		—
Leptonic and semileptonic modes			
$e^+ \nu_e$	< 8.8 $\times 10^{-6}$	CL=90%	935
$\mu^+ \nu_\mu$	(3.82 $\pm$ 0.33) $\times 10^{-4}$		932
$\tau^+ \nu_\tau$	< 1.2 $\times 10^{-3}$	CL=90%	91
$\bar{K}^0 e^+ \nu_e$	(8.83 $\pm$ 0.22) %		869
$\bar{K}^0 \mu^+ \nu_\mu$	(9.2 $\pm$ 0.6) %		865
$K^- \pi^+ e^+ \nu_e$	(4.00 $\pm$ 0.10) %		864

$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.68±0.10) %		722
$(K^- \pi^+)_{S-wave} e^+ \nu_e$	(2.32±0.10) × 10 ⁻³		—
$\bar{K}^*(1410)^0 e^+ \nu_e,$ $\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6 × 10 ⁻³	CL=90%	—
$\bar{K}_2^*(1430)^0 e^+ \nu_e,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5 × 10 ⁻⁴	CL=90%	—
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 × 10 ⁻³	CL=90%	864
$K^- \pi^+ \mu^+ \nu_\mu$	(3.8 ±0.4) %		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52±0.10) %		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(2.0 ±0.5) × 10 ⁻³		851
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.6 × 10 ⁻³	CL=90%	825
$\pi^0 e^+ \nu_e$	(4.05±0.18) × 10 ⁻³		930
$\eta e^+ \nu_e$	(1.14±0.10) × 10 ⁻³		855
$\rho^0 e^+ \nu_e$	(2.2 ±0.4) × 10 ⁻³		774
$\rho^0 \mu^+ \nu_\mu$	(2.4 ±0.4) × 10 ⁻³		770
$\omega e^+ \nu_e$	(1.6 ^{+0.7} _{-0.6}) × 10 ⁻³		771
$\eta'(958) e^+ \nu_e$	(2.2 ±0.5) × 10 ⁻⁴		689
$\phi e^+ \nu_e$	< 9 × 10 ⁻⁵	CL=90%	657

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\bar{K}^*(892)^0 e^+ \nu_e$	(5.52±0.15) %		722
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.28±0.15) %		717
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.4 × 10 ⁻⁴		380
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5 × 10 ⁻³		105

Hadronic modes with a $\bar{K}$ or $\bar{K}K\bar{K}$

$K_S^0 \pi^+$	(1.47±0.07) %	S=2.0	863
$K_L^0 \pi^+$	(1.46±0.05) %		863
$K^- 2\pi^+$	[c] (9.13±0.19) %		846
$(K^- \pi^+)_{S-wave} \pi^+$	(7.32±0.19) %		846
$\bar{K}_0^*(1430)^0 \pi^+,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] (1.21±0.06) %		382
$\bar{K}^*(892)^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.01±0.11) %		714
$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+$	not seen		381
$\bar{K}_2^*(1430)^0 \pi^+,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[d] (2.2 ±0.7) × 10 ⁻⁴		371
$\bar{K}^*(1680)^0 \pi^+,$ $\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] (2.1 ±1.1) × 10 ⁻⁴		58

$K^-(2\pi^+)_{I=2}$	(1.41±0.26) %	—
$K_S^0 \pi^+ \pi^0$	[c] (6.99±0.27) %	845
$K_S^0 \rho^+$	(4.8 ±1.0) %	677
$\bar{K}^*(892)^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	(1.3 ±0.6) %	714
$K_S^0 \pi^+ \pi^0$ nonresonant	(9 ±7) × 10 ⁻³	845
$K^- 2\pi^+ \pi^0$	[e] (5.99±0.18) %	816
$K_S^0 2\pi^+ \pi^-$	[e] (3.12±0.11) %	814
$K^- 3\pi^+ \pi^-$	[c] (5.6 ±0.5) × 10 ⁻³	S=1.1 772
$\bar{K}^*(892)^0 2\pi^+ \pi^-,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.2 ±0.4) × 10 ⁻³	645
$\bar{K}^*(892)^0 \rho^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(2.2 ±0.4) × 10 ⁻³	239
$\bar{K}^*(892)^0 a_1(1260)^+$	[f] (9.0 ±1.8) × 10 ⁻³	†
$K^- \rho^0 2\pi^+$	(1.68±0.27) × 10 ⁻³	524
$K^- 3\pi^+ \pi^-$ nonresonant	(3.9 ±2.9) × 10 ⁻⁴	772
$K^+ 2K_S^0$	(4.5 ±2.0) × 10 ⁻³	545
$K^+ K^- K_S^0 \pi^+$	(2.4 ±0.6) × 10 ⁻⁴	436

Pionic modes

$\pi^+ \pi^0$	(1.19±0.06) × 10 ⁻³	925
$2\pi^+ \pi^-$	(3.18±0.18) × 10 ⁻³	909
$\rho^0 \pi^+$	(8.1 ±1.5) × 10 ⁻⁴	767
$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	(1.78±0.16) × 10 ⁻³	909
$\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$	(1.34±0.12) × 10 ⁻³	—
$f_0(980) \pi^+,$ $f_0(980) \rightarrow \pi^+ \pi^-$	(1.52±0.33) × 10 ⁻⁴	669
$f_0(1370) \pi^+,$ $f_0(1370) \rightarrow \pi^+ \pi^-$	(8 ±4) × 10 ⁻⁵	—
$f_2(1270) \pi^+,$ $f_2(1270) \rightarrow \pi^+ \pi^-$	(4.9 ±0.9) × 10 ⁻⁴	485
$\rho(1450)^0 \pi^+,$ $\rho(1450)^0 \rightarrow \pi^+ \pi^-$	< 8 × 10 ⁻⁵	CL=95% 338
$f_0(1500) \pi^+,$ $f_0(1500) \rightarrow \pi^+ \pi^-$	(1.1 ±0.4) × 10 ⁻⁴	—
$f_0(1710) \pi^+,$ $f_0(1710) \rightarrow \pi^+ \pi^-$	< 5 × 10 ⁻⁵	CL=95% —
$f_0(1790) \pi^+,$ $f_0(1790) \rightarrow \pi^+ \pi^-$	< 6 × 10 ⁻⁵	CL=95% —
$(\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	< 1.2 × 10 ⁻⁴	CL=95% 909
$2\pi^+ \pi^-$ nonresonant	< 1.1 × 10 ⁻⁴	CL=95% 909
$\pi^+ 2\pi^0$	(4.6 ±0.4) × 10 ⁻³	910
$2\pi^+ \pi^- \pi^0$	(1.13±0.08) %	883

$\eta\pi^+, \eta \rightarrow \pi^+\pi^-\pi^0$	$(8.0 \pm 0.5) \times 10^{-4}$		848
$\omega\pi^+, \omega \rightarrow \pi^+\pi^-\pi^0$	$< 3 \times 10^{-4}$	CL=90%	763
$3\pi^+2\pi^-$	$(1.61 \pm 0.16) \times 10^{-3}$		845

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\eta\pi^+$	$(3.53 \pm 0.21) \times 10^{-3}$		848
$\eta\pi^+\pi^0$	$(1.38 \pm 0.35) \times 10^{-3}$		830
$\omega\pi^+$	$< 3.4 \times 10^{-4}$	CL=90%	764
$\eta'(958)\pi^+$	$(4.67 \pm 0.29) \times 10^{-3}$		681
$\eta'(958)\pi^+\pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$		654

Hadronic modes with a $K\bar{K}$ pair

$K^+K_S^0$	$(2.83 \pm 0.16) \times 10^{-3}$	S=2.2	793
$K^+K^-\pi^+$	[c] $(9.54 \pm 0.26) \times 10^{-3}$	S=1.1	744
$\phi\pi^+, \phi \rightarrow K^+K^-$	$(2.65^{+0.08}_{-0.09}) \times 10^{-3}$		647
$K^+\bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(2.45^{+0.09}_{-0.14}) \times 10^{-3}$		613
$K^+\bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^-\pi^+$	$(1.79 \pm 0.34) \times 10^{-3}$		—
$K^+\bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^-\pi^+$	$(1.6^{+1.2}_{-0.8}) \times 10^{-4}$		—
$K^+\bar{K}_0^*(800), \bar{K}_0^* \rightarrow K^-\pi^+$	$(6.7^{+3.4}_{-2.1}) \times 10^{-4}$		—
$a_0(1450)^0\pi^+, a_0^0 \rightarrow K^+K^-$	$(4.4^{+7.0}_{-1.8}) \times 10^{-4}$		—
$\phi(1680)\pi^+, \phi \rightarrow K^+K^-$	$(4.9^{+4.0}_{-1.9}) \times 10^{-5}$		—
$K^+K^-\pi^+$ nonresonant	not seen		744
$K^+K_S^0\pi^+\pi^-$	$(1.75 \pm 0.18) \times 10^{-3}$		678
$K_S^0K^-\pi^+\pi^-$	$(2.40 \pm 0.18) \times 10^{-3}$		678
$K^+K^-\pi^+\pi^-$	$(2.2 \pm 1.2) \times 10^{-4}$		600

A few poorly measured branching fractions:

$\phi\pi^+\pi^0$	$(2.3 \pm 1.0) \%$		619
$\phi\rho^+$	$< 1.5 \%$	CL=90%	260
$K^+K^-\pi^+\pi^0$ non- ϕ	$(1.5^{+0.7}_{-0.6}) \%$		682
$K^*(892)^+K_S^0$	$(1.6 \pm 0.7) \%$		612

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(1.83 \pm 0.26) \times 10^{-4}$	S=1.4	864
$K^+ \eta$	$(1.08 \pm 0.17) \times 10^{-4}$		776
$K^+ \eta'(958)$	$(1.76 \pm 0.22) \times 10^{-4}$		571
$K^+ \pi^+ \pi^-$	$(5.27 \pm 0.23) \times 10^{-4}$		846
$K^+ \rho^0$	$(2.0 \pm 0.5) \times 10^{-4}$		679
$K^{*}(892)^0 \pi^+, K^{*}(892)^0 \rightarrow$	$(2.5 \pm 0.4) \times 10^{-4}$		714
$K^+ \pi^-$			
$K^+ f_0(980), f_0(980) \rightarrow$	$(4.7 \pm 2.8) \times 10^{-5}$		—
$\pi^+ \pi^-$			
$K_2^{*}(1430)^0 \pi^+, K_2^{*}(1430)^0 \rightarrow$	$(4.2 \pm 2.9) \times 10^{-5}$		—
$K^+ \pi^-$			
$K^+ \pi^+ \pi^-$ nonresonant	not seen		846
$2K^+ K^-$	$(8.7 \pm 2.0) \times 10^{-5}$		550

 **$\Delta C = 1$ weak neutral current ($C1$) modes, or
Lepton Family number (LF) or Lepton number (L) violating modes**

$\pi^+ e^+ e^-$	$C1$	< 1.1	$\times 10^{-6}$	CL=90%	930
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[g]	$(1.7^{+1.4}_{-0.9}) \times 10^{-6}$			—
$\pi^+ \mu^+ \mu^-$	$C1$	< 3.9	$\times 10^{-6}$	CL=90%	918
$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[g]	$(1.8 \pm 0.8) \times 10^{-6}$			—
$\rho^+ \mu^+ \mu^-$	$C1$	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^+ e^+ e^-$	[h]	< 1.0	$\times 10^{-6}$	CL=90%	870
$K^+ \mu^+ \mu^-$	[h]	< 4.3	$\times 10^{-6}$	CL=90%	856
$\pi^+ e^+ \mu^-$	LF	< 2.9	$\times 10^{-6}$	CL=90%	927
$\pi^+ e^- \mu^+$	LF	< 3.6	$\times 10^{-6}$	CL=90%	927
$K^+ e^+ \mu^-$	LF	< 1.2	$\times 10^{-6}$	CL=90%	866
$K^+ e^- \mu^+$	LF	< 2.8	$\times 10^{-6}$	CL=90%	866
$\pi^- 2e^+$	L	< 1.1	$\times 10^{-6}$	CL=90%	930
$\pi^- 2\mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%	918
$\pi^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%	927
$\rho^- 2\mu^+$	L	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^- 2e^+$	L	< 9	$\times 10^{-7}$	CL=90%	870
$K^- 2\mu^+$	L	< 1.0	$\times 10^{-5}$	CL=90%	856
$K^- e^+ \mu^+$	L	< 1.9	$\times 10^{-6}$	CL=90%	866
$K^{*}(892)^- 2\mu^+$	L	< 8.5	$\times 10^{-4}$	CL=90%	703



$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1864.86 \pm 0.13 \text{ MeV}$$

$$m_{D^\pm} - m_{D^0} = 4.76 \pm 0.10 \text{ MeV} \quad (S = 1.1)$$

$$\text{Mean life } \tau = (410.1 \pm 1.5) \times 10^{-15} \text{ s}$$

$$c\tau = 122.9 \mu\text{m}$$

$$|m_{D_1^0} - m_{D_2^0}| = (1.18^{+0.43}_{-0.47}) \times 10^{10} \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2\gamma = (1.43 \pm 0.19) \times 10^{-2}$$

$$|q/p| = 0.67^{+0.18}_{-0.14}$$

$$A_\Gamma = (-0.22 \pm 1.61) \times 10^{-3}$$

$$K^+ \pi^- \text{ relative strong phase: } \cos \delta = 0.81^{+0.23}_{-0.19}$$

$$K^- \pi^+ \pi^0 \text{ coherence factor } R_{K \pi \pi^0} = 0.78^{+0.11}_{-0.25}$$

$$K^- \pi^+ \pi^0 \text{ average relative strong phase } \delta^{K \pi \pi^0} = (239^{+32}_{-28})^\circ$$

$$K^- \pi^- 2\pi^+ \text{ coherence factor } R_{K 3\pi} = 0.36^{+0.24}_{-0.30}$$

$$K^- \pi^- 2\pi^+ \text{ average relative strong phase } \delta^{K 3\pi} = (118^{+60}_{-50})^\circ$$

$$K_S^0 K^+ \pi^- \text{ coherence factor } R_{K_S^0 K \pi} = 0.73 \pm 0.08$$

$$K_S^0 K^+ \pi^- \text{ average relative strong phase } \delta^{K_S^0 K \pi} = (8 \pm 15)^\circ$$

$$K^* K \text{ coherence factor } R_{K^* K} = 1.00 \pm 0.16$$

$$K^* K \text{ average relative strong phase } \delta^{K^* K} = (26 \pm 16)^\circ$$

CP-violation decay-rate asymmetries (labeled by the D^0 decay)

$$A_{CP}(K^+ K^-) = (-0.21 \pm 0.17)\%$$

$$A_{CP}(2K_S^0) = (-23 \pm 19)\%$$

$$A_{CP}(\pi^+ \pi^-) = (0.22 \pm 0.21)\%$$

$$A_{CP}(2\pi^0) = (0 \pm 5)\%$$

$$A_{CP}(\pi^+ \pi^- \pi^0) = (0.3 \pm 0.4)\%$$

$$A_{CP}(\rho(770)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (1.2 \pm 0.9)\% [i]$$

$$A_{CP}(\rho(770)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-3.1 \pm 3.0)\% [i]$$

$$A_{CP}(\rho(770)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (-1.0 \pm 1.7)\% [i]$$

$$A_{CP}(\rho(1450)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 70)\% [i]$$

$$A_{CP}(\rho(1450)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-20 \pm 40)\% [i]$$

$$A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 9)\% [i]$$

$$A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (-5 \pm 14)\% [i]$$

$$A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (13 \pm 9)\% [i]$$

$$A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (8 \pm 11)\% [i]$$

$$A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 35)\% [i]$$

$$A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (25 \pm 18)\% [i]$$

$$A_{CP}(f_0(1500) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 18)\% [i]$$

$$A_{CP}(f_0(1710) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 24)\% [i]$$

$$A_{CP}(f_2(1270) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-4 \pm 6)\% [i]$$

$$\begin{aligned}
 A_{CP}(\sigma(400)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (6 \pm 8)\% [i] \\
 A_{CP}(\text{nonresonant } \pi^+\pi^-\pi^0) &= (-13 \pm 23)\% [i] \\
 A_{CP}(K^+K^-\pi^0) &= (-1.0 \pm 1.7)\% \\
 A_{CP}(K^*(892)^+K^- \rightarrow K^+K^-\pi^0) &= (-0.9 \pm 1.3)\% [i] \\
 A_{CP}(K^*(1410)^+K^- \rightarrow K^+K^-\pi^0) &= (-21 \pm 24)\% [i] \\
 A_{CP}((K^+\pi^0)_{S\text{-wave}}K^- \rightarrow K^+K^-\pi^0) &= (7 \pm 15)\% [i] \\
 A_{CP}(\phi(1020)\pi^0 \rightarrow K^+K^-\pi^0) &= (1.1 \pm 2.2)\% [i] \\
 A_{CP}(f_0(980)\pi^0 \rightarrow K^+K^-\pi^0) &= (-3 \pm 19)\% [i] \\
 A_{CP}(a_0(980)^0\pi^0 \rightarrow K^+K^-\pi^0) &= (-5 \pm 16)\% [i] \\
 A_{CP}(f'_2(1525)\pi^0 \rightarrow K^+K^-\pi^0) &= (0 \pm 160)\% [i] \\
 A_{CP}(K^*(892)^-K^+ \rightarrow K^+K^-\pi^0) &= (-5 \pm 4)\% [i] \\
 A_{CP}(K^*(1410)^-K^+ \rightarrow K^+K^-\pi^0) &= (-17 \pm 29)\% [i] \\
 A_{CP}((K^-\pi^0)_{S\text{-wave}}K^+ \rightarrow K^+K^-\pi^0) &= (-10 \pm 40)\% [i] \\
 A_{CP}(K_S^0\pi^0) &= (-0.27 \pm 0.21)\% \\
 A_{CP}(K_S^0\eta) &= (0.5 \pm 0.5)\% \\
 A_{CP}(K_S^0\eta') &= (1.0 \pm 0.7)\% \\
 A_{CP}(K_S^0\phi) &= (-3 \pm 9)\% \\
 A_{CP}(K^-\pi^+) &= (0.1 \pm 0.7)\% \\
 A_{CP}(K^+\pi^-) &= (2.2 \pm 3.2)\% \\
 A_{CP}(K^-\pi^+\pi^0) &= (0.2 \pm 0.9)\% \\
 A_{CP}(K^+\pi^-\pi^0) &= (0 \pm 5)\% \\
 A_{CP}(K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.8)\% \\
 A_{CP}(K^*(892)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (0.4 \pm 0.5)\% \\
 A_{CP}(K^*(892)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (1 \pm 6)\% \\
 A_{CP}(\bar{K}^0\rho^0 \rightarrow K_S^0\pi^+\pi^-) &= (-0.1 \pm 0.5)\% \\
 A_{CP}(\bar{K}^0\omega \rightarrow K_S^0\pi^+\pi^-) &= (-13 \pm 7)\% \\
 A_{CP}(\bar{K}^0f_0(980) \rightarrow K_S^0\pi^+\pi^-) &= (-0.4 \pm 2.7)\% \\
 A_{CP}(\bar{K}^0f_2(1270) \rightarrow K_S^0\pi^+\pi^-) &= (-4 \pm 5)\% \\
 A_{CP}(\bar{K}^0f_0(1370) \rightarrow K_S^0\pi^+\pi^-) &= (-1 \pm 9)\% \\
 A_{CP}(\bar{K}^0\rho^0(1450) \rightarrow K_S^0\pi^+\pi^-) &= (-4 \pm 10)\% \\
 A_{CP}(\bar{K}^0f_0(600) \rightarrow K_S^0\pi^+\pi^-) &= (-3 \pm 5)\% \\
 A_{CP}(\bar{K}^0f_2(1270) \rightarrow K_S^0\pi^+\pi^-) &= (-7 \pm 8)\% \\
 A_{CP}(K^*(1410)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (-2 \pm 9)\% \\
 A_{CP}(K_0^*(1430)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (4 \pm 4)\% \\
 A_{CP}(K_0^*(1430)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (12 \pm 15)\% \\
 A_{CP}(K_2^*(1430)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (3 \pm 6)\% \\
 A_{CP}(K_2^*(1430)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (-10 \pm 32)\% \\
 A_{CP}(K^*(1680)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (-10 \pm 32)\% \\
 A_{CP}(K^-\pi^+\pi^+\pi^-) &= (0.7 \pm 1.0)\% \\
 A_{CP}(K^+\pi^-\pi^+\pi^-) &= (-2 \pm 4)\% \\
 A_{CP}(K^+K^-\pi^+\pi^-) &= (-8 \pm 7)\% \\
 A_{CP}(K_1^*(1270)^+K^- \rightarrow K^{*0}\pi^+K^-) &= (-1 \pm 10)\%
 \end{aligned}$$

$$A_{CP}(K_1^*(1270)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) = (-10 \pm 32)\%$$

$$A_{CP}(K_1^*(1270)^+ K^- \rightarrow \rho^0 K^+ K^-) = (-7 \pm 17)\%$$

$$A_{CP}(K_1^*(1270)^- K^+ \rightarrow \rho^0 K^- K^+) = (10 \pm 13)\%$$

$$A_{CP}(K^*(1410)^+ K^- \rightarrow K^{*0} \pi^+ K^-) = (-20 \pm 17)\%$$

$$A_{CP}(K^*(1410)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) = (-1 \pm 14)\%$$

$$A_{CP}(K^{*0} \bar{K}^{*0} \text{ S-wave}) = (10 \pm 14)\%$$

$$A_{CP}(\phi \rho^0 \text{ S-wave}) = (-3 \pm 5)\%$$

$$A_{CP}(\phi \rho^0 \text{ D-wave}) = (-37 \pm 19)\%$$

$$A_{CP}(\phi(\pi^+ \pi^-)_{S\text{-wave}}) = (-9 \pm 10)\%$$

$$A_{CP}((K^- \pi^+)_{P\text{-wave}} (K^+ \pi^-)_{S\text{-wave}}) = (3 \pm 11)\%$$

CP-violation asymmetry difference

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.68 \pm 0.16)\%$$

T-violation decay-rate asymmetry

$$A_T(K^+ K^- \pi^+ \pi^-) = (1 \pm 7) \times 10^{-3} [b]$$

CPT-violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$r_V \equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 1.7 \pm 0.8$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 0.9 \pm 0.4$$

$$f_+(0) \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.727 \pm 0.011$$

$$f_+(0)|V_{cs}| \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.726 \pm 0.009$$

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = -2.65 \pm 0.35$$

$$r_2 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 13 \pm 9$$

$$f_+(0)|V_{cd}| \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.152 \pm 0.005$$

$$r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -2.8 \pm 0.5$$

$$r_2 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 6 \pm 3.0$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as $\bar{K}^0$ modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Topological modes			
0-prongs	[j] (15 $\pm$ 6) %		—
2-prongs	(70 $\pm$ 6) %		—
4-prongs	[k] (14.5 $\pm$ 0.5) %		—
6-prongs	[l] (6.4 $\pm$ 1.3) $\times 10^{-4}$		—
Inclusive modes			
e^+ anything	[n] (6.49 $\pm$ 0.11) %		—
μ^+ anything	(6.7 $\pm$ 0.6) %		—
K^- anything	(54.7 $\pm$ 2.8) %	S=1.3	—
$\bar{K}^0$ anything + K^0 anything	(47 $\pm$ 4) %		—
K^+ anything	(3.4 $\pm$ 0.4) %		—
$K^*(892)^-$ anything	(15 $\pm$ 9) %		—
$\bar{K}^*(892)^0$ anything	(9 $\pm$ 4) %		—
$K^*(892)^+$ anything	< 3.6 %	CL=90%	—
$K^*(892)^0$ anything	(2.8 $\pm$ 1.3) %		—
η anything	(9.5 $\pm$ 0.9) %		—
η' anything	(2.48 $\pm$ 0.27) %		—
ϕ anything	(1.05 $\pm$ 0.11) %		—
Semileptonic modes			
$K^- e^+ \nu_e$	(3.55 $\pm$ 0.05) %	S=1.2	867
$K^- \mu^+ \nu_\mu$	(3.31 $\pm$ 0.13) %		864
$K^*(892)^- e^+ \nu_e$	(2.16 $\pm$ 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$	(1.91 $\pm$ 0.24) %		714
$K^- \pi^0 e^+ \nu_e$	(1.6 $\pm$ 1.3 $\pm$ 0.5) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	(2.7 $\pm$ 0.9 $\pm$ 0.7) %		860
$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 $\pm$ 1.4 $\pm$ 1.1) $\times 10^{-4}$		843
$K_1(1270)^- e^+ \nu_e$	(7.6 $\pm$ 4.0 $\pm$ 3.1) $\times 10^{-4}$		498
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2 $\times 10^{-3}$	CL=90%	821
$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.4 $\times 10^{-3}$	CL=90%	692
$\pi^- e^+ \nu_e$	(2.89 $\pm$ 0.08) $\times 10^{-3}$	S=1.1	927
$\pi^- \mu^+ \nu_\mu$	(2.37 $\pm$ 0.24) $\times 10^{-3}$		924
$\rho^- e^+ \nu_e$	(1.9 $\pm$ 0.4) $\times 10^{-3}$		771

Hadronic modes with one $\bar{K}$

$K^- \pi^+$	(3.88 $\pm$ 0.05) %	S=1.1	861
$K^+ \pi^-$	(1.37 $\pm$ 0.06) $\times 10^{-4}$		861
$K_S^0 \pi^0$	(1.19 $\pm$ 0.04) %		860
$K_L^0 \pi^0$	(10.0 $\pm$ 0.7) $\times 10^{-3}$		860
$K_S^0 \pi^+ \pi^-$	[c] (2.83 $\pm$ 0.20) %	S=1.1	842
$K_S^0 \rho^0$	(6.3 $\pm$ 0.7 $\pm$ 0.8) $\times 10^{-3}$		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	(2.1 $\pm$ 0.6) $\times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S\text{-wave}}$	(3.4 $\pm$ 0.8) $\times 10^{-3}$		842
$K_S^0 f_0(980),$ $f_0(980) \rightarrow \pi^+ \pi^-$	(1.22 $\pm$ 0.40 $\pm$ 0.24) $\times 10^{-3}$		549
$K_S^0 f_0(1370),$ $f_0(1370) \rightarrow \pi^+ \pi^-$	(2.8 $\pm$ 0.9 $\pm$ 1.3) $\times 10^{-3}$		†
$K_S^0 f_2(1270),$ $f_2(1270) \rightarrow \pi^+ \pi^-$	(9 $\pm$ 10 $\pm$ 6) $\times 10^{-5}$		262
$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K_S^0 \pi^-$	(1.66 $\pm$ 0.15 $\pm$ 0.17) %		711
$K_0^*(1430)^- \pi^+,$ $K_0^*(1430)^- \rightarrow K_S^0 \pi^-$	(2.70 $\pm$ 0.40 $\pm$ 0.34) $\times 10^{-3}$		378
$K_2^*(1430)^- \pi^+,$ $K_2^*(1430)^- \rightarrow K_S^0 \pi^-$	(3.4 $\pm$ 1.9 $\pm$ 1.0) $\times 10^{-4}$		367
$K^*(1680)^- \pi^+,$ $K^*(1680)^- \rightarrow K_S^0 \pi^-$	(4 $\pm$ 4) $\times 10^{-4}$		46
$K^*(892)^+ \pi^-,$ $K^*(892)^+ \rightarrow K_S^0 \pi^+$	[o] (1.14 $\pm$ 0.60 $\pm$ 0.34) $\times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-,$ $K_0^*(1430)^+ \rightarrow K_S^0 \pi^+$	[o] < 1.4 $\times 10^{-5}$	CL=95%	—
$K_2^*(1430)^+ \pi^-,$ $K_2^*(1430)^+ \rightarrow K_S^0 \pi^+$	[o] < 3.4 $\times 10^{-5}$	CL=95%	—
$K_S^0 \pi^+ \pi^-$ nonresonant	(2.5 $\pm$ 6.0 $\pm$ 1.6) $\times 10^{-4}$		842
$K^- \pi^+ \pi^0$	[c] (13.9 $\pm$ 0.5) %	S=1.7	844
$K^- \rho^+$	(10.8 $\pm$ 0.7) %		675
$K^- \rho(1700)^+,$ $\rho(1700)^+ \rightarrow \pi^+ \pi^0$	(7.9 $\pm$ 1.7) $\times 10^{-3}$		†
$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K^- \pi^0$	(2.22 $\pm$ 0.40 $\pm$ 0.19) %		711
$\bar{K}^*(892)^0 \pi^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.88 $\pm$ 0.23) %		711

$K_0^*(1430)^- \pi^+$,	$(4.6 \pm 2.1) \times 10^{-3}$		378
$K_0^*(1430)^- \rightarrow K^- \pi^0$			
$\bar{K}_0^*(1430)^0 \pi^0$,	$(5.7 \pm 5.0 \pm 1.5) \times 10^{-3}$		379
$\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$			
$K^*(1680)^- \pi^+$,	$(1.8 \pm 0.7) \times 10^{-3}$		46
$K^*(1680)^- \rightarrow K^- \pi^0$			
$K^- \pi^+ \pi^0$ nonresonant	$(1.11 \pm 0.50 \pm 0.19) \%$		844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2	843
$K_S^0(2\pi^0)$ -S-wave	$(2.6 \pm 0.7) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \pi^0$,	$(7.8 \pm 0.7) \times 10^{-3}$		711
$\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$			
$\bar{K}^*(1430)^0 \pi^0, \bar{K}^{*0} \rightarrow$	$(4 \pm 23) \times 10^{-5}$		—
$K_S^0 \pi^0$			
$\bar{K}^*(1680)^0 \pi^0, \bar{K}^{*0} \rightarrow$	$(1.0 \pm 0.4) \times 10^{-3}$		—
$K_S^0 \pi^0$			
$K_S^0 f_2(1270), f_2 \rightarrow 2\pi^0$	$(2.3 \pm 1.1) \times 10^{-4}$		—
$2K_S^0$, one $K_S^0 \rightarrow 2\pi^0$	$(3.2 \pm 1.1) \times 10^{-4}$		—
$K^- 2\pi^+ \pi^-$	[c] $(8.08 \pm 0.21 \pm 0.19) \%$	S=1.3	813
$K^- \pi^+ \rho^0$ total	$(6.75 \pm 0.33) \%$		609
$K^- \pi^+ \rho^0$ 3-body	$(5.1 \pm 2.3) \times 10^{-3}$		609
$\bar{K}^*(892)^0 \rho^0$,	$(1.05 \pm 0.23) \%$		416
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$K^- a_1(1260)^+$,	$(3.6 \pm 0.6) \%$		327
$a_1(1260)^+ \rightarrow 2\pi^+ \pi^-$			
$\bar{K}^*(892)^0 \pi^+ \pi^-$ total,	$(1.6 \pm 0.4) \%$		685
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body,	$(9.9 \pm 2.3) \times 10^{-3}$		685
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$K_1(1270)^- \pi^+$,	[p] $(2.9 \pm 0.3) \times 10^{-3}$		484
$K_1(1270)^- \rightarrow K^- \pi^+ \pi^-$			
$K^- 2\pi^+ \pi^-$ nonresonant	$(1.88 \pm 0.26) \%$		813
$K_S^0 \pi^+ \pi^- \pi^0$	[q] $(5.2 \pm 0.6) \%$		813
$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	$(1.02 \pm 0.09) \times 10^{-3}$		772
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	$(9.9 \pm 0.5) \times 10^{-3}$		670
$K^- 2\pi^+ \pi^- \pi^0$	$(4.2 \pm 0.4) \%$		771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$,	$(1.3 \pm 0.6) \%$		643
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$			
$K^- \pi^+ \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	$(2.7 \pm 0.5) \%$		605
$\bar{K}^*(892)^0 \omega$,	$(6.5 \pm 3.0) \times 10^{-3}$		410
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$,			
$\omega \rightarrow \pi^+ \pi^- \pi^0$			
$K_S^0 \eta \pi^0$	$(5.5 \pm 1.1) \times 10^{-3}$		721

$K_S^0 a_0(980), a_0(980) \rightarrow \eta \pi^0$	$(6.5 \pm 2.0) \times 10^{-3}$	—
$\overline{K}^{*0}(892) \eta,$	$(1.6 \pm 0.5) \times 10^{-3}$	—
$\overline{K}^{*0}(892) \rightarrow K_S^0 \pi$		
$K_S^0 2\pi^+ 2\pi^-$	$(2.69 \pm 0.31) \times 10^{-3}$	768
$K_S^0 \rho^0 \pi^+ \pi^-, \text{ no } K^*(892)^-$	$(1.1 \pm 0.7) \times 10^{-3}$	—
$K^*(892)^- 2\pi^+ \pi^-,$	$(5 \pm 8) \times 10^{-4}$	642
$K^*(892)^- \rightarrow K_S^0 \pi^-,$		
no ρ^0		
$K^*(892)^- \rho^0 \pi^+,$	$(1.6 \pm 0.6) \times 10^{-3}$	230
$K^*(892)^- \rightarrow K_S^0 \pi^-$		
$K_S^0 2\pi^+ 2\pi^- \text{ nonresonant}$	$< 1.2 \times 10^{-3}$	CL=90% 768
$K^- 3\pi^+ 2\pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	713

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\overline{K}^*(892)\rho$ submodes only appear below.)

$K_S^0 \eta$	$(4.79 \pm 0.30) \times 10^{-3}$	772
$K_S^0 \omega$	$(1.11 \pm 0.06) \%$	670
$K_S^0 \eta'(958)$	$(9.4 \pm 0.5) \times 10^{-3}$	565
$K^- a_1(1260)^+$	$(7.8 \pm 1.1) \%$	327
$K^- a_2(1320)^+$	$< 2 \times 10^{-3}$	CL=90% 198
$\overline{K}^{*0}(892) \pi^+ \pi^- \text{ total}$	$(2.4 \pm 0.5) \%$	685
$\overline{K}^{*0}(892) \pi^+ \pi^- 3\text{-body}$	$(1.48 \pm 0.34) \%$	685
$\overline{K}^{*0}(892) \rho^0$	$(1.58 \pm 0.34) \%$	417
$\overline{K}^{*0}(892) \rho^0 \text{ transverse}$	$(1.7 \pm 0.6) \%$	417
$\overline{K}^{*0}(892) \rho^0 S\text{-wave}$	$(3.0 \pm 0.6) \%$	417
$\overline{K}^{*0}(892) \rho^0 S\text{-wave long.}$	$< 3 \times 10^{-3}$	CL=90% 417
$\overline{K}^{*0}(892) \rho^0 P\text{-wave}$	$< 3 \times 10^{-3}$	CL=90% 417
$\overline{K}^{*0}(892) \rho^0 D\text{-wave}$	$(2.1 \pm 0.6) \%$	417
$K_1(1270)^- \pi^+$	[p] $(1.6 \pm 0.8) \%$	484
$K_1(1400)^- \pi^+$	$< 1.2 \%$	CL=90% 386
$\overline{K}^{*0}(892) \pi^+ \pi^- \pi^0$	$(1.9 \pm 0.9) \%$	644
$K^- \pi^+ \omega$	$(3.0 \pm 0.6) \%$	605
$\overline{K}^{*0}(892) \omega$	$(1.1 \pm 0.5) \%$	410
$K^- \pi^+ \eta'(958)$	$(7.5 \pm 1.9) \times 10^{-3}$	479
$\overline{K}^{*0}(892) \eta'(958)$	$< 1.1 \times 10^{-3}$	CL=90% 120

Hadronic modes with three K's

$K_S^0 K^+ K^-$	$(4.47 \pm 0.34) \times 10^{-3}$	544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	$(3.0 \pm 0.4) \times 10^{-3}$	—
$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$	$(6.0 \pm 1.8) \times 10^{-4}$	—
$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$	$< 1.1 \times 10^{-4}$	CL=95% —
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	$< 9 \times 10^{-5}$	CL=95% —

$K_S^0 \phi, \phi \rightarrow K^+ K^-$	$(2.05 \pm 0.16) \times 10^{-3}$	520
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	$(1.7 \pm 1.1) \times 10^{-4}$	—
$3K_S^0$	$(9.1 \pm 1.3) \times 10^{-4}$	539
$K^+ 2K^- \pi^+$	$(2.21 \pm 0.31) \times 10^{-4}$	434
$K^+ K^- \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(4.4 \pm 1.7) \times 10^{-5}$	†
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.7) \times 10^{-5}$	422
$\phi \bar{K}^*(892)^0,$ $\phi \rightarrow K^+ K^-,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.06 \pm 0.20) \times 10^{-4}$	†
$K^+ 2K^- \pi^+$ nonresonant	$(3.3 \pm 1.5) \times 10^{-5}$	434
$2K_S^0 K^\pm \pi^\mp$	$(6.0 \pm 1.3) \times 10^{-4}$	427

Pionic modes

$\pi^+ \pi^-$	$(1.402 \pm 0.026) \times 10^{-3}$	S=1.1	922
$2\pi^0$	$(8.20 \pm 0.35) \times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	$(1.43 \pm 0.06) \%$	S=1.9	907
$\rho^+ \pi^-$	$(9.8 \pm 0.4) \times 10^{-3}$		764
$\rho^0 \pi^0$	$(3.72 \pm 0.22) \times 10^{-3}$		764
$\rho^- \pi^+$	$(4.96 \pm 0.24) \times 10^{-3}$		764
$\rho(1450)^+ \pi^-, \rho(1450)^+ \rightarrow$ $\pi^+ \pi^0$	$(1.6 \pm 2.0) \times 10^{-5}$		—
$\rho(1450)^0 \pi^0, \rho(1450)^0 \rightarrow$ $\pi^+ \pi^-$	$(4.3 \pm 1.9) \times 10^{-5}$		—
$\rho(1450)^- \pi^+, \rho(1450)^- \rightarrow$ $\pi^- \pi^0$	$(2.6 \pm 0.4) \times 10^{-4}$		—
$\rho(1700)^+ \pi^-, \rho(1700)^+ \rightarrow$ $\pi^+ \pi^0$	$(5.9 \pm 1.4) \times 10^{-4}$		—
$\rho(1700)^0 \pi^0, \rho(1700)^0 \rightarrow$ $\pi^+ \pi^-$	$(7.2 \pm 1.7) \times 10^{-4}$		—
$\rho(1700)^- \pi^+, \rho(1700)^- \rightarrow$ $\pi^- \pi^0$	$(4.6 \pm 1.1) \times 10^{-4}$		—
$f_0(980) \pi^0, f_0(980) \rightarrow$ $\pi^+ \pi^-$	$(3.6 \pm 0.8) \times 10^{-5}$		—
$f_0(500) \pi^0, f_0(500) \rightarrow$ $\pi^+ \pi^-$	$(1.18 \pm 0.21) \times 10^{-4}$		—
$f_0(1370) \pi^0, f_0(1370) \rightarrow$ $\pi^+ \pi^-$	$(5.3 \pm 2.1) \times 10^{-5}$		—
$f_0(1500) \pi^0, f_0(1500) \rightarrow$ $\pi^+ \pi^-$	$(5.6 \pm 1.5) \times 10^{-5}$		—
$f_0(1710) \pi^0, f_0(1710) \rightarrow$ $\pi^+ \pi^-$	$(4.4 \pm 1.5) \times 10^{-5}$		—
$f_2(1270) \pi^0, f_2(1270) \rightarrow$ $\pi^+ \pi^-$	$(1.89 \pm 0.20) \times 10^{-4}$		—
$\pi^+ \pi^- \pi^0$ nonresonant	$(1.20 \pm 0.35) \times 10^{-4}$		907
$3\pi^0$	$< 3.5 \times 10^{-4}$	CL=90%	908
$2\pi^+ 2\pi^-$	$(7.42 \pm 0.21) \times 10^{-3}$	S=1.1	880

$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	$(4.45 \pm 0.31) \times 10^{-3}$	—
$2\pi^+ \pi^-$ total		
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	$(3.21 \pm 0.25) \times 10^{-3}$	—
$\rho^0 \pi^+$ S-wave		
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	$(1.9 \pm 0.5) \times 10^{-4}$	—
$\rho^0 \pi^+$ D-wave		
$a_1(1260)^+ \pi^-, a_1^+ \rightarrow$	$(6.2 \pm 0.7) \times 10^{-4}$	—
$\sigma \pi^+$		
$2\rho^0$ total	$(1.82 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.2 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.8 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.25 \pm 0.10) \times 10^{-3}$	—
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$	$(1.48 \pm 0.12) \times 10^{-3}$	—
3-body total		
$\sigma \pi^+ \pi^-$	$(6.1 \pm 0.9) \times 10^{-4}$	—
$f_0(980) \pi^+ \pi^-, f_0 \rightarrow$	$(1.8 \pm 0.5) \times 10^{-4}$	—
$\pi^+ \pi^-$		
$f_2(1270) \pi^+ \pi^-, f_2 \rightarrow$	$(3.6 \pm 0.6) \times 10^{-4}$	—
$\pi^+ \pi^-$		
$\pi^+ \pi^- 2\pi^0$	$(1.00 \pm 0.09) \%$	882
$\eta \pi^0$	[r] $(6.8 \pm 0.7) \times 10^{-4}$	846
$\omega \pi^0$	[r] $< 2.6 \times 10^{-4}$	CL=90% 761
$2\pi^+ 2\pi^- \pi^0$	$(4.1 \pm 0.5) \times 10^{-3}$	844
$\eta \pi^+ \pi^-$	[r] $(1.09 \pm 0.16) \times 10^{-3}$	827
$\omega \pi^+ \pi^-$	[r] $(1.6 \pm 0.5) \times 10^{-3}$	738
$3\pi^+ 3\pi^-$	$(4.2 \pm 1.2) \times 10^{-4}$	795
$\eta'(958) \pi^0$	$(9.0 \pm 1.4) \times 10^{-4}$	678
$\eta'(958) \pi^+ \pi^-$	$(4.5 \pm 1.7) \times 10^{-4}$	650
2η	$(1.67 \pm 0.20) \times 10^{-3}$	755
$\eta \eta'(958)$	$(1.05 \pm 0.26) \times 10^{-3}$	537

Hadronic modes with a $K\bar{K}$ pair

$K^+ K^-$	$(3.96 \pm 0.08) \times 10^{-3}$	S=1.4	791
$2K_S^0$	$(1.7 \pm 0.4) \times 10^{-4}$	S=2.5	789
$K_S^0 K^- \pi^+$	$(3.5 \pm 0.5) \times 10^{-3}$	S=1.2	739
$\bar{K}^{*0}(892)^0 K_S^0, \bar{K}^{*0} \rightarrow$	$< 5 \times 10^{-4}$	CL=90%	608
$K^- \pi^+$			
$K_S^0 K^+ \pi^-$	$(2.1 \pm 0.4) \times 10^{-3}$	S=1.3	739
$K^{*0}(892)^0 K_S^0, K^{*0} \rightarrow$	$< 1.8 \times 10^{-4}$	CL=90%	608
$K^+ \pi^-$			
$K^+ K^- \pi^0$	$(3.29 \pm 0.14) \times 10^{-3}$		743
$K^{*}(892)^+ K^-, K^{*}(892)^+ \rightarrow$	$(1.46 \pm 0.07) \times 10^{-3}$		—
$K^+ \pi^0$			
$K^{*}(892)^- K^+, K^{*}(892)^- \rightarrow$	$(5.2 \pm 0.4) \times 10^{-4}$		—
$K^- \pi^0$			

$(K^+\pi^0)_{S\text{-wave}} K^-$	$(2.34 \pm 0.17) \times 10^{-3}$		743
$(K^-\pi^0)_{S\text{-wave}} K^+$	$(1.3 \pm 0.4) \times 10^{-4}$		743
$f_0(980)\pi^0, f_0 \rightarrow K^+ K^-$	$(3.5 \pm 0.6) \times 10^{-4}$		—
$\phi\pi^0, \phi \rightarrow K^+ K^-$	$(6.4 \pm 0.4) \times 10^{-4}$		—
$2K_S^0\pi^0$	$< 5.9 \times 10^{-4}$		740
$K^+ K^- \pi^+ \pi^-$	$(2.43 \pm 0.12) \times 10^{-3}$		677
$\phi(\pi^+\pi^-)_{S\text{-wave}}, \phi \rightarrow K^+ K^-$	$(2.50 \pm 0.33) \times 10^{-4}$		614
$(\phi\rho^0)_{S\text{-wave}}, \phi \rightarrow K^+ K^-$	$(9.3 \pm 1.2) \times 10^{-4}$		250
$(\phi\rho^0)_{D\text{-wave}}, \phi \rightarrow K^+ K^-$	$(8.3 \pm 2.3) \times 10^{-5}$		—
$(K^{*0}\bar{K}^{*0})_{S\text{-wave}}, K^{*0} \rightarrow K^\pm \pi^\mp$	$(1.48 \pm 0.30) \times 10^{-4}$		—
$(K^-\pi^+)_{P\text{-wave}}, (K^+\pi^-)_{S\text{-wave}},$	$(2.6 \pm 0.5) \times 10^{-4}$		—
$K_1(1270)^+ K^-$	$(1.8 \pm 0.5) \times 10^{-4}$		—
$K_1(1270)^+ \rightarrow K^{*0}\pi^+$			
$K_1(1270)^+ K^-$	$(1.14 \pm 0.26) \times 10^{-4}$		—
$K_1(1270)^+ \rightarrow \rho^0 K^+$			
$K_1(1270)^- K^+$	$(2.2 \pm 1.2) \times 10^{-5}$		—
$K_1(1270)^- \rightarrow \bar{K}^{*0}\pi^-$			
$K_1(1270)^- K^+$	$(1.46 \pm 0.25) \times 10^{-4}$		—
$K_1(1270)^- \rightarrow \rho^0 K^-$			
$K^*(1410)^+ K^-$	$(1.02 \pm 0.26) \times 10^{-4}$		—
$K^*(1410)^+ \rightarrow K^{*0}\pi^+$			
$K^*(1410)^- K^+$	$(1.14 \pm 0.25) \times 10^{-4}$		—
$K^*(1410)^- \rightarrow \bar{K}^{*0}\pi^-$			
$2K_S^0\pi^+\pi^-$	$(1.23 \pm 0.24) \times 10^{-3}$		673
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.5 \times 10^{-4}$	CL=90%	595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$		600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi\eta$	$(1.4 \pm 0.5) \times 10^{-4}$		489
$\phi\omega$	$< 2.1 \times 10^{-3}$	CL=90%	238

Radiative modes

$\rho^0\gamma$	$< 2.4 \times 10^{-4}$	CL=90%	771
$\omega\gamma$	$< 2.4 \times 10^{-4}$	CL=90%	768
$\phi\gamma$	$(2.70 \pm 0.35) \times 10^{-5}$		654
$\bar{K}^*(892)^0\gamma$	$(3.27 \pm 0.34) \times 10^{-4}$		719

**Doubly Cabibbo suppressed (DC) modes or
 $\Delta C = 2$ forbidden via mixing (C2M) modes**

$K^+ \ell^- \bar{\nu}_\ell$ via $\bar{D}^0$		< 2.2	$\times 10^{-5}$	CL=90%	—
K^+ or $K^*(892)^+$ $e^- \bar{\nu}_e$ via $\bar{D}^0$		< 6	$\times 10^{-5}$	CL=90%	—
$K^+ \pi^-$	DC	$(1.47 \pm 0.07) \times 10^{-4}$		S=2.8	861
$K^+ \pi^-$ via DCS		$(1.31 \pm 0.08) \times 10^{-4}$			—
$K^+ \pi^-$ via $\bar{D}^0$		< 1.6	$\times 10^{-5}$	CL=95%	861
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$		< 1.8	$\times 10^{-4}$	CL=95%	—
$K^*(892)^+ \pi^-$, $K^*(892)^+ \rightarrow K_S^0 \pi^+$	DC	$(1.14 \pm_{-0.34}^{+0.60}) \times 10^{-4}$			711
$K_0^*(1430)^+ \pi^-$, $K_0^*(1430)^+ \rightarrow K_S^0 \pi^+$	DC	< 1.4	$\times 10^{-5}$		—
$K_2^*(1430)^+ \pi^-$, $K_2^*(1430)^+ \rightarrow K_S^0 \pi^+$	DC	< 3.4	$\times 10^{-5}$		—
$K^+ \pi^- \pi^0$	DC	$(3.04 \pm 0.17) \times 10^{-4}$			844
$K^+ \pi^- \pi^0$ via $\bar{D}^0$		$(7.3 \pm 0.5) \times 10^{-4}$			—
$K^+ \pi^+ 2\pi^-$	DC	$(2.62 \pm_{-0.19}^{+0.21}) \times 10^{-4}$			813
$K^+ \pi^+ 2\pi^-$ via $\bar{D}^0$		< 4	$\times 10^{-4}$	CL=90%	812
μ^- anything via $\bar{D}^0$		< 4	$\times 10^{-4}$	CL=90%	—

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton Family number (LF) violating modes,
Lepton (L) or Baryon (B) number violating modes**

$\gamma\gamma$	C1	< 2.2	$\times 10^{-6}$	CL=90%	932
$e^+ e^-$	C1	< 7.9	$\times 10^{-8}$	CL=90%	932
$\mu^+ \mu^-$	C1	< 1.4	$\times 10^{-7}$	CL=90%	926
$\pi^0 e^+ e^-$	C1	< 4.5	$\times 10^{-5}$	CL=90%	928
$\pi^0 \mu^+ \mu^-$	C1	< 1.8	$\times 10^{-4}$	CL=90%	915
$\eta e^+ e^-$	C1	< 1.1	$\times 10^{-4}$	CL=90%	852
$\eta \mu^+ \mu^-$	C1	< 5.3	$\times 10^{-4}$	CL=90%	838
$\pi^+ \pi^- e^+ e^-$	C1	< 3.73	$\times 10^{-4}$	CL=90%	922
$\rho^0 e^+ e^-$	C1	< 1.0	$\times 10^{-4}$	CL=90%	771
$\pi^+ \pi^- \mu^+ \mu^-$	C1	< 3.0	$\times 10^{-5}$	CL=90%	894
$\rho^0 \mu^+ \mu^-$	C1	< 2.2	$\times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	C1	< 1.8	$\times 10^{-4}$	CL=90%	768
$\omega \mu^+ \mu^-$	C1	< 8.3	$\times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	C1	< 3.15	$\times 10^{-4}$	CL=90%	791
$\phi e^+ e^-$	C1	< 5.2	$\times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	C1	< 3.3	$\times 10^{-5}$	CL=90%	710
$\phi \mu^+ \mu^-$	C1	< 3.1	$\times 10^{-5}$	CL=90%	631
$\bar{K}^0 e^+ e^-$	[h]	< 1.1	$\times 10^{-4}$	CL=90%	866
$\bar{K}^0 \mu^+ \mu^-$	[h]	< 2.6	$\times 10^{-4}$	CL=90%	852

$K^- \pi^+ e^+ e^-$	CI	< 3.85	$\times 10^{-4}$	CL=90%	861
$\overline{K}^*(892)^0 e^+ e^-$	$[h]$	< 4.7	$\times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	CI	< 3.59	$\times 10^{-4}$	CL=90%	829
$\overline{K}^*(892)^0 \mu^+ \mu^-$	$[h]$	< 2.4	$\times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	CI	< 8.1	$\times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	LF	$[s] < 2.6$	$\times 10^{-7}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	LF	$[s] < 8.6$	$\times 10^{-5}$	CL=90%	924
$\eta e^\pm \mu^\mp$	LF	$[s] < 1.0$	$\times 10^{-4}$	CL=90%	848
$\pi^+ \pi^- e^\pm \mu^\mp$	LF	$[s] < 1.5$	$\times 10^{-5}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	LF	$[s] < 4.9$	$\times 10^{-5}$	CL=90%	767
$\omega e^\pm \mu^\mp$	LF	$[s] < 1.2$	$\times 10^{-4}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	LF	$[s] < 1.8$	$\times 10^{-4}$	CL=90%	754
$\phi e^\pm \mu^\mp$	LF	$[s] < 3.4$	$\times 10^{-5}$	CL=90%	648
$\overline{K}^0 e^\pm \mu^\mp$	LF	$[s] < 1.0$	$\times 10^{-4}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	LF	$[s] < 5.53$	$\times 10^{-4}$	CL=90%	848
$\overline{K}^*(892)^0 e^\pm \mu^\mp$	LF	$[s] < 8.3$	$\times 10^{-5}$	CL=90%	714
$2\pi^- 2e^+ + \text{c.c.}$	L	< 1.12	$\times 10^{-4}$	CL=90%	922
$2\pi^- 2\mu^+ + \text{c.c.}$	L	< 2.9	$\times 10^{-5}$	CL=90%	894
$K^- \pi^- 2e^+ + \text{c.c.}$	L	< 2.06	$\times 10^{-4}$	CL=90%	861
$K^- \pi^- 2\mu^+ + \text{c.c.}$	L	< 3.9	$\times 10^{-4}$	CL=90%	829
$2K^- 2e^+ + \text{c.c.}$	L	< 1.52	$\times 10^{-4}$	CL=90%	791
$2K^- 2\mu^+ + \text{c.c.}$	L	< 9.4	$\times 10^{-5}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	L	< 7.9	$\times 10^{-5}$	CL=90%	911
$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	L	< 2.18	$\times 10^{-4}$	CL=90%	848
$2K^- e^+ \mu^+ + \text{c.c.}$	L	< 5.7	$\times 10^{-5}$	CL=90%	754
$p e^-$	L, B	$[t] < 1.0$	$\times 10^{-5}$	CL=90%	696
$\overline{p} e^+$	L, B	$[u] < 1.1$	$\times 10^{-5}$	CL=90%	696

 $D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

 I, J, P need confirmation.Mass $m = 2006.99 \pm 0.15$ MeV $m_{D^{*0}} - m_{D^0} = 142.12 \pm 0.07$ MeVFull width $\Gamma < 2.1$ MeV, CL = 90% $\overline{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^0$	$(61.9 \pm 2.9) \%$	43
$D^0 \gamma$	$(38.1 \pm 2.9) \%$	137

$D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

 I, J, P need confirmation.Mass $m = 2010.29 \pm 0.13$ MeV

$$m_{D^*(2010)^+} - m_{D^+} = 140.66 \pm 0.10 \text{ MeV} \quad (S = 1.1)$$

$$m_{D^*(2010)^+} - m_{D^0} = 145.421 \pm 0.010 \text{ MeV} \quad (S = 1.1)$$

Full width $\Gamma = 96 \pm 22$ keV $D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	$(67.7 \pm 0.5) \%$	39
$D^+ \pi^0$	$(30.7 \pm 0.5) \%$	38
$D^+ \gamma$	$(1.6 \pm 0.4) \%$	136

 $D_0^*(2400)^0$

$$I(J^P) = \frac{1}{2}(0^+)$$

Mass $m = 2318 \pm 29$ MeV ($S = 1.7$)Full width $\Gamma = 267 \pm 40$ MeV

$D_0^*(2400)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+ \pi^-$	seen	385

 $D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

 I needs confirmation.Mass $m = 2421.4 \pm 0.6$ MeV ($S = 1.2$)

$$m_{D_1^0} - m_{D^{*+}} = 411.1 \pm 0.6 \text{ MeV} \quad (S = 1.2)$$

Full width $\Gamma = 27.4 \pm 2.5$ MeV ($S = 2.3$) $\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

$D_1(2420)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^*(2010)^+ \pi^-$	seen	354
$D^0 \pi^+ \pi^-$	seen	425
$D^+ \pi^-$	not seen	473
$D^{*0} \pi^+ \pi^-$	not seen	280

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2462.6 \pm 0.6 \text{ MeV} \quad (S = 1.2)$$

$$m_{D_2^{*0}} - m_{D^+} = 593.0 \pm 0.6 \text{ MeV} \quad (S = 1.3)$$

$$m_{D_2^{*0}} - m_{D^{*+}} = 452.3 \pm 0.6 \text{ MeV} \quad (S = 1.3)$$

$$\text{Full width } \Gamma = 49.0 \pm 1.3 \text{ MeV} \quad (S = 1.5)$$

 $\bar{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

$D_2^*(2460)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+ \pi^-$	seen	507
$D^*(2010)^+ \pi^-$	seen	391
$D^0 \pi^+ \pi^-$	not seen	463
$D^{*0} \pi^+ \pi^-$	not seen	326

 $D_2^*(2460)^\pm$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2464.3 \pm 1.6 \text{ MeV} \quad (S = 1.7)$$

$$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7 \text{ MeV}$$

$$\text{Full width } \Gamma = 37 \pm 6 \text{ MeV} \quad (S = 1.4)$$

 $D_2^*(2460)^-$ modes are charge conjugates of modes below.

$D_2^*(2460)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	seen	512
$D^{*0} \pi^+$	seen	395
$D^+ \pi^+ \pi^-$	not seen	461
$D^{*+} \pi^+ \pi^-$	not seen	324

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- 2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] Submodes of the $D^+ \rightarrow K^- 2\pi^+ \pi^0$ and $K_S^0 2\pi^+ \pi^-$ modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 *Review*, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\bar{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
- [n] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.19 ± 0.17 %.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.
- [q] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [r] This branching fraction includes all the decay modes of the resonance in the final state.

[s] The value is for the sum of the charge states or particle/antiparticle states indicated.

[t] This limit is for either D^0 or $\bar{D}^0$ to $p e^-$.

[u] This limit is for either D^0 or $\bar{D}^0$ to $\bar{p} e^+$.