

$\omega(1420)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\omega(1420)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>(1400–1450) OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1470 \pm 50$	13.1k	<sup>1</sup> AULCHENKO 15A	SND	$1.05\text{--}1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$1382 \pm 23 \pm 70$		AUBERT 07AU	BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
$1350 \pm 20 \pm 20$		AUBERT,B 04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$1400 \pm 50 \pm 130$	1.2M	<sup>2</sup> ACHASOV 03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$1450 \pm 10$		<sup>3</sup> HENNER 02	RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
$1373 \pm 70$	177	<sup>4</sup> AKHMETSHIN 00D	CMD2	$1.2\text{--}1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
$1370 \pm 25$	5095	ANISOVICH 00H	SPEC	$0.0 \rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
$1400^{+100}_{-200}$		<sup>5</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$\sim 1400$		<sup>6</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
$\sim 1460$		<sup>7</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow K^+K^-$
$1440 \pm 70$		<sup>8</sup> CLEGG 94	RVUE	
$1419 \pm 31$	315	<sup>9</sup> ANTONELLI 92	DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi$

<sup>1</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .<sup>2</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.<sup>3</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.<sup>4</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.<sup>5</sup> Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.<sup>6</sup> Using the data from ANTONELLI 92.<sup>7</sup> Using the data from IVANOV 81 and BISELLO 88B.<sup>8</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.<sup>9</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega, \phi$  tails with fixed (+, -, +) phases.

### $\omega(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>(180–250) OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$880 \pm 170$	13.1k	<sup>10</sup> AULCHENKO 15A	SND	$1.05\text{--}1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$130 \pm 50 \pm 100$		AUBERT 07AU	BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
$450 \pm 70 \pm 70$		AUBERT,B 04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$870^{+500}_{-300} \pm 450$	1.2M	<sup>11</sup> ACHASOV 03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$199 \pm 15$		<sup>12</sup> HENNER 02	RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

188 ± 45	177	<sup>13</sup> AKHMETSHIN 00D	CMD2	1.2–1.38	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
360 <sup>+100</sup> <sub>-60</sub>	5095	ANISOVICH 00H	SPEC	0.0	$\rho \bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$
240 ± 70		<sup>14</sup> CLEGG 94	RVUE		
174 ± 59	315	<sup>15</sup> ANTONELLI 92	DM2	1.34–2.4	$e^+ e^- \rightarrow \rho \pi$

<sup>10</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .  
<sup>11</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+ \pi^- \pi^0$  and ANTONELLI 92 on the  $\omega \pi^+ \pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.  
<sup>12</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.  
<sup>13</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho \pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.  
<sup>14</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.  
<sup>15</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega, \phi$  tails with fixed (+, -, +) phases.

### $\omega(1420)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho \pi$	dominant
$\Gamma_2$ $\omega \pi \pi$	seen
$\Gamma_3$ $\omega \eta$	
$\Gamma_4$ $b_1(1235) \pi$	seen
$\Gamma_5$ $e^+ e^-$	seen
$\Gamma_6$ $\pi^0 \gamma$	

### $\omega(1420) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

$\Gamma(\rho \pi) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_1 / \Gamma \times \Gamma_5 / \Gamma$			
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.73 ± 0.08	13.1k	<sup>16</sup> AULCHENKO 15A	SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.82 ± 0.05 ± 0.06		AUBERT,B 04N	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
0.65 ± 0.13 ± 0.21	1.2M	<sup>17,18</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.625 ± 0.160		<sup>19,20</sup> CLEGG 94	RVUE	
0.466 ± 0.178		<sup>21,22</sup> ANTONELLI 92	DM2	1.34–2.4 $e^+ e^- \rightarrow \rho \pi$

- <sup>16</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .  
<sup>17</sup> Calculated by us from the cross section at the peak.  
<sup>18</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+ \pi^- \pi^0$  and ANTONELLI 92 on the  $\omega \pi^+ \pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.  
<sup>19</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.  
<sup>20</sup> From the partial and leptonic width given by the authors.  
<sup>21</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega, \phi$  tails with fixed (+, -, +) phases.  
<sup>22</sup> From the product of the leptonic width and partial branching ratio given by the authors.

**$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma \times \Gamma_5/\Gamma$**

VALUE (units  $10^{-8}$ )      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.7 ± 5.7      AUBERT      07AU BABR      10.6  $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$   
 1.9 ± 1.9      <sup>23</sup> AKHMETSHIN 00D      CMD2      1.2–2.4  $e^+e^- \rightarrow \omega\pi^+\pi^-$

<sup>23</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

**$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$**

VALUE (units  $10^{-8}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6<sup>+0.9</sup><sub>-0.7</sub>      898      <sup>24</sup> ACHASOV      16B      SND      1.34–2.00  $e^+e^- \rightarrow \omega\eta$

<sup>24</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass and the width of  $\omega(1420)$  are fixed to the 2014 edition (PDG 14) of this review.

**$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma \times \Gamma_5/\Gamma$**

VALUE (units  $10^{-8}$ )      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.03<sup>+0.70</sup><sub>-0.75</sub>      <sup>25</sup> AKHMETSHIN 05      CMD2      0.60–1.38  $e^+e^- \rightarrow \pi^0\gamma$

<sup>25</sup> Using 1420 MeV and 220 MeV for the  $\omega(1420)$  mass and width.

**$\omega(1420)$  BRANCHING RATIOS**

**$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$**

VALUE      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.301 ± 0.029      <sup>26</sup> HENNER      02      RVUE      1.2–2.0  $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$   
 possibly seen      AKHMETSHIN 00D      CMD2       $e^+e^- \rightarrow \omega\pi^+\pi^-$

**$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$   $\Gamma_2/\Gamma_4$**

VALUE      EVTS      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.60 ± 0.16      5095      ANISOVICH      00H      SPEC      0.0  $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

**$\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$**

VALUE      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.699 ± 0.029      <sup>26</sup> HENNER      02      RVUE      1.2–2.0  $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 6.6$	1.2M	<sup>27,28</sup> ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$23 \pm 1$		<sup>26</sup> HENNER	02 RVUE	$1.2\text{--}2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

<sup>26</sup> Assuming that the  $\omega(1420)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

<sup>27</sup> Calculated by us from the cross section at the peak.

<sup>28</sup> Assuming that the  $\omega(1420)$  decays into  $\rho\pi$  only.

### $\omega(1420)$ REFERENCES

ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
		Translated from ZETF 148 34.		
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)