

$\Sigma(1620) 1/2^-$  $I(J^P) = 1(\frac{1}{2}^-)$  Status: \*

## OMITTED FROM SUMMARY TABLE

The  $S_{11}$  state at 1697 MeV reported by VANHORN 75 is tentatively listed under the  $\Sigma(1750)$ . CARROLL 76 sees two bumps in the isospin-1 total cross section near this mass. GAO 12 sees no evidence for this resonance.

Production experiments are listed separately in the next entry.

 **$\Sigma(1620)$  POLE POSITION****REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1501	ZHANG	13A	DPWA Multichannel

**-2×IMAGINARY PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
171	ZHANG	13A	DPWA Multichannel

 **$\Sigma(1620)$  MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>≈ 1620 OUR ESTIMATE</b>			
1600 ± 15	ZHANG	13A	DPWA Multichannel
1600 ± 6	<sup>1</sup> MORRIS	78	DPWA $K^- n \rightarrow \Lambda \pi^-$
1608 ± 5	<sup>2</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
1633 ± 10	<sup>3</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
1630 ± 10	LANGBEIN	72	IPWA $\bar{K}N$ multichannel
1620	KIM	71	DPWA K-matrix analysis

 **$\Sigma(1620)$  WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
400 ± 152	ZHANG	13A	DPWA Multichannel
87 ± 19	<sup>1</sup> MORRIS	78	DPWA $K^- n \rightarrow \Lambda \pi^-$
15	<sup>2</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
10	<sup>3</sup> CARROLL	76	DPWA Isospin-1 total $\sigma$
65 ± 20	LANGBEIN	72	IPWA $\bar{K}N$ multichannel
40	KIM	71	DPWA K-matrix analysis

## Σ(1620) DECAY MODES

Mode
Γ <sub>1</sub> $N\bar{K}$
Γ <sub>2</sub> $\Lambda\pi$
Γ <sub>3</sub> $\Sigma\pi$

## Σ(1620) BRANCHING RATIOS

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.59±0.10	ZHANG    13A    DPWA    Multichannel
0.22±0.02	LANGBEIN    72    IPWA $\bar{K}N$ multichannel
0.05	KIM    71    DPWA    K-matrix analysis

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Lambda\pi$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.12±0.02	<sup>1</sup> MORRIS    78    DPWA $K^-n \rightarrow \Lambda\pi^-$
not seen	BAILLON    75    IPWA $\bar{K}N \rightarrow \Lambda\pi$
0.15	KIM    71    DPWA    K-matrix analysis

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Sigma\pi$	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.32±0.03	ZHANG    13A    DPWA    Multichannel
not seen	HEPP    76B    DPWA $K^-N \rightarrow \Sigma\pi$
+0.40±0.06	LANGBEIN    72    IPWA $\bar{K}N$ multichannel
+0.08	KIM    71    DPWA    K-matrix analysis

## Σ(1620) FOOTNOTES

- <sup>1</sup> MORRIS 78 obtains an equally good fit without including this resonance.  
<sup>2</sup> Total cross-section bump with  $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$  is 0.06 seen by CARROLL 76.  
<sup>3</sup> Total cross-section bump with  $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$  is 0.04 seen by CARROLL 76.

## Σ(1620) REFERENCES

ZHANG    13A    PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
GAO    12    PR C86 025201	P. Gao, J. Shi, B.S. Zou	(BHEP, BEIJT)
Also	P. Gao, B.S. Zou, A. Sibirtsev	(BHEP, BEIJT+)
MORRIS    78    PR D17 55	W.A. Morris <i>et al.</i>	(FSU) IJP
CARROLL    76    PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
HEPP    76B    PL 65B 487	V. Hepp <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
BAILLON    75    NP B94 39	P.H. Baillon, P.J. Litchfield	(CERN, RHEL) IJP
VANHORN    75    NP B87 145	A.J. van Horn	(LBL) IJP
Also	A.J. van Horn	(LBL) IJP
LANGBEIN    72    NP B47 477	W. Langbein, F. Wagner	(MPIM) IJP
KIM    71    PRL 27 356	J.K. Kim	(HARV) IJP
Also	J.K. Kim	(HARV) IJP
Hyperon Resonances, 1970		