

$\Upsilon(1D)$

$$I^G(J^{PC}) = 0^-(2^{- -})$$

First observed by BONVICINI 04 in the decay to $\gamma\gamma \Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+\pi^- \Upsilon(1S)$. Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

$\Upsilon(1D)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10163.7 ± 1.4 OUR AVERAGE		Error includes scale factor of 1.7.		
10164.5 ± 0.8 ± 0.5		DEL-AMO-SA...10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1 ± 0.6 ± 1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Upsilon(1D)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma\gamma \Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta \Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

$\Upsilon(1D)$ BRANCHING RATIOS

$\Gamma(\eta \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_3/Γ_1
<u>VALUE</u>	<u>CL%</u>
<0.25	90
<u>DOCUMENT ID</u>	<u>TECN</u>
BONVICINI 04	CLE3
<u>COMMENT</u>	
$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>
$0.66^{+0.15}_{-0.14} \pm 0.06$	¹ DEL-AMO-SA...10R
<u>TECN</u>	<u>COMMENT</u>
BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$

¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(1D))$.

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_4/Γ_1
<u>VALUE</u>	<u>CL%</u>
<1.2	90
<u>DOCUMENT ID</u>	<u>TECN</u>
² BONVICINI 04	CLE3
<u>COMMENT</u>	
$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

² Assuming $J = 2$.

$\Upsilon(1D)$ REFERENCES

DEL-AMO-SA... 10R	PR D82 111102	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
BONVICINI 04	PR D70 032001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)