

$b\bar{b}$ MESONS

$\eta_b(1S)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

Mass $m = 9399.0 \pm 2.3$ MeV (S = 1.6)

Full width $\Gamma = 10^{+5}_{-4}$ MeV

$\eta_b(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
hadrons	seen		—
$3h^+3h^-$	not seen		4673
$2h^+2h^-$	not seen		4689
$\gamma\gamma$	not seen		4700
$\mu^+\mu^-$	$<9 \times 10^{-3}$	90%	4698
$\tau^+\tau^-$	$<8\%$	90%	4351

$\Upsilon(1S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 9460.30 \pm 0.26$ MeV (S = 3.3)

Full width $\Gamma = 54.02 \pm 1.25$ keV

$\Gamma_{ee} = 1.340 \pm 0.018$ keV

$\Upsilon(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\tau^+\tau^-$	(2.60 \pm 0.10) %		4384
e^+e^-	(2.38 \pm 0.11) %		4730
$\mu^+\mu^-$	(2.48 \pm 0.05) %		4729

Hadronic decays

ggg	(81.7 \pm 0.7) %		—
γgg	(2.2 \pm 0.6) %		—
$\eta'(958)$ anything	(2.94 \pm 0.24) %		—
$J/\psi(1S)$ anything	(5.4 \pm 0.4) $\times 10^{-4}$	S=1.4	4223
$J/\psi(1S)\eta_c$	< 2.2	$\times 10^{-6}$ CL=90%	3624
$J/\psi(1S)\chi_{c0}$	< 3.4	$\times 10^{-6}$ CL=90%	3429
$J/\psi(1S)\chi_{c1}$	(3.9 \pm 1.2) $\times 10^{-6}$		3382
$J/\psi(1S)\chi_{c2}$	< 1.4	$\times 10^{-6}$ CL=90%	3359
$J/\psi(1S)\eta_c(2S)$	< 2.2	$\times 10^{-6}$ CL=90%	3316
$J/\psi(1S)X(3940)$	< 5.4	$\times 10^{-6}$ CL=90%	3148
$J/\psi(1S)X(4160)$	< 5.4	$\times 10^{-6}$ CL=90%	3018
$X(4350)$ anything, $X \rightarrow$ $J/\psi(1S)\phi$	< 8.1	$\times 10^{-6}$ CL=90%	—

$X(3900)^\pm$ anything, $X \rightarrow J/\psi(1S)\pi^\pm$	< 1.3	$\times 10^{-5}$	CL=90%	—
$X(4200)^\pm$ anything, $X \rightarrow J/\psi(1S)\pi^\pm$	< 6.0	$\times 10^{-5}$	CL=90%	—
$X(4430)^\pm$ anything, $X \rightarrow J/\psi(1S)\pi^\pm$	< 4.9	$\times 10^{-5}$	CL=90%	—
X_{cs}^\pm anything, $X \rightarrow J/\psi K^\pm$	< 5.7	$\times 10^{-6}$	CL=90%	—
$X(3872)$ anything, $X \rightarrow J/\psi(1S)\pi^+\pi^-$	< 9.5	$\times 10^{-6}$	CL=90%	—
$X(4260)$ anything, $X \rightarrow J/\psi(1S)\pi^+\pi^-$	< 3.8	$\times 10^{-5}$	CL=90%	—
$X(4260)$ anything, $X \rightarrow J/\psi(1S)K^+K^-$	< 7.5	$\times 10^{-6}$	CL=90%	—
$X(4140)$ anything, $X \rightarrow J/\psi(1S)\phi$	< 5.2	$\times 10^{-6}$	CL=90%	—
χ_{c0} anything	< 4	$\times 10^{-3}$	CL=90%	—
χ_{c1} anything	(1.90 ± 0.35)	$\times 10^{-4}$		—
χ_{c2} anything	(2.8 ± 0.8)	$\times 10^{-4}$		—
$\psi(2S)$ anything	(1.23 ± 0.20)	$\times 10^{-4}$		—
$\psi(2S)\eta_c$	< 3.6	$\times 10^{-6}$	CL=90%	3345
$\psi(2S)\chi_{c0}$	< 6.5	$\times 10^{-6}$	CL=90%	3124
$\psi(2S)\chi_{c1}$	< 4.5	$\times 10^{-6}$	CL=90%	3070
$\psi(2S)\chi_{c2}$	< 2.1	$\times 10^{-6}$	CL=90%	3043
$\psi(2S)\eta_c(2S)$	< 3.2	$\times 10^{-6}$	CL=90%	2993
$\psi(2S)X(3940)$	< 2.9	$\times 10^{-6}$	CL=90%	2797
$\psi(2S)X(4160)$	< 2.9	$\times 10^{-6}$	CL=90%	2642
$X(4260)$ anything, $X \rightarrow \psi(2S)\pi^+\pi^-$	< 7.9	$\times 10^{-5}$	CL=90%	—
$X(4360)$ anything, $X \rightarrow \psi(2S)\pi^+\pi^-$	< 5.2	$\times 10^{-5}$	CL=90%	—
$X(4660)$ anything, $X \rightarrow \psi(2S)\pi^+\pi^-$	< 2.2	$\times 10^{-5}$	CL=90%	—
$X(4050)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	< 8.8	$\times 10^{-5}$	CL=90%	—
$X(4430)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	< 6.7	$\times 10^{-5}$	CL=90%	—
$\rho\pi$	< 3.68	$\times 10^{-6}$	CL=90%	4697
$\omega\pi^0$	< 3.90	$\times 10^{-6}$	CL=90%	4697
$\pi^+\pi^-$	< 5	$\times 10^{-4}$	CL=90%	4728
K^+K^-	< 5	$\times 10^{-4}$	CL=90%	4704
$p\bar{p}$	< 5	$\times 10^{-4}$	CL=90%	4636
$\pi^+\pi^-\pi^0$	(2.1 ± 0.8)	$\times 10^{-6}$		4725
ϕK^+K^-	(2.4 ± 0.5)	$\times 10^{-6}$		4622

$\omega \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$	4694
$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(4.4 \pm 0.8) \times 10^{-6}$	4667
$\phi f'_2(1525)$	$< 1.63 \times 10^{-6}$ CL=90%	4549
$\omega f_2(1270)$	$< 1.79 \times 10^{-6}$ CL=90%	4611
$\rho(770) a_2(1320)$	$< 2.24 \times 10^{-6}$ CL=90%	4605
$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(3.0 \pm 0.8) \times 10^{-6}$	4579
$K_1(1270)^\pm K^\mp$	$< 2.41 \times 10^{-6}$ CL=90%	4631
$K_1(1400)^\pm K^\mp$	$(1.0 \pm 0.4) \times 10^{-6}$	4613
$b_1(1235)^\pm \pi^\mp$	$< 1.25 \times 10^{-6}$ CL=90%	4649
$\pi^+ \pi^- \pi^0 \pi^0$	$(1.28 \pm 0.30) \times 10^{-5}$	4720
$K_S^0 K^+ \pi^- + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-6}$	4696
$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(2.9 \pm 0.9) \times 10^{-6}$	4675
$K^*(892)^- K^+ + \text{c.c.}$	$< 1.11 \times 10^{-6}$ CL=90%	4675
$D^*(2010)^\pm \text{ anything}$	$(2.52 \pm 0.20) \%$	—
${}^2H \text{ anything}$	$(2.85 \pm 0.25) \times 10^{-5}$	—
Sum of 100 exclusive modes	$(1.200 \pm 0.017) \%$	—

Radiative decays

$\gamma \pi^+ \pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	4728
$\gamma \pi^0 \pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	4728
$\gamma \pi^0 \eta$	$< 2.4 \times 10^{-6}$ CL=90%	4713
$\gamma K^+ K^-$	[a] $(1.14 \pm 0.13) \times 10^{-5}$	4704
$\gamma p \bar{p}$	[b] $< 6 \times 10^{-6}$ CL=90%	4636
$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	4720
$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	4703
$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	4679
$\gamma \pi^+ \pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	4686
$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	4720
$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	4703
$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	4658
$\gamma \pi^+ \pi^- p \bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	4604
$\gamma 2\pi^+ 2\pi^- p \bar{p}$	$(4 \pm 6) \times 10^{-5}$	4563
$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	4601
$\gamma \eta'(958)$	$< 1.9 \times 10^{-6}$ CL=90%	4682
$\gamma \eta$	$< 1.0 \times 10^{-6}$ CL=90%	4714
$\gamma f_0(980)$	$< 3 \times 10^{-5}$ CL=90%	4678
$\gamma f'_2(1525)$	$(3.8 \pm 0.9) \times 10^{-5}$	4607
$\gamma f_2(1270)$	$(1.01 \pm 0.09) \times 10^{-4}$	4644
$\gamma \eta(1405)$	$< 8.2 \times 10^{-5}$ CL=90%	4625
$\gamma f_0(1500)$	$< 1.5 \times 10^{-5}$ CL=90%	4611
$\gamma f_0(1710)$	$< 2.6 \times 10^{-4}$ CL=90%	4573
$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	$< 7 \times 10^{-6}$ CL=90%	—
$\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0$	$< 1.4 \times 10^{-6}$ CL=90%	—
$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$< 1.8 \times 10^{-6}$ CL=90%	—

$\gamma f_4(2050)$	< 5.3	$\times 10^{-5}$	CL=90%	4515
$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	< 2	$\times 10^{-4}$	CL=90%	4475
$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	< 8	$\times 10^{-7}$	CL=90%	4469
$\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-$	< 6	$\times 10^{-7}$	CL=90%	—
$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	< 1.1	$\times 10^{-6}$	CL=90%	—
$\gamma \eta(2225) \rightarrow \gamma \phi \phi$	< 3	$\times 10^{-3}$	CL=90%	4469
$\gamma \eta_c(1S)$	< 5.7	$\times 10^{-5}$	CL=90%	4260
$\gamma \chi_{c0}$	< 6.5	$\times 10^{-4}$	CL=90%	4114
$\gamma \chi_{c1}$	< 2.3	$\times 10^{-5}$	CL=90%	4079
$\gamma \chi_{c2}$	< 7.6	$\times 10^{-6}$	CL=90%	4062
$\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi$	< 1.6	$\times 10^{-6}$	CL=90%	—
$\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	—
$\gamma X(3915) \rightarrow \omega J/\psi$	< 3.0	$\times 10^{-6}$	CL=90%	—
$\gamma X(4140) \rightarrow \phi J/\psi$	< 2.2	$\times 10^{-6}$	CL=90%	—
γX	[c] < 4.5	$\times 10^{-6}$	CL=90%	—
$\gamma X \bar{X} (m_X < 3.1 \text{ GeV})$	[d] < 1	$\times 10^{-3}$	CL=90%	—
$\gamma X \bar{X} (m_X < 4.5 \text{ GeV})$	[e] < 2.4	$\times 10^{-4}$	CL=90%	—
$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[f] < 1.78	$\times 10^{-4}$	CL=95%	—
$\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$	[g] < 9	$\times 10^{-6}$	CL=90%	—
$\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$	[a] < 1.30	$\times 10^{-4}$	CL=90%	—
$\gamma a_1^0 \rightarrow \gamma g g$	[h] < 1	%	CL=90%	—
$\gamma a_1^0 \rightarrow \gamma s \bar{s}$	[h] < 1	$\times 10^{-3}$	CL=90%	—

Lepton Family number (LF) violating modes

$\mu^\pm \tau^\mp$	LF	< 6.0	$\times 10^{-6}$	CL=95%	4563
--------------------	----	---------	------------------	--------	------

Other decays

invisible	< 3.0	$\times 10^{-4}$	CL=90%	—
-----------	---------	------------------	--------	---

$\chi_{b0}(1P)$ [i]

$I^G(J^{PC}) = 0^+(0^{++})$
J needs confirmation.

Mass $m = 9859.44 \pm 0.42 \pm 0.31 \text{ MeV}$

Mass $m = 32.5 \pm 0.9 \text{ MeV}$

$\chi_{b0}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	$\frac{P}{\text{MeV}/c}$
$\gamma \Upsilon(1S)$	(1.94 ± 0.27) %		†
$D^0 X$	< 10.4	%	90%
$\pi^+ \pi^- K^+ K^- \pi^0$	< 1.6	$\times 10^{-4}$	90%
$2\pi^+ \pi^- K^- K_S^0$	< 5	$\times 10^{-5}$	90%
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 5	$\times 10^{-4}$	90%
$2\pi^+ 2\pi^- 2\pi^0$	< 2.1	$\times 10^{-4}$	90%
$2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.6) $\times 10^{-4}$		†
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.7	$\times 10^{-4}$	90%

$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 5	$\times 10^{-4}$	90%	†
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 1.6	$\times 10^{-4}$	90%	†
$3\pi^+ 3\pi^-$	< 8	$\times 10^{-5}$	90%	†
$3\pi^+ 3\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	†
$3\pi^+ 3\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$			†
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 1.0	$\times 10^{-3}$	90%	†
$4\pi^+ 4\pi^-$	< 8	$\times 10^{-5}$	90%	†
$4\pi^+ 4\pi^- 2\pi^0$	< 2.1	$\times 10^{-3}$	90%	†
$J/\psi J/\psi$	< 7	$\times 10^{-5}$	90%	†
$J/\psi \psi(2S)$	< 1.2	$\times 10^{-4}$	90%	†
$\psi(2S) \psi(2S)$	< 3.1	$\times 10^{-5}$	90%	†

$\chi_{b1}(1P)$ [1]

$I^G(J^{PC}) = 0^+(1^{++})$
J needs confirmation.

Mass $m = 9892.78 \pm 0.26 \pm 0.31$ MeV

$\chi_{b1}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\gamma \Upsilon(1S)$	$(35.0 \pm 2.1) \%$		423
$D^0 X$	$(12.6 \pm 2.2) \%$		—
$\pi^+ \pi^- K^+ K^- \pi^0$	$(2.0 \pm 0.6) \times 10^{-4}$		4892
$2\pi^+ \pi^- K^- K_S^0$	$(1.3 \pm 0.5) \times 10^{-4}$		4892
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$< 6 \times 10^{-4}$	90%	4863
$2\pi^+ 2\pi^- 2\pi^0$	$(8.0 \pm 2.5) \times 10^{-4}$		4921
$2\pi^+ 2\pi^- K^+ K^-$	$(1.5 \pm 0.5) \times 10^{-4}$		4878
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(3.5 \pm 1.2) \times 10^{-4}$		4863
$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(8.6 \pm 3.2) \times 10^{-4}$		4845
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$(9.3 \pm 3.3) \times 10^{-4}$		4844
$3\pi^+ 3\pi^-$	$(1.9 \pm 0.6) \times 10^{-4}$		4921
$3\pi^+ 3\pi^- 2\pi^0$	$(1.7 \pm 0.5) \times 10^{-3}$		4898
$3\pi^+ 3\pi^- K^+ K^-$	$(2.6 \pm 0.8) \times 10^{-4}$		4844
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(7.5 \pm 2.6) \times 10^{-4}$		4825
$4\pi^+ 4\pi^-$	$(2.6 \pm 0.9) \times 10^{-4}$		4897
$4\pi^+ 4\pi^- 2\pi^0$	$(1.4 \pm 0.6) \times 10^{-3}$		4867
$J/\psi J/\psi$	$< 2.7 \times 10^{-5}$	90%	3857
$J/\psi \psi(2S)$	$< 1.7 \times 10^{-5}$	90%	3594
$\psi(2S) \psi(2S)$	$< 6 \times 10^{-5}$	90%	3298

$h_b(1P)$

$$I^G(J^{PC}) = ??(1^{+-})$$

Mass $m = 9899.3 \pm 0.8$ MeV

$h_b(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\eta_b(1S)\gamma$	$(52_{-5}^{+6})\%$	488

$\chi_{b2}(1P)$ [i]

$$I^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

Mass $m = 9912.21 \pm 0.26 \pm 0.31$ MeV

$\chi_{b2}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\gamma \Upsilon(1S)$	$(18.8 \pm 1.1)\%$		442
$D^0 X$	$< 7.9\%$	90%	—
$\pi^+ \pi^- K^+ K^- \pi^0$	$(8 \pm 5) \times 10^{-5}$		4902
$2\pi^+ \pi^- K^- K_S^0$	$< 1.0 \times 10^{-4}$	90%	4901
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$(5.3 \pm 2.4) \times 10^{-4}$		4873
$2\pi^+ 2\pi^- 2\pi^0$	$(3.5 \pm 1.4) \times 10^{-4}$		4931
$2\pi^+ 2\pi^- K^+ K^-$	$(1.1 \pm 0.4) \times 10^{-4}$		4888
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.1 \pm 0.9) \times 10^{-4}$		4872
$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(3.9 \pm 1.8) \times 10^{-4}$		4855
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 5 \times 10^{-4}$	90%	4854
$3\pi^+ 3\pi^-$	$(7.0 \pm 3.1) \times 10^{-5}$		4931
$3\pi^+ 3\pi^- 2\pi^0$	$(1.0 \pm 0.4) \times 10^{-3}$		4908
$3\pi^+ 3\pi^- K^+ K^-$	$< 8 \times 10^{-5}$	90%	4854
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(3.6 \pm 1.5) \times 10^{-4}$		4835
$4\pi^+ 4\pi^-$	$(8 \pm 4) \times 10^{-5}$		4907
$4\pi^+ 4\pi^- 2\pi^0$	$(1.8 \pm 0.7) \times 10^{-3}$		4877
$J/\psi J/\psi$	$< 4 \times 10^{-5}$	90%	3869
$J/\psi \psi(2S)$	$< 5 \times 10^{-5}$	90%	3608
$\psi(2S) \psi(2S)$	$< 1.6 \times 10^{-5}$	90%	3313

$\Upsilon(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 10023.26 \pm 0.31$ MeV

$m_{\Upsilon(3S)} - m_{\Upsilon(2S)} = 331.50 \pm 0.13$ MeV

Full width $\Gamma = 31.98 \pm 2.63$ keV

$\Gamma_{ee} = 0.612 \pm 0.011$ keV

$\Upsilon(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\Upsilon(1S)\pi^+\pi^-$	$(17.85 \pm 0.26) \%$		475
$\Upsilon(1S)\pi^0\pi^0$	$(8.6 \pm 0.4) \%$		480
$\tau^+\tau^-$	$(2.00 \pm 0.21) \%$		4686
$\mu^+\mu^-$	$(1.93 \pm 0.17) \%$	S=2.2	5011
e^+e^-	$(1.91 \pm 0.16) \%$		5012
$\Upsilon(1S)\pi^0$	< 4	$\times 10^{-5}$ CL=90%	531
$\Upsilon(1S)\eta$	$(2.9 \pm 0.4) \times 10^{-4}$	S=2.0	126
$J/\psi(1S)$ anything	< 6	$\times 10^{-3}$ CL=90%	4533
$J/\psi(1S)\eta_c$	< 5.4	$\times 10^{-6}$ CL=90%	3984
$J/\psi(1S)\chi_{c0}$	< 3.4	$\times 10^{-6}$ CL=90%	3808
$J/\psi(1S)\chi_{c1}$	< 1.2	$\times 10^{-6}$ CL=90%	3765
$J/\psi(1S)\chi_{c2}$	< 2.0	$\times 10^{-6}$ CL=90%	3744
$J/\psi(1S)\eta_c(2S)$	< 2.5	$\times 10^{-6}$ CL=90%	3706
$J/\psi(1S)X(3940)$	< 2.0	$\times 10^{-6}$ CL=90%	3555
$J/\psi(1S)X(4160)$	< 2.0	$\times 10^{-6}$ CL=90%	3440
χ_{c1} anything	$(2.2 \pm 0.5) \times 10^{-4}$		—
χ_{c2} anything	$(2.3 \pm 0.8) \times 10^{-4}$		—
$\psi(2S)\eta_c$	< 5.1	$\times 10^{-6}$ CL=90%	3732
$\psi(2S)\chi_{c0}$	< 4.7	$\times 10^{-6}$ CL=90%	3536
$\psi(2S)\chi_{c1}$	< 2.5	$\times 10^{-6}$ CL=90%	3488
$\psi(2S)\chi_{c2}$	< 1.9	$\times 10^{-6}$ CL=90%	3464
$\psi(2S)\eta_c(2S)$	< 3.3	$\times 10^{-6}$ CL=90%	3421
$\psi(2S)X(3940)$	< 3.9	$\times 10^{-6}$ CL=90%	3250
$\psi(2S)X(4160)$	< 3.9	$\times 10^{-6}$ CL=90%	3118
2H anything	$(2.78^{+0.30}_{-0.26}) \times 10^{-5}$	S=1.2	—
hadrons	$(94 \pm 11) \%$		—
ggg	$(58.8 \pm 1.2) \%$		—
γgg	$(1.87 \pm 0.28) \%$		—
ϕK^+K^-	$(1.6 \pm 0.4) \times 10^{-6}$		4910
$\omega\pi^+\pi^-$	< 2.58	$\times 10^{-6}$ CL=90%	4977
$K^*(892)^0 K^-\pi^+ + c.c.$	$(2.3 \pm 0.7) \times 10^{-6}$		4952
$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$ CL=90%	4841
$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$ CL=90%	4899
$\rho(770)a_2(1320)$	< 8.8	$\times 10^{-7}$ CL=90%	4894
$K^*(892)^0 \bar{K}_2^*(1430)^0 + c.c.$	$(1.5 \pm 0.6) \times 10^{-6}$		4869
$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$ CL=90%	4918
$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$ CL=90%	4901
$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$ CL=90%	4935
$\rho\pi$	< 1.16	$\times 10^{-6}$ CL=90%	4981
$\pi^+\pi^-\pi^0$	< 8.0	$\times 10^{-7}$ CL=90%	5007
$\omega\pi^0$	< 1.63	$\times 10^{-6}$ CL=90%	4980

$\pi^+\pi^-\pi^0\pi^0$	$(1.30 \pm 0.28) \times 10^{-5}$	5002
$K_S^0 K^+\pi^- + \text{c.c.}$	$(1.14 \pm 0.33) \times 10^{-6}$	4979
$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$< 4.22 \times 10^{-6}$	CL=90% 4959
$K^*(892)^- K^+ + \text{c.c.}$	$< 1.45 \times 10^{-6}$	CL=90% 4960
Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$	—

Radiative decays

$\gamma\chi_{b1}(1P)$	$(6.9 \pm 0.4) \%$	130
$\gamma\chi_{b2}(1P)$	$(7.15 \pm 0.35) \%$	110
$\gamma\chi_{b0}(1P)$	$(3.8 \pm 0.4) \%$	5012
$\gamma f_0(1710)$	$< 5.9 \times 10^{-4}$	CL=90% 4864
$\gamma f_2'(1525)$	$< 5.3 \times 10^{-4}$	CL=90% 4896
$\gamma f_2(1270)$	$< 2.41 \times 10^{-4}$	CL=90% 4930
$\gamma\eta_c(1S)$	$< 2.7 \times 10^{-5}$	CL=90% 4568
$\gamma\chi_{c0}$	$< 1.0 \times 10^{-4}$	CL=90% 4430
$\gamma\chi_{c1}$	$< 3.6 \times 10^{-6}$	CL=90% 4397
$\gamma\chi_{c2}$	$< 1.5 \times 10^{-5}$	CL=90% 4381
$\gamma X(3872) \rightarrow \pi^+\pi^- J/\psi$	$< 8 \times 10^{-7}$	CL=90% —
$\gamma X(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi$	$< 2.4 \times 10^{-6}$	CL=90% —
$\gamma X(3915) \rightarrow \omega J/\psi$	$< 2.8 \times 10^{-6}$	CL=90% —
$\gamma X(4140) \rightarrow \phi J/\psi$	$< 1.2 \times 10^{-6}$	CL=90% —
$\gamma X(4350) \rightarrow \phi J/\psi$	$< 1.3 \times 10^{-6}$	CL=90% —
$\gamma\eta_b(1S)$	$(3.9 \pm 1.5) \times 10^{-4}$	605
$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	$< 3.7 \times 10^{-6}$	CL=90% —
$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	$< 4.9 \times 10^{-6}$	CL=90% —
$\gamma X \rightarrow \gamma + \geq 4$ prongs	$[j] < 1.95 \times 10^{-4}$	CL=95% —
$\gamma A^0 \rightarrow \gamma$ hadrons	$< 8 \times 10^{-5}$	CL=90% —
$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	$< 8.3 \times 10^{-6}$	CL=90% —

Lepton Family number (LF) violating modes

$e^\pm \tau^\mp$	LF	$< 3.2 \times 10^{-6}$	CL=90%	4854
$\mu^\pm \tau^\mp$	LF	$< 3.3 \times 10^{-6}$	CL=90%	4854

$\Upsilon(1D)$

$$I^G(J^{PC}) = 0^-(2^{--})$$

$$\text{Mass } m = 10163.7 \pm 1.4 \text{ MeV} \quad (S = 1.7)$$

$\Upsilon(1D)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma\gamma \Upsilon(1S)$	seen	679
$\gamma\chi_{bJ}(1P)$	seen	5082
$\eta \Upsilon(1S)$	not seen	426
$\pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$	623

$\chi_{b0}(2P)$ [i]

$$J^G(J^{PC}) = 0^+(0^{++})$$

J needs confirmation.

$$\text{Mass } m = 10232.5 \pm 0.4 \pm 0.5 \text{ MeV}$$

$$\text{Mass } m = 23.8 \pm 1.7 \text{ MeV}$$

$\chi_{b0}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\gamma \Upsilon(2S)$	$(1.38 \pm 0.30) \%$		†
$\gamma \Upsilon(1S)$	$(3.8 \pm 1.7) \times 10^{-3}$		†
$D^0 X$	$< 8.2 \%$	90%	—
$\pi^+ \pi^- K^+ K^- \pi^0$	$< 3.4 \times 10^{-5}$	90%	†
$2\pi^+ \pi^- K^- K_S^0$	$< 5 \times 10^{-5}$	90%	†
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$< 2.2 \times 10^{-4}$	90%	†
$2\pi^+ 2\pi^- 2\pi^0$	$< 2.4 \times 10^{-4}$	90%	†
$2\pi^+ 2\pi^- K^+ K^-$	$< 1.5 \times 10^{-4}$	90%	†
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$< 2.2 \times 10^{-4}$	90%	†
$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$< 1.1 \times 10^{-3}$	90%	†
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 7 \times 10^{-4}$	90%	†
$3\pi^+ 3\pi^-$	$< 7 \times 10^{-5}$	90%	†
$3\pi^+ 3\pi^- 2\pi^0$	$< 1.2 \times 10^{-3}$	90%	†
$3\pi^+ 3\pi^- K^+ K^-$	$< 1.5 \times 10^{-4}$	90%	†
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$< 7 \times 10^{-4}$	90%	†
$4\pi^+ 4\pi^-$	$< 1.7 \times 10^{-4}$	90%	†
$4\pi^+ 4\pi^- 2\pi^0$	$< 6 \times 10^{-4}$	90%	†

$\chi_{b1}(2P)$ [i]

$$J^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

$$\text{Mass } m = 10255.46 \pm 0.22 \pm 0.50 \text{ MeV}$$

$$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)} = 23.5 \pm 1.0 \text{ MeV}$$

$\chi_{b1}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\omega \Upsilon(1S)$	$(1.63^{+0.40}_{-0.34}) \%$	135
$\gamma \Upsilon(2S)$	$(18.1 \pm 1.9) \%$	230
$\gamma \Upsilon(1S)$	$(9.9 \pm 1.0) \%$	764
$\pi\pi \chi_{b1}(1P)$	$(9.1 \pm 1.3) \times 10^{-3}$	238
$D^0 X$	$(8.8 \pm 1.7) \%$	—
$\pi^+ \pi^- K^+ K^- \pi^0$	$(3.1 \pm 1.0) \times 10^{-4}$	5075
$2\pi^+ \pi^- K^- K_S^0$	$(1.1 \pm 0.5) \times 10^{-4}$	5075
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$(7.7 \pm 3.2) \times 10^{-4}$	5047
$2\pi^+ 2\pi^- 2\pi^0$	$(5.9 \pm 2.0) \times 10^{-4}$	5104

$2\pi^+ 2\pi^- K^+ K^-$	$(10 \pm 4) \times 10^{-5}$	5062
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(5.5 \pm 1.8) \times 10^{-4}$	5047
$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(10 \pm 4) \times 10^{-4}$	5030
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$(6.7 \pm 2.6) \times 10^{-4}$	5029
$3\pi^+ 3\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	5103
$3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	5081
$3\pi^+ 3\pi^- K^+ K^-$	$(2.0 \pm 0.8) \times 10^{-4}$	5029
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(6.1 \pm 2.2) \times 10^{-4}$	5011
$4\pi^+ 4\pi^-$	$(1.7 \pm 0.6) \times 10^{-4}$	5080
$4\pi^+ 4\pi^- 2\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$	5051

$\chi_{b2}(2P)$ [i]

$$I^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

$$\text{Mass } m = 10268.65 \pm 0.22 \pm 0.50 \text{ MeV}$$

$$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)} = 13.10 \pm 0.24 \text{ MeV}$$

$\chi_{b2}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\omega \Upsilon(1S)$	$(1.10^{+0.34}_{-0.30})\%$		194
$\gamma \Upsilon(2S)$	$(8.9 \pm 1.2)\%$		242
$\gamma \Upsilon(1S)$	$(6.6 \pm 0.8)\%$		777
$\pi\pi \chi_{b2}(1P)$	$(5.1 \pm 0.9) \times 10^{-3}$		229
$D^0 X$	$< 2.4\%$	90%	—
$\pi^+ \pi^- K^+ K^- \pi^0$	$< 1.1 \times 10^{-4}$	90%	5082
$2\pi^+ \pi^- K^- K_S^0$	$< 9 \times 10^{-5}$	90%	5082
$2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$< 7 \times 10^{-4}$	90%	5054
$2\pi^+ 2\pi^- 2\pi^0$	$(3.9 \pm 1.6) \times 10^{-4}$		5110
$2\pi^+ 2\pi^- K^+ K^-$	$(9 \pm 4) \times 10^{-5}$		5068
$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.4 \pm 1.1) \times 10^{-4}$		5054
$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(4.7 \pm 2.3) \times 10^{-4}$		5037
$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 4 \times 10^{-4}$	90%	5036
$3\pi^+ 3\pi^-$	$(9 \pm 4) \times 10^{-5}$		5110
$3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$		5088
$3\pi^+ 3\pi^- K^+ K^-$	$(1.4 \pm 0.7) \times 10^{-4}$		5036
$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(4.2 \pm 1.7) \times 10^{-4}$		5017
$4\pi^+ 4\pi^-$	$(9 \pm 5) \times 10^{-5}$		5087
$4\pi^+ 4\pi^- 2\pi^0$	$(1.3 \pm 0.5) \times 10^{-3}$		5058

$\Upsilon(3S)$

$$J^{PC} = 0^{-}(1^{- -})$$

Mass $m = 10355.2 \pm 0.5$ MeV
 $m_{\Upsilon(3S)} - m_{\Upsilon(2S)} = 331.50 \pm 0.13$ MeV
 Full width $\Gamma = 20.32 \pm 1.85$ keV
 $\Gamma_{ee} = 0.443 \pm 0.008$ keV

$\Upsilon(3S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\Upsilon(2S)$ anything	(10.6 \pm 0.8) %		296
$\Upsilon(2S)\pi^+\pi^-$	(2.82 \pm 0.18) %	S=1.6	177
$\Upsilon(2S)\pi^0\pi^0$	(1.85 \pm 0.14) %		190
$\Upsilon(2S)\gamma\gamma$	(5.0 \pm 0.7) %		327
$\Upsilon(2S)\pi^0$	< 5.1	$\times 10^{-4}$ CL=90%	298
$\Upsilon(1S)\pi^+\pi^-$	(4.37 \pm 0.08) %		813
$\Upsilon(1S)\pi^0\pi^0$	(2.20 \pm 0.13) %		816
$\Upsilon(1S)\eta$	< 1	$\times 10^{-4}$ CL=90%	677
$\Upsilon(1S)\pi^0$	< 7	$\times 10^{-5}$ CL=90%	846
$h_b(1P)\pi^0$	< 1.2	$\times 10^{-3}$ CL=90%	426
$h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 \pm 1.4) $\times 10^{-4}$		—
$h_b(1P)\pi^+\pi^-$	< 1.2	$\times 10^{-4}$ CL=90%	353
$\tau^+\tau^-$	(2.29 \pm 0.30) %		4863
$\mu^+\mu^-$	(2.18 \pm 0.21) %	S=2.1	5177
e^+e^-	(2.18 \pm 0.20) %		5178
hadrons	(93 \pm 12) %		—
ggg	(35.7 \pm 2.6) %		—
γgg	(9.7 \pm 1.8) $\times 10^{-3}$		—
2H anything	(2.33 \pm 0.33) $\times 10^{-5}$		—

Radiative decays

$\gamma\chi_{b2}(2P)$	(13.1 \pm 1.6) %	S=3.4	86
$\gamma\chi_{b1}(2P)$	(12.6 \pm 1.2) %	S=2.4	99
$\gamma\chi_{b0}(2P)$	(5.9 \pm 0.6) %	S=1.4	5178
$\gamma\chi_{b2}(1P)$	(9.9 \pm 1.2) $\times 10^{-3}$	S=1.9	434
$\gamma\chi_{b1}(1P)$	(9 \pm 5) $\times 10^{-4}$	S=1.8	452
$\gamma\chi_{b0}(1P)$	(2.7 \pm 0.4) $\times 10^{-3}$		5178
$\gamma\eta_b(2S)$	< 6.2	$\times 10^{-4}$ CL=90%	350
$\gamma\eta_b(1S)$	(5.1 \pm 0.7) $\times 10^{-4}$		912
$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$ CL=90%	—
$\gamma X \rightarrow \gamma + \geq 4$ prongs	[k] < 2.2	$\times 10^{-4}$ CL=95%	—
$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5	$\times 10^{-6}$ CL=90%	—
$\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[l] < 1.6	$\times 10^{-4}$ CL=90%	—

Lepton Family number (*LF*) violating modes

$e^\pm \tau^\mp$	<i>LF</i>	< 4.2	$\times 10^{-6}$	CL=90%	5025
$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.1	$\times 10^{-6}$	CL=90%	5025

$\chi_{b1}(3P)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

Mass $m = 10512.1 \pm 2.3$ MeV

$\chi_{b1}(3P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Upsilon(1S)\gamma$	seen	999
$\Upsilon(2S)\gamma$	seen	477
$\Upsilon(3S)\gamma$	seen	156

**$\Upsilon(4S)$
or $\Upsilon(10580)$**

$$J^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 10579.4 \pm 1.2$ MeV

Full width $\Gamma = 20.5 \pm 2.5$ MeV

$\Gamma_{ee} = 0.272 \pm 0.029$ keV ($S = 1.5$)

$\Upsilon(4S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$B\bar{B}$	> 96 %	95%	326
B^+B^-	(51.4 \pm 0.6) %		331
D_s^+ anything + c.c.	(17.8 \pm 2.6) %		—
$B^0\bar{B}^0$	(48.6 \pm 0.6) %		326
$J/\psi K_S^0 + (J/\psi, \eta_c)K_S^0$	< 4 $\times 10^{-7}$	90%	—
non- $B\bar{B}$	< 4 %	95%	—
e^+e^-	(1.57 \pm 0.08) $\times 10^{-5}$		5290
$\rho^+\rho^-$	< 5.7 $\times 10^{-6}$	90%	5233
$K^*(892)^0\bar{K}^0$	< 2.0 $\times 10^{-6}$	90%	5240
$J/\psi(1S)$ anything	< 1.9 $\times 10^{-4}$	95%	—
D^{*+} anything + c.c.	< 7.4 %	90%	5099
ϕ anything	(7.1 \pm 0.6) %		5240
$\phi\eta$	< 1.8 $\times 10^{-6}$	90%	5226
$\phi\eta'$	< 4.3 $\times 10^{-6}$	90%	5196
$\rho\eta$	< 1.3 $\times 10^{-6}$	90%	5247
$\rho\eta'$	< 2.5 $\times 10^{-6}$	90%	5217
$\Upsilon(1S)$ anything	< 4 $\times 10^{-3}$	90%	1053
$\Upsilon(1S)\pi^+\pi^-$	(8.1 \pm 0.6) $\times 10^{-5}$		1026
$\Upsilon(1S)\eta$	(1.96 \pm 0.28) $\times 10^{-4}$		924

$\Upsilon(2S)\pi^+\pi^-$	$(8.6 \pm 1.3) \times 10^{-5}$	468
$h_b(1P)\pi^+\pi^-$	not seen	600
$h_b(1P)\eta$	$(2.18 \pm 0.21) \times 10^{-3}$	390
2H anything	$< 1.3 \times 10^{-5}$	90% —

X(10610)[±]

$$I^G(J^P) = 1^+(1^+)$$

Mass $m = 10607.2 \pm 2.0$ MeV

Full width $\Gamma = 18.4 \pm 2.4$ MeV

X(10610)[−] decay modes are charge conjugates of the modes below.

X(10610)⁺ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Upsilon(1S)\pi^+$	$(5.4^{+1.9}_{-1.5}) \times 10^{-3}$	1077
$\Upsilon(2S)\pi^+$	$(3.6^{+1.1}_{-0.8})\%$	551
$\Upsilon(3S)\pi^+$	$(2.1^{+0.8}_{-0.6})\%$	207
$h_b(1P)\pi^+$	$(3.5^{+1.2}_{-0.9})\%$	671
$h_b(2P)\pi^+$	$(4.7^{+1.7}_{-1.3})\%$	313
$B^+\bar{B}^0$	not seen	505
$B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	$(85.6^{+2.1}_{-2.9})\%$	—
$B^{*+}\bar{B}^{*0}$	not seen	†

X(10610)⁰

$$I^G(J^P) = 1^+(1^+)$$

Mass $m = 10609 \pm 6$ MeV

X(10610)⁰ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Upsilon(1S)\pi^0$	not seen	1079
$\Upsilon(2S)\pi^0$	seen	554
$\Upsilon(3S)\pi^0$	seen	212

$\Upsilon(10860)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 10889.9^{+3.2}_{-2.6}$ MeV

Full width $\Gamma = 51^{+6}_{-7}$ MeV

$\Gamma_{ee} = 0.31 \pm 0.07$ keV ($S = 1.3$)

$T(10860)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$B\bar{B}X$	(76.2 $^{+2.7}_{-4.0}$) %		–
$B\bar{B}$	(5.5 ± 1.0) %		1332
$B\bar{B}^* + \text{c.c.}$	(13.7 ± 1.6) %		–
$B^*\bar{B}^*$	(38.1 ± 3.4) %		1138
$B\bar{B}^{(*)}\pi$	< 19.7 %	90%	1027
$B\bar{B}\pi$	(0.0 ± 1.2) %		1027
$B^*\bar{B}\pi + B\bar{B}^*\pi$	(7.3 ± 2.3) %		–
$B^*\bar{B}^*\pi$	(1.0 ± 1.4) %		756
$B\bar{B}\pi\pi$	< 8.9 %	90%	574
$B_s^{(*)}\bar{B}_s^{(*)}$	(20.1 ± 3.1) %		919
$B_s\bar{B}_s$	(5 ± 5) $\times 10^{-3}$		919
$B_s\bar{B}_s^* + \text{c.c.}$	(1.35 ± 0.32) %		–
$B_s^*\bar{B}_s^*$	(17.6 ± 2.7) %		566
no open-bottom	(3.8 $^{+5.0}_{-0.5}$) %		–
e^+e^-	(6.1 ± 1.6) $\times 10^{-6}$		5445
$K^*(892)^0\bar{K}^0$	< 1.0 $\times 10^{-5}$	90%	5397
$\Upsilon(1S)\pi^+\pi^-$	(5.3 ± 0.6) $\times 10^{-3}$		1310
$\Upsilon(2S)\pi^+\pi^-$	(7.8 ± 1.3) $\times 10^{-3}$		788
$\Upsilon(3S)\pi^+\pi^-$	(4.8 $^{+1.9}_{-1.7}$) $\times 10^{-3}$		445
$\Upsilon(1S)K^+K^-$	(6.1 ± 1.8) $\times 10^{-4}$		965
$h_b(1P)\pi^+\pi^-$	(3.5 $^{+1.0}_{-1.3}$) $\times 10^{-3}$		907
$h_b(2P)\pi^+\pi^-$	(5.7 $^{+1.7}_{-2.1}$) $\times 10^{-3}$		548
$\chi_{b0}(1P)\pi^+\pi^-\pi^0$	< 6.3 $\times 10^{-3}$	90%	5440
$\chi_{b0}(1P)\omega$	< 3.9 $\times 10^{-3}$	90%	5417
$\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	< 4.8 $\times 10^{-3}$	90%	–
$\chi_{b1}(1P)\pi^+\pi^-\pi^0$	(1.85 ± 0.33) $\times 10^{-3}$		865
$\chi_{b1}(1P)\omega$	(1.57 ± 0.30) $\times 10^{-3}$		589
$\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	(5.2 ± 1.9) $\times 10^{-4}$		–
$\chi_{b2}(1P)\pi^+\pi^-\pi^0$	(1.17 ± 0.30) $\times 10^{-3}$		846
$\chi_{b2}(1P)\omega$	(6.0 ± 2.7) $\times 10^{-4}$		559
$\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	(6 ± 4) $\times 10^{-4}$		–
$\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega$	< 3.8 $\times 10^{-5}$	90%	–

Inclusive Decays.

These decay modes are submodes of one or more of the decay modes above.

ϕ anything	(13.8 $^{+2.4}_{-1.7}$) %		–
D^0 anything + c.c.	(108 ± 8) %		–

D_s anything + c.c.	(46 ± 6) %	—
J/ψ anything	(2.06 ± 0.21) %	—
B^0 anything + c.c.	(77 ± 8) %	—
B^+ anything + c.c.	(72 ± 6) %	—

$\Upsilon(11020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$$\text{Mass } m = 10992.9^{+10.0}_{-3.1} \text{ MeV}$$

$$\text{Full width } \Gamma = 49^{+9}_{-15} \text{ MeV}$$

$$\Gamma_{ee} = 0.130 \pm 0.030 \text{ keV}$$

$\Upsilon(11020)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$e^+ e^-$	$(2.7^{+1.0}_{-0.8}) \times 10^{-6}$	5496

NOTES

[a] $2m_\tau < M(\tau^+ \tau^-) < 9.2 \text{ GeV}$

[b] $2 \text{ GeV} < m_{K^+ K^-} < 3 \text{ GeV}$

[c] $X = \text{scalar with } m < 8.0 \text{ GeV}$

[d] $X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV}$

[e] X and $\bar{X} = \text{zero spin with } m < 4.5 \text{ GeV}$

[f] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[g] $201 \text{ MeV} < M(\mu^+ \mu^-) < 3565 \text{ MeV}$

[h] $0.5 \text{ GeV} < m_X < 9.0 \text{ GeV}$, where m_X is the invariant mass of the hadronic final state.

[i] Spectroscopic labeling for these states is theoretical, pending experimental information.

[j] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[k] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[l] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV.