



$$I(J^P) = 0(0^-)$$

$I, J, P$  need confirmation. Quantum numbers shown are quark-model predictions.

### $B_s^0$ MASS

| VALUE (MeV)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | EVTS | DOCUMENT ID         | TECN      | COMMENT                           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|---------------------|-----------|-----------------------------------|
| <b>5366.88 ± 0.14 OUR FIT</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |      |                     |           |                                   |
| <b>5366.84 ± 0.15 OUR AVERAGE</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |      |                     |           |                                   |
| 5366.85 ± 0.19 ± 0.13                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |      | <sup>1</sup> AAIJ   | 19U LHCb  | $pp$ at 7, 8, 13 TeV              |
| 5366.83 ± 0.25 ± 0.27                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |      | <sup>2</sup> AAIJ   | 18AC LHCb | $pp$ at 7, 8, 13 TeV              |
| 5367.08 ± 0.38 ± 0.15                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | 128  | <sup>3</sup> AAIJ   | 16U LHCb  | $pp$ at 7, 8 TeV                  |
| 5366.90 ± 0.28 ± 0.23                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |      | <sup>4</sup> AAIJ   | 12E LHCb  | $pp$ at 7 TeV                     |
| 5364.4 ± 1.3 ± 0.7                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |      | LOUVOT              | 09 BELL   | $e^+e^- \rightarrow \Upsilon(5S)$ |
| 5366.01 ± 0.73 ± 0.33                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |      | <sup>5</sup> ACOSTA | 06 CDF    | $p\bar{p}$ at 1.96 TeV            |
| 5369.9 ± 2.3 ± 1.3                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 32   | <sup>6</sup> ABE    | 96B CDF   | $p\bar{p}$ at 1.8 TeV             |
| 5374 ± 16 ± 2                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 3    | ABREU               | 94D DLPH  | $e^+e^- \rightarrow Z$            |
| 5359 ± 19 ± 7                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | 1    | <sup>6</sup> AKERS  | 94J OPAL  | $e^+e^- \rightarrow Z$            |
| 5368.6 ± 5.6 ± 1.5                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 2    | BUSKULIC            | 93G ALEP  | $e^+e^- \rightarrow Z$            |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |      |                     |           |                                   |
| 5370 ± 1 ± 3                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |      | DRUTSKOY            | 07A BELL  | Repl. by LOUVOT 09                |
| 5370 ± 40                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | 6    | <sup>7</sup> AKERS  | 94J OPAL  | $e^+e^- \rightarrow Z$            |
| 5383.3 ± 4.5 ± 5.0                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | 14   | ABE                 | 93F CDF   | Repl. by ABE 96B                  |
| <sup>1</sup> Uses $B_s^0 \rightarrow J/\psi p\bar{p}$ decays.<br><sup>2</sup> Uses $B_s \rightarrow \chi_{c1} K^+ K^-$ mode.<br><sup>3</sup> Uses $J/\psi \rightarrow \mu^+ \mu^-$ , $\phi \rightarrow K^+ K^-$ decays, and observes $128 \pm 13$ events of $B_s^0 \rightarrow J/\psi \phi \phi$ .<br><sup>4</sup> Uses $B_s^0 \rightarrow J/\psi \phi$ fully reconstructed decays.<br><sup>5</sup> Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+ \mu^-$ decays.<br><sup>6</sup> From the decay $B_s \rightarrow J/\psi(1S) \phi$ .<br><sup>7</sup> From the decay $B_s \rightarrow D_s^- \pi^+$ . |      |                     |           |                                   |

### $m_{B_s^0} - m_B$

$m_B$  is the average of our  $B$  masses  $(m_{B^\pm} + m_{B^0})/2$ .

| VALUE (MeV)                                                          | CL% | DOCUMENT ID         | TECN     | COMMENT                |
|----------------------------------------------------------------------|-----|---------------------|----------|------------------------|
| <b>87.38 ± 0.16 OUR FIT</b>                                          |     |                     |          |                        |
| <b>87.42 ± 0.24 OUR AVERAGE</b>                                      |     |                     |          |                        |
| 87.60 ± 0.44 ± 0.09                                                  |     | <sup>1</sup> AAIJ   | 15U LHCb | $pp$ at 7, 8 TeV       |
| 87.42 ± 0.30 ± 0.09                                                  |     | <sup>2</sup> AAIJ   | 12E LHCb | $pp$ at 7 TeV          |
| 86.64 ± 0.80 ± 0.08                                                  |     | <sup>3</sup> ACOSTA | 06 CDF   | $p\bar{p}$ at 1.96 TeV |
| • • • We use the following data for averages but not for fits. • • • |     |                     |          |                        |
| 89.7 ± 2.7 ± 1.2                                                     |     | ABE                 | 96B CDF  | $p\bar{p}$ at 1.8 TeV  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

80 to 130 68 LEE-FRANZINI 90 CSB2  $e^+e^- \rightarrow \Upsilon(5S)$

<sup>1</sup> The reported result is  $m_{B_s^0} - m_{B^0} = 87.45 \pm 0.44 \pm 0.09$  MeV. We convert it to the mass difference with respect to the average of  $(m_{B^\pm} + m_{B^0})/2$ . Uses the mode  $B_s^0 \rightarrow \psi(2S) K^- \pi^+$ .

<sup>2</sup> The reported result is  $m_{B_s^0} - m_{B^+} = 87.52 \pm 0.30 \pm 0.12$  MeV. We convert it to the mass difference with respect to the average of  $(m_{B^\pm} + m_{B^0})/2$ .

<sup>3</sup> The reported result is  $m_{B_s^0} - m_{B^0} = 86.38 \pm 0.90 \pm 0.06$  MeV. We convert it to the mass difference with respect to the average of  $(m_{B^\pm} + m_{B^0})/2$ .

$$m_{B_{sH}^0} - m_{B_{sL}^0}$$

See the  $B_s^0\text{-}\bar{B}_s^0$  MIXING section near the end of these  $B_s^0$  Listings.

## $B_s^0$ MEAN LIFE

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>). It is derived from the average of  $\Gamma_{B_s^0}$ .

| VALUE ( $10^{-12}$ s)                                                         | EVTS | DOCUMENT ID             | TECN     | COMMENT                |
|-------------------------------------------------------------------------------|------|-------------------------|----------|------------------------|
| <b>1.515<math>\pm</math>0.004 OUR EVALUATION</b>                              |      |                         |          |                        |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                         |          |                        |
| 1.518 $\pm$ 0.041 $\pm$ 0.027                                                 |      | <sup>1</sup> AALTONEN   | 11AP CDF | $p\bar{p}$ at 1.96 TeV |
| 1.398 $\pm$ 0.044 $^{+0.028}_{-0.025}$                                        |      | <sup>2</sup> ABAZOV     | 06V D0   | $p\bar{p}$ at 1.96 TeV |
| 1.42 $^{+0.14}_{-0.13}$ $\pm$ 0.03                                            |      | <sup>3</sup> ABREU      | 00Y DLPH | $e^+e^- \rightarrow Z$ |
| 1.53 $^{+0.16}_{-0.15}$ $\pm$ 0.07                                            |      | <sup>4</sup> ABREU,P    | 00G DLPH | $e^+e^- \rightarrow Z$ |
| 1.36 $\pm$ 0.09 $^{+0.06}_{-0.05}$                                            |      | <sup>5</sup> ABE        | 99D CDF  | $p\bar{p}$ at 1.8 TeV  |
| 1.72 $^{+0.20}_{-0.19}$ $^{+0.18}_{-0.17}$                                    |      | <sup>6</sup> ACKERSTAFF | 98F OPAL | $e^+e^- \rightarrow Z$ |
| 1.50 $^{+0.16}_{-0.15}$ $\pm$ 0.04                                            |      | <sup>5</sup> ACKERSTAFF | 98G OPAL | $e^+e^- \rightarrow Z$ |
| 1.47 $\pm$ 0.14 $\pm$ 0.08                                                    |      | <sup>4</sup> BARATE     | 98C ALEP | $e^+e^- \rightarrow Z$ |
| 1.51 $\pm$ 0.11                                                               |      | <sup>7</sup> BARATE     | 98C ALEP | $e^+e^- \rightarrow Z$ |
| 1.56 $^{+0.29}_{-0.26}$ $^{+0.08}_{-0.07}$                                    |      | <sup>5</sup> ABREU      | 96F DLPH | Repl. by ABREU 00Y     |
| 1.65 $^{+0.34}_{-0.31}$ $\pm$ 0.12                                            |      | <sup>4</sup> ABREU      | 96F DLPH | Repl. by ABREU 00Y     |
| 1.76 $\pm$ 0.20 $^{+0.15}_{-0.10}$                                            |      | <sup>8</sup> ABREU      | 96F DLPH | Repl. by ABREU 00Y     |
| 1.60 $\pm$ 0.26 $^{+0.13}_{-0.15}$                                            |      | <sup>9</sup> ABREU      | 96F DLPH | Repl. by ABREU,P 00G   |
| 1.67 $\pm$ 0.14                                                               |      | <sup>10</sup> ABREU     | 96F DLPH | $e^+e^- \rightarrow Z$ |
| 1.61 $^{+0.30}_{-0.29}$ $^{+0.18}_{-0.16}$                                    | 90   | <sup>4</sup> BUSKULIC   | 96E ALEP | Repl. by BARATE 98C    |

|      |                    |            |     |                       |     |      |                          |
|------|--------------------|------------|-----|-----------------------|-----|------|--------------------------|
| 1.54 | $+0.14$<br>$-0.13$ | $\pm 0.04$ |     | <sup>5</sup> BUSKULIC | 96M | ALEP | $e^+e^- \rightarrow Z$   |
| 1.42 | $+0.27$<br>$-0.23$ | $\pm 0.11$ | 76  | <sup>5</sup> ABE      | 95R | CDF  | Repl. by ABE 99D         |
| 1.74 | $+1.08$<br>$-0.69$ | $\pm 0.07$ | 8   | <sup>11</sup> ABE     | 95R | CDF  | Sup. by ABE 96N          |
| 1.54 | $+0.25$<br>$-0.21$ | $\pm 0.06$ | 79  | <sup>5</sup> AKERS    | 95G | OPAL | Repl. by ACKER-STAFF 98G |
| 1.59 | $+0.17$<br>$-0.15$ | $\pm 0.03$ | 134 | <sup>5</sup> BUSKULIC | 95O | ALEP | Sup. by BUSKULIC 96M     |
| 0.96 | $\pm 0.37$         |            | 41  | <sup>12</sup> ABREU   | 94E | DLPH | Sup. by ABREU 96F        |
| 1.92 | $+0.45$<br>$-0.35$ | $\pm 0.04$ | 31  | <sup>5</sup> BUSKULIC | 94C | ALEP | Sup. by BUSKULIC 95O     |
| 1.13 | $+0.35$<br>$-0.26$ | $\pm 0.09$ | 22  | <sup>5</sup> ACTON    | 93H | OPAL | Sup. by AKERS 95G        |

<sup>1</sup> AALTONEN 11AP combines the fully reconstructed  $B_s^0 \rightarrow D_s^- \pi^+$  decays and partially reconstructed  $B_s^0 \rightarrow D_s X$  decays.

<sup>2</sup> Measured using  $D_s \mu^+$  vertices.

<sup>3</sup> Uses  $D_s^- \ell^+$ , and  $\phi \ell^+$  vertices.

<sup>4</sup> Measured using  $D_s$  hadron vertices.

<sup>5</sup> Measured using  $D_s^- \ell^+$  vertices.

<sup>6</sup> ACKERSTAFF 98F use fully reconstructed  $D_s^- \rightarrow \phi \pi^-$  and  $D_s^- \rightarrow K^{*0} K^-$  in the inclusive  $B_s^0$  decay.

<sup>7</sup> Combined results from  $D_s^- \ell^+$  and  $D_s$  hadron.

<sup>8</sup> Measured using  $\phi \ell$  vertices.

<sup>9</sup> Measured using inclusive  $D_s$  vertices.

<sup>10</sup> Combined result for the four ABREU 96F methods.

<sup>11</sup> Exclusive reconstruction of  $B_s \rightarrow \psi \phi$ .

<sup>12</sup> ABREU 94E uses the flight-distance distribution of  $D_s$  vertices,  $\phi$ -lepton vertices, and  $D_s \mu$  vertices.

## $\Gamma_{B_s^0}$

"OUR EVALUATION" is an average performed by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) as described in our "Review on  $B$ - $\bar{B}$  Mixing" in the  $B^0$  section of these Listings. It includes the measurements of  $\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  listed in this section, as well as constraints from effective lifetimes with pure  $CP$  modes and flavor-specific modes.

| VALUE ( $10^{12} \text{ s}^{-1}$ )                   | DOCUMENT ID                                                 | TECN      | COMMENT                |
|------------------------------------------------------|-------------------------------------------------------------|-----------|------------------------|
| <b><math>0.6600 \pm 0.0016</math> OUR EVALUATION</b> |                                                             |           |                        |
| <b><math>0.661 \pm 0.004</math> OUR AVERAGE</b>      | Error includes scale factor of 2.2. See the ideogram below. |           |                        |
| $0.6563 \pm 0.0021$                                  | <sup>1</sup> AAIJ                                           | 19Q LHCb  | $pp$ at 7, 8, 13 TeV   |
| $0.675 \pm 0.003 \pm 0.003$                          | <sup>2</sup> AAD                                            | 16AP ATLS | $pp$ at 7, 8 TeV       |
| $0.668 \pm 0.011 \pm 0.006$                          | <sup>3</sup> AAIJ                                           | 16AK LHCb | $pp$ at 7, 8 TeV       |
| $0.6704 \pm 0.0043 \pm 0.0055$                       | <sup>2</sup> KHACHATRY...                                   | 16S CMS   | $pp$ at 8 TeV          |
| $0.654 \pm 0.008 \pm 0.004$                          | <sup>2</sup> AALTONEN                                       | 12AJ CDF  | $p\bar{p}$ at 1.96 TeV |
| $0.693 \pm 0.018$<br>$-0.017$                        | <sup>2</sup> ABAZOV                                         | 12D D0    | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                          |                         |        |      |                        |
|------------------------------------------|-------------------------|--------|------|------------------------|
| $0.650 \pm 0.006 \pm 0.004$              | <sup>1</sup> AAIJ       | 17V    | LHCB | Repl. by AAIJ 19Q      |
| $0.6603 \pm 0.0027 \pm 0.0015$           | <sup>4</sup> AAIJ       | 15I    | LHCB | Repl. by AAIJ 19Q      |
| $0.677 \pm 0.007 \pm 0.004$              | <sup>2</sup> AAD        | 14U    | ATLS | Repl. by AAD 16AP      |
| $0.661 \pm 0.004 \pm 0.006$              | <sup>5</sup> AAIJ       | 13AR   | LHCB | Repl. by AAIJ 15I      |
| $0.677 \pm 0.007 \pm 0.004$              | <sup>2</sup> AAD        | 12CV   | ATLS | Repl. by AAD 14U       |
| $0.657 \pm 0.009 \pm 0.008$              | <sup>2</sup> AAIJ       | 12D    | LHCB | Repl. by AAIJ 13AR     |
| $0.654 \pm 0.011 \pm 0.005$              | <sup>2,6</sup> AALTONEN | 12D    | CDF  | Repl. by AALTONEN 12AJ |
| $0.672 \pm 0.027 \pm 0.013$              | <sup>2</sup> ABAZOV     | 09E    | D0   | Repl. by ABAZOV 08AM   |
| $0.658 \pm 0.017 \pm 0.009$              | <sup>2,7</sup> AALTONEN | 08J    | CDF  | Repl. by AALTONEN 12D  |
| $0.658 \pm 0.022 \pm 0.004$              | <sup>2</sup> ABAZOV     | 08AMD0 |      | Repl. by ABAZOV 12D    |
| $0.658 \pm 0.035 \pm 0.0130$<br>$-0.004$ | <sup>2,7</sup> ABAZOV   | 07     | D0   | Repl. by ABAZOV 09E    |
| $0.714 \pm 0.007 \pm 0.010$<br>$-0.008$  | <sup>2,7</sup> ACOSTA   | 05     | CDF  | Repl. by AALTONEN 08J  |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  in the region  $m(KK) > 1.05$  GeV.

<sup>2</sup> Measured using a time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.


<sup>3</sup> Measured using a time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S) \phi$  decays.

<sup>4</sup> Measured using a time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>5</sup> Measured using a combined time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  and  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays.

<sup>6</sup> Assuming CPV phase  $\phi_s = -0.04$ .

<sup>7</sup> Assuming CPV phase  $\phi_s = 0$ .



$\Delta\Gamma_{B_s^0}$ 

"OUR EVALUATION" is an average performed by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) as described in our "Review on  $B$ - $\bar{B}$  Mixing" in the  $B^0$  section of these Listings. It includes the measurements of  $\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  listed in this section, as well as constraints from effective lifetimes with pure  $CP$  modes and flavor-specific modes.

| VALUE ( $10^{12} \text{ s}^{-1}$ )                                            | DOCUMENT ID              | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|--------------------------|-----------|------------------------|
| <b>0.085 <math>\pm</math> 0.004 OUR EVALUATION</b>                            |                          |           |                        |
| <b>0.081 <math>\pm</math> 0.006 OUR AVERAGE</b>                               |                          |           |                        |
| 0.077 $\pm$ 0.008 $\pm$ 0.003                                                 | <sup>1</sup> AAIJ        | 19Q LHCb  | $pp$ at 13 TeV         |
| 0.066 $\pm$ 0.018 $\pm$ 0.010                                                 | <sup>2</sup> AAIJ        | 17V LHCb  | $pp$ at 7, 8 TeV       |
| 0.085 $\pm$ 0.011 $\pm$ 0.007                                                 | <sup>3</sup> AAD         | 16AP ATLS | $pp$ at 7, 8 TeV       |
| 0.066 $^{+0.041}_{-0.044}$ $\pm$ 0.007                                        | <sup>4</sup> AAIJ        | 16AK LHCb | $pp$ at 7, 8 TeV       |
| 0.095 $\pm$ 0.013 $\pm$ 0.007                                                 | <sup>3</sup> KHACHATRYAN | 16S CMS   | $pp$ at 8 TeV          |
| 0.068 $\pm$ 0.026 $\pm$ 0.009                                                 | <sup>3</sup> AALTONEN    | 12AJ CDF  | $p\bar{p}$ at 1.96 TeV |
| 0.163 $^{+0.065}_{-0.064}$                                                    | <sup>3,5</sup> ABAZOV    | 12D D0    | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                          |           |                        |
| 0.0805 $\pm$ 0.0091 $\pm$ 0.0032                                              | <sup>1</sup> AAIJ        | 15I LHCb  | Repl. by AAIJ 19Q      |
| 0.053 $\pm$ 0.021 $\pm$ 0.010                                                 | <sup>3</sup> AAD         | 14U ATLS  | Repl. by AAD 16AP      |
| 0.106 $\pm$ 0.011 $\pm$ 0.007                                                 | <sup>6</sup> AAIJ        | 13AR LHCb | Repl. by AAIJ 15I      |
| 0.053 $\pm$ 0.021 $\pm$ 0.010                                                 | <sup>3</sup> AAD         | 12CV ATLS | Repl. by AAD 14U       |
| 0.123 $\pm$ 0.029 $\pm$ 0.011                                                 | <sup>3</sup> AAIJ        | 12D LHCb  | Repl. by AAIJ 13AR     |
| 0.075 $\pm$ 0.035 $\pm$ 0.006                                                 | <sup>7</sup> AALTONEN    | 12D CDF   | Repl. by AALTONEN 12AJ |
| 0.085 $^{+0.072}_{-0.078}$ $\pm$ 0.001                                        | <sup>8</sup> ABAZOV      | 09E D0    | Repl. by ABAZOV 08AM   |
| 0.076 $^{+0.059}_{-0.063}$ $\pm$ 0.006                                        | <sup>9</sup> AALTONEN    | 08J CDF   | Repl. by AALTONEN 12D  |
| 0.19 $\pm$ 0.07 $^{+0.02}_{-0.01}$                                            | <sup>3,10</sup> ABAZOV   | 08AMD0    | Repl. by ABAZOV 12D    |
| 0.12 $^{+0.08}_{-0.10}$ $\pm$ 0.02                                            | <sup>9,11</sup> ABAZOV   | 07 D0     | Repl. by ABAZOV 07N    |
| 0.13 $\pm$ 0.09                                                               | <sup>12</sup> ABAZOV     | 07N D0    | Repl. by ABAZOV 09E    |
| 0.47 $^{+0.19}_{-0.24}$ $\pm$ 0.01                                            | <sup>9</sup> ACOSTA      | 05 CDF    | Repl. by AALTONEN 08J  |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>2</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  in the region  $m(KK) > 1.05 \text{ GeV}$ .

<sup>3</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>4</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S)\phi$  decays.

<sup>5</sup> The error includes both statistical and systematic uncertainties.

<sup>6</sup> AAIJ 13AR result comes from a combined fit to  $B_s^0 \rightarrow J/\psi K^+ K^-$  and  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  data sets. Also reports  $\Delta\Gamma_s = 0.100 \pm 0.016 \pm 0.003 \text{ ps}^{-1}$  from a fit to  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>7</sup> Uses the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays and assuming  $CP$ -violating angle  $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$ .

- <sup>8</sup> Measured the angular and lifetime parameters for the time-dependent angular untagged decays  $B_d^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi \phi$ .  
<sup>9</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays and assuming  $CP$ -violating phase  $\phi_s = 0$ .  
<sup>10</sup> Obtains 90% CL interval  $-0.06 < \Delta\Gamma_s < 0.30$ .  
<sup>11</sup> ABAZOV 07 reports  $0.17 \pm 0.09 \pm 0.02$  with  $CP$ -violating phase  $\phi_s$  as a free parameter.  
<sup>12</sup> Combines  $D^0$  measurements of time-dependent angular distributions in  $B_s^0 \rightarrow J/\psi \phi$  and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

$$\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$$

$\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  are the decay rate average and difference between two  $B_s^0$   $CP$  eigenstates (light – heavy).

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>). It is derived from the averages of  $\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  (and their correlation).

| VALUE                                                                         | CL% | DOCUMENT ID            | TECN     | COMMENT                           |
|-------------------------------------------------------------------------------|-----|------------------------|----------|-----------------------------------|
| <b>0.129 ± 0.006 OUR EVALUATION</b>                                           |     |                        |          |                                   |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |                        |          |                                   |
| $0.090 \pm 0.009 \pm 0.023$                                                   |     | <sup>1</sup> ESEN      | 13 BELL  | $e^+e^- \rightarrow \Upsilon(5S)$ |
|                                                                               |     | <sup>2</sup> AAIJ      | 12D LHCB | $pp$ at 7 TeV                     |
|                                                                               |     | <sup>3</sup> AALTONEN  | 12D CDF  | $p\bar{p}$ at 1.96 TeV            |
|                                                                               |     | <sup>4</sup> ABAZOV    | 12D D0   | $p\bar{p}$ at 1.96 TeV            |
| $0.147^{+0.036+0.042}_{-0.030-0.041}$                                         |     | <sup>1</sup> ESEN      | 10 BELL  | $e^+e^- \rightarrow \Upsilon(5S)$ |
|                                                                               |     | <sup>5</sup> ABAZOV    | 09I D0   | $p\bar{p}$ at 1.96 TeV            |
| $0.072 \pm 0.021 \pm 0.022$                                                   | 95  | <sup>5</sup> AALTONEN  | 08F CDF  | $p\bar{p}$ at 1.96 TeV            |
| $>0.012$                                                                      |     | <sup>6</sup> AALTONEN  | 08J CDF  | Repl. by AALTONEN 12D             |
| $0.116^{+0.09}_{-0.10} \pm 0.010$                                             |     | <sup>6</sup> AALTONEN  | 08J CDF  | Repl. by AALTONEN 12D             |
| $0.079^{+0.038+0.031}_{-0.035-0.030}$                                         |     | <sup>5</sup> ABAZOV    | 07Y D0   | Repl. by ABAZOV 09I               |
| $0.24^{+0.28+0.03}_{-0.38-0.04}$                                              |     | <sup>6,7</sup> ABAZOV  | 05W D0   | Repl. by ABAZOV 08AM              |
| $0.65^{+0.25}_{-0.33} \pm 0.01$                                               |     | <sup>6</sup> ACOSTA    | 05 CDF   | Repl. by AALTONEN 08J             |
| $<0.46$                                                                       | 95  | <sup>8</sup> ABREU     | 00Y DLPH | $e^+e^- \rightarrow Z$            |
| $<0.69$                                                                       | 95  | <sup>9</sup> ABREU,P   | 00G DLPH | $e^+e^- \rightarrow Z$            |
| $0.25^{+0.21}_{-0.14}$                                                        |     | <sup>10</sup> BARATE   | 00K ALEP | $e^+e^- \rightarrow Z$            |
| $<0.83$                                                                       | 95  | <sup>11</sup> ABE      | 99D CDF  | $p\bar{p}$ at 1.8 TeV             |
| $<0.67$                                                                       | 95  | <sup>12</sup> ACCIARRI | 98S L3   | $e^+e^- \rightarrow Z$            |

<sup>1</sup> Assumes  $CP$  violation is negligible.

<sup>2</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>3</sup> Uses the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays and assuming  $CP$ -violating angle  $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$ .

<sup>4</sup> Measured using fully reconstructed  $B_s \rightarrow J/\psi \phi$  decays.

<sup>5</sup> Assumes  $2\text{B}(B_s^0 \rightarrow D_s^{(*)} D_s^{(*)}) \simeq \Delta\Gamma_s^{CP} / \Gamma_s$ .

<sup>6</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>7</sup> Uses  $|A_0|^2 - |A_{||}|^2 = 0.355 \pm 0.066$  from ACOSTA 05.

<sup>8</sup> Uses  $D_s^- \ell^+$ , and  $\phi \ell^+$  vertices.

<sup>9</sup> Measured using  $D_s$  hadron vertices.

<sup>10</sup> Uses  $\phi\phi$  correlations from  $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$ .

<sup>11</sup> ABE 99D assumes  $\tau_{B_s^0} = 1.55 \pm 0.05$  ps.

<sup>12</sup> ACCIARRI 98S assumes  $\tau_{B_s^0} = 1.49 \pm 0.06$  ps and PDG 98 values of  $b$  production fraction.

## $B_{sH}^0$ MEAN LIFE

$B_{sH}^0$  is the heavy mass state of two  $B_s^0$   $CP$  eigenstates.

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>). It is derived from the averages of  $\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  (and their correlation).

| VALUE ( $10^{-12}$ s)                                                         | DOCUMENT ID             | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-------------------------|-----------|------------------------|
| <b>1.620<math>\pm</math>0.007 OUR EVALUATION</b>                              |                         |           |                        |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                         |           |                        |
| 1.677 $\pm$ 0.034 $\pm$ 0.011                                                 | <sup>1</sup> SIRUNYAN   | 18BY CMS  | $pp$ at 8 TeV          |
| 2.04 $\pm$ 0.44 $\pm$ 0.05                                                    | <sup>2</sup> AAIJ       | 17AI LHCb | $pp$ at 7, 8, 13 TeV   |
| 1.70 $\pm$ 0.14 $\pm$ 0.05                                                    | <sup>3</sup> ABAZOV     | 16C D0    | $p\bar{p}$ at 1.96 TeV |
| 1.75 $\pm$ 0.12 $\pm$ 0.07                                                    | <sup>4</sup> AAIJ       | 13AB LHCb | $pp$ at 7 TeV          |
| 1.652 $\pm$ 0.024 $\pm$ 0.024                                                 | <sup>5</sup> AAIJ       | 13AR LHCb | $pp$ at 7 TeV          |
| 1.700 $\pm$ 0.040 $\pm$ 0.026                                                 | <sup>6</sup> AAIJ       | 12AN LHCb | $pp$ at 7 TeV          |
|                                                                               | <sup>7</sup> AALTONEN   | 12D CDF   | $p\bar{p}$ at 1.96 TeV |
| 1.70 $^{+0.12}_{-0.11}$ $\pm$ 0.03                                            | <sup>6</sup> AALTONEN   | 11AB CDF  | $p\bar{p}$ at 1.96 TeV |
| 1.613 $^{+0.123}_{-0.113}$                                                    | <sup>8,9</sup> AALTONEN | 08J CDF   | Repl. by AALTONEN 12D  |
| 1.58 $^{+0.39}_{-0.42}$ $^{+0.01}_{-0.02}$                                    | <sup>9</sup> ABAZOV     | 05W D0    | Repl. by ABAZOV 08AM   |
| 2.07 $^{+0.58}_{-0.46}$ $\pm$ 0.03                                            | <sup>9</sup> ACOSTA     | 05 CDF    | Repl. by AALTONEN 08J  |

<sup>1</sup> Measured using  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays with  $0.9240 < m(\pi\pi) < 1.0204$  GeV, which is dominated by the  $f_0(980)$  resonance, making it a  $CP$ -odd state.

<sup>2</sup> Measured using  $B_s \rightarrow \mu^+ \mu^-$  decays which, in the Standard Model, correspond to  $B_{sH}^0$  decays. Assumes  $-2 \operatorname{Re}(\lambda)/(1+|\lambda|^2) = 1$ .

<sup>3</sup> Measured using  $J/\psi \pi^+ \pi^-$  mode with  $0.880 < m(\pi\pi) < 1.080$  GeV/ $c^2$ , which is mostly  $J/\psi f(0)(980)$  mode, a pure  $CP$ -odd final state.

<sup>4</sup> Measured using a pure  $CP$ -odd final state  $J/\psi K_S^0$  with the assumption that contributions from penguin diagrams are small.

<sup>5</sup> Measured using  $B_s \rightarrow J/\psi \pi^+ \pi^-$  decays which, in the limit of  $\phi_s = 0$  and  $|\lambda| = 1$ , correspond to  $B_{sH}^0$  decays.

<sup>6</sup> Measured using a pure  $CP$ -odd final state  $J/\psi f_0(980)$ .

<sup>7</sup> Uses the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays assuming  $CP$ -violating angle  $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$ .

<sup>8</sup> Obtained from  $\Delta\Gamma_s$  and  $\Gamma_s$  fit with a correlation of 0.6.

<sup>9</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

## $B_{sL}^0$ MEAN LIFE

$B_{sL}^0$  is the light mass state of two  $B_s^0$  CP eigenstates.

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>). It is derived from the averages of  $\Gamma_{B_s^0}$  and  $\Delta\Gamma_{B_s^0}$  (and their correlation).

| VALUE ( $10^{-12}$ s)                                                         | DOCUMENT ID             | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-------------------------|-----------|------------------------|
| <b>1.423 ± 0.005 OUR EVALUATION</b>                                           |                         |           |                        |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                         |           |                        |
| 1.40 ± 0.02                                                                   | <sup>1</sup> SIRUNYAN   | 18BY CMS  | $pp$ at 8 TeV          |
| 1.479 ± 0.034 ± 0.011                                                         | <sup>2</sup> AAIJ       | 16AL LHCb | $pp$ at 7, 8 TeV       |
| 1.379 ± 0.026 ± 0.017                                                         | <sup>3</sup> AAIJ       | 14F LHCb  | $pp$ at 7, 8 TeV       |
| 1.407 ± 0.016 ± 0.007                                                         | <sup>4</sup> AAIJ       | 14R LHCb  | $pp$ at 7 TeV          |
| 1.440 ± 0.096 ± 0.009                                                         | <sup>4</sup> AAIJ       | 12 LHCb   | Repl. by AAIJ 14R      |
| 1.455 ± 0.046 ± 0.006                                                         | <sup>4</sup> AAIJ       | 12R LHCb  | Repl. by AAIJ 14R      |
|                                                                               | <sup>5</sup> AALTONEN   | 12D CDF   | $p\bar{p}$ at 1.96 TeV |
| 1.437 <sup>+0.054</sup> <sub>-0.047</sub>                                     | <sup>6,7</sup> AALTONEN | 08J CDF   | Repl. by AALTONEN 12D  |
| 1.24 <sup>+0.14</sup> <sub>-0.11</sub> <sup>+0.01</sup> <sub>-0.02</sub>      | <sup>7</sup> ABAZOV     | 05W D0    | Repl. by ABAZOV 08AM   |
| 1.05 <sup>+0.16</sup> <sub>-0.13</sub> ± 0.02                                 | <sup>7</sup> ACOSTA     | 05 CDF    | Repl. by AALTONEN 08J  |
| 1.27 ± 0.33 ± 0.08                                                            | <sup>8</sup> BARATE     | 00K ALEP  | $e^+e^- \rightarrow Z$ |

<sup>1</sup> Measured using results in SIRUNYAN 18BY for the heavy  $B_s^0$  lifetime obtained from  $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$  decays and the average effective  $B_s^0 \rightarrow J/\psi \phi$  lifetime, and magnitude squared of the CP-odd amplitude  $|A_{\perp}|^2 = 0.250 \pm 0.006$ . The uncertainty includes all statistical and systematic contributions.

<sup>2</sup> Measured using  $B_s^0 \rightarrow J/\psi \eta$  decays.

<sup>3</sup> Measured using  $B_s^0 \rightarrow D_s^- D_s^+$ . The effective lifetime is translated into a decay width of  $\Gamma_L = 0.725 \pm 0.014 \pm 0.009$  ps<sup>-1</sup>.

<sup>4</sup> Measured using  $B_s^0 \rightarrow K^+ K^-$  decays. There may still be CPV in the decay.

<sup>5</sup> Uses the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays and assuming CP-violating angle  $\beta_s(B^0 \rightarrow J/\psi \phi) = 0.02$ .

<sup>6</sup> Obtained from  $\Delta\Gamma_s$  and  $\Gamma_s$  fit with a correlation of 0.6.

<sup>7</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>8</sup> Uses  $\phi\phi$  correlations from  $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$ .

## $B_s^0$ MEAN LIFE (Flavor specific)

| VALUE ( $10^{-12}$ s)                                                                        | DOCUMENT ID         | TECN      | COMMENT                |
|----------------------------------------------------------------------------------------------|---------------------|-----------|------------------------|
| <b>1.527 ± 0.011 OUR EVALUATION</b>                                                          |                     |           |                        |
| <b>1.526 ± 0.015 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below. |                     |           |                        |
| 1.547 ± 0.013 ± 0.011                                                                        | <sup>1</sup> AAIJ   | 17AN LHCb | $pp$ at 7, 8 TeV       |
| 1.479 ± 0.010 ± 0.021                                                                        | <sup>2</sup> ABAZOV | 15A D0    | $p\bar{p}$ at 1.96 TeV |
| 1.535 ± 0.015 ± 0.014                                                                        | <sup>3</sup> AAIJ   | 14AX LHCb | $pp$ at 7 TeV          |



|                                 |                         |      |      |                        |
|---------------------------------|-------------------------|------|------|------------------------|
| $1.52 \pm 0.15 \pm 0.01$        | <sup>4</sup> AAIJ       | 14F  | LHCB | $pp$ at 7, 8 TeV       |
| $1.60 \pm 0.06 \pm 0.01$        | <sup>5</sup> AAIJ       | 14R  | LHCB | $pp$ at 7 TeV          |
| $1.518 \pm 0.041 \pm 0.027$     | <sup>6</sup> AALTONEN   | 11AP | CDF  | $p\bar{p}$ at 1.96 TeV |
| $1.42^{+0.14}_{-0.13} \pm 0.03$ | <sup>7</sup> ABREU      | 00Y  | DLPH | $e^+e^- \rightarrow Z$ |
| $1.36 \pm 0.09^{+0.06}_{-0.05}$ | <sup>8</sup> ABE        | 99D  | CDF  | $p\bar{p}$ at 1.8 TeV  |
| $1.50^{+0.16}_{-0.15} \pm 0.04$ | <sup>8</sup> ACKERSTAFF | 98G  | OPAL | $e^+e^- \rightarrow Z$ |
| $1.54^{+0.14}_{-0.13} \pm 0.04$ | <sup>8</sup> BUSKULIC   | 96M  | ALEP | $e^+e^- \rightarrow Z$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                     |                     |     |    |                     |
|-------------------------------------|---------------------|-----|----|---------------------|
| $1.398 \pm 0.044^{+0.028}_{-0.025}$ | <sup>9</sup> ABAZOV | 06V | D0 | Repl. by ABAZOV 15A |
|-------------------------------------|---------------------|-----|----|---------------------|

<sup>1</sup> AAIJ 17AN value was measured using  $B_S^0 \rightarrow D_S^{(*)-} \mu^+ \nu_\mu$  decays relative to  $B^0 \rightarrow D^{(*)-} \mu^+ \nu_\mu$  decays.

<sup>2</sup> Measured using  $B_S^0 \rightarrow D_S^- \mu^+ \nu_\mu X$  decays.

<sup>3</sup> Measured using the  $B_S^0 \rightarrow D_S^- \pi^+$  decays.

<sup>4</sup> Measured using  $B_S^0 \rightarrow D^+ D_S^-$ .


<sup>5</sup> Measured using  $B_S^0 \rightarrow \pi^+ K^-$  decays.

<sup>6</sup> AALTONEN 11AP combines the fully reconstructed  $B_S^0 \rightarrow D_S^- \pi^+$  decays and partially reconstructed  $B_S^0 \rightarrow D_S X$  decays.

<sup>7</sup> Uses  $D_S^- \ell^+$ , and  $\phi \ell^+$  vertices.

<sup>8</sup> Measured using  $D_S^- \ell^+$  vertices.

<sup>9</sup> Measured using  $D_S^- \mu^+$  vertices.



### $B_s^0$ MEAN LIFE ( $B_s \rightarrow J/\psi\phi$ )

| VALUE ( $10^{-12}$ s)                                                         | DOCUMENT ID           | TECN     | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|----------|------------------------|
| <b><math>1.480 \pm 0.007</math> OUR EVALUATION</b>                            |                       |          |                        |
| <b><math>1.480 \pm 0.007</math> OUR AVERAGE</b>                               |                       |          |                        |
| $1.481 \pm 0.007 \pm 0.005$                                                   | <sup>1</sup> SIRUNYAN | 18BY CMS | $pp$ at 8 TeV          |
| $1.480 \pm 0.011 \pm 0.005$                                                   | <sup>1</sup> AAIJ     | 14E LHCb | $pp$ at 7 TeV          |
| $1.444^{+0.098}_{-0.090} \pm 0.020$                                           | <sup>1</sup> ABAZOV   | 05B D0   | $p\bar{p}$ at 1.96 TeV |
| $1.34^{+0.23}_{-0.19} \pm 0.05$                                               | <sup>2</sup> ABE      | 98B CDF  | $p\bar{p}$ at 1.8 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |          |                        |
| $1.39^{+0.13}_{-0.16}^{+0.01}_{-0.02}$                                        | <sup>2</sup> ABAZOV   | 05W D0   | $p\bar{p}$ at 1.96 TeV |
| $1.34^{+0.23}_{-0.19} \pm 0.05$                                               | <sup>3</sup> ABE      | 96N CDF  | Repl. by ABE 98B       |

<sup>1</sup> Measured using fully reconstructed  $B_s \rightarrow J/\psi\phi$  decays.

<sup>2</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays.

<sup>3</sup> ABE 96N uses  $58 \pm 12$  exclusive  $B_s \rightarrow J/\psi\phi$  events.

### $\tau_{B_s^0}/\tau_{B^0}$ MEAN LIFE RATIO

$\tau_{B_s^0}/\tau_{B^0}$  (direct measurements)

| VALUE                                                                                                            | DOCUMENT ID           | TECN     | COMMENT       |
|------------------------------------------------------------------------------------------------------------------|-----------------------|----------|---------------|
| <b><math>0.980 \pm 0.006 \pm 0.003</math></b>                                                                    | <sup>1</sup> SIRUNYAN | 18BY CMS | $pp$ at 8 TeV |
| <sup>1</sup> Measured using $B_s^0 \rightarrow J/\psi\phi(1020)$ and $B^0 \rightarrow J/\psi K^*(892)^0$ decays. |                       |          |               |

$\Gamma_{B_s^0} - \Gamma_{B^0}$ 

| VALUE ( $10^{12} \text{ s}^{-1}$ )                                                                        | DOCUMENT ID       | TECN     | COMMENT        |
|-----------------------------------------------------------------------------------------------------------|-------------------|----------|----------------|
| $-0.0041 \pm 0.0024 \pm 0.0015$                                                                           | <sup>1</sup> AAIJ | 19Q LHCb | $pp$ at 13 TeV |
| <sup>1</sup> Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays. |                   |          |                |

 $\Gamma_{B_{sH}^0} - \Gamma_{B^0}$ 

| VALUE ( $10^{12} \text{ s}^{-1}$ )                                      | DOCUMENT ID       | TECN      | COMMENT              |
|-------------------------------------------------------------------------|-------------------|-----------|----------------------|
| $-0.05 \pm 0.004 \pm 0.004$                                             | <sup>1</sup> AAIJ | 19AF LHCb | $pp$ at 7, 8, 13 TeV |
| <sup>1</sup> Measured in $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays. |                   |           |                      |

 $B_s^0$  DECAY MODES

These branching fractions all scale with  $B(\bar{b} \rightarrow B_s^0)$ .

The branching fraction  $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything})$  is not a pure measurement since the measured product branching fraction  $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything})$  was used to determine  $B(\bar{b} \rightarrow B_s^0)$ , as described in the note on “ $B^0$ - $\bar{B}^0$  Mixing”

For inclusive branching fractions, e.g.,  $B \rightarrow D^\pm \text{ anything}$ , the values usually are multiplicities, not branching fractions. They can be greater than one.

| Mode                                                                               | Fraction ( $\Gamma_i/\Gamma$ )       | Scale factor/<br>Confidence level |
|------------------------------------------------------------------------------------|--------------------------------------|-----------------------------------|
| $\Gamma_1 \quad D_s^- \text{ anything}$                                            | (93 $\pm$ 25 ) %                     |                                   |
| $\Gamma_2 \quad \ell \nu_\ell X$                                                   | ( 9.6 $\pm$ 0.8 ) %                  |                                   |
| $\Gamma_3 \quad e^+ \nu X^-$                                                       | ( 9.1 $\pm$ 0.8 ) %                  |                                   |
| $\Gamma_4 \quad \mu^+ \nu X^-$                                                     | (10.2 $\pm$ 1.0 ) %                  |                                   |
| $\Gamma_5 \quad D_s^- \ell^+ \nu_\ell \text{ anything}$                            | [a] ( 8.1 $\pm$ 1.3 ) %              |                                   |
| $\Gamma_6 \quad D_s^{*-} \ell^+ \nu_\ell \text{ anything}$                         | ( 5.4 $\pm$ 1.1 ) %                  |                                   |
| $\Gamma_7 \quad D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D_s^{*-} K_S^0$ | ( 2.7 $\pm$ 0.7 ) $\times 10^{-3}$   |                                   |
| $\Gamma_8 \quad D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+$    | ( 4.4 $\pm$ 1.3 ) $\times 10^{-3}$   |                                   |
| $\Gamma_9 \quad D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+$    | ( 2.7 $\pm$ 1.0 ) $\times 10^{-3}$   |                                   |
| $\Gamma_{10} \quad D_s^- \pi^+$                                                    | ( 3.00 $\pm$ 0.23 ) $\times 10^{-3}$ |                                   |
| $\Gamma_{11} \quad D_s^- \rho^+$                                                   | ( 6.9 $\pm$ 1.4 ) $\times 10^{-3}$   |                                   |
| $\Gamma_{12} \quad D_s^- \pi^+ \pi^+ \pi^-$                                        | ( 6.1 $\pm$ 1.0 ) $\times 10^{-3}$   |                                   |
| $\Gamma_{13} \quad D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-$   | ( 2.5 $\pm$ 0.8 ) $\times 10^{-5}$   |                                   |
| $\Gamma_{14} \quad D_s^\mp K^\pm$                                                  | ( 2.27 $\pm$ 0.19 ) $\times 10^{-4}$ |                                   |

|               |                                                              |                                         |        |
|---------------|--------------------------------------------------------------|-----------------------------------------|--------|
| $\Gamma_{15}$ | $D_s^- K^+ \pi^+ \pi^-$                                      | $(3.2 \pm 0.6) \times 10^{-4}$          |        |
| $\Gamma_{16}$ | $D_s^+ D_s^-$                                                | $(4.4 \pm 0.5) \times 10^{-3}$          |        |
| $\Gamma_{17}$ | $D_s^- D^+$                                                  | $(2.8 \pm 0.5) \times 10^{-4}$          |        |
| $\Gamma_{18}$ | $D^+ D^-$                                                    | $(2.2 \pm 0.6) \times 10^{-4}$          |        |
| $\Gamma_{19}$ | $D^0 \bar{D}^0$                                              | $(1.9 \pm 0.5) \times 10^{-4}$          |        |
| $\Gamma_{20}$ | $D_s^{*-} \pi^+$                                             | $(2.0 \pm 0.5) \times 10^{-3}$          |        |
| $\Gamma_{21}$ | $D_s^{*\mp} K^\pm$                                           | $(1.33 \pm 0.35) \times 10^{-4}$        |        |
| $\Gamma_{22}$ | $D_s^{*-} \rho^+$                                            | $(9.6 \pm 2.1) \times 10^{-3}$          |        |
| $\Gamma_{23}$ | $D_s^{*+} D_s^- + D_s^{*-} D_s^+$                            | $(1.39 \pm 0.17) \%$                    |        |
| $\Gamma_{24}$ | $D_s^{*+} D_s^{*-}$                                          | $(1.44 \pm 0.21) \%$                    | S=1.1  |
| $\Gamma_{25}$ | $D_s^{(*)+} D_s^{(*)-}$                                      | $(4.5 \pm 1.4) \%$                      |        |
| $\Gamma_{26}$ | $\bar{D}^{*0} \bar{K}^0$                                     | $(2.8 \pm 1.1) \times 10^{-4}$          |        |
| $\Gamma_{27}$ | $\bar{D}^0 \bar{K}^0$                                        | $(4.3 \pm 0.9) \times 10^{-4}$          |        |
| $\Gamma_{28}$ | $\bar{D}^0 K^- \pi^+$                                        | $(1.04 \pm 0.13) \times 10^{-3}$        |        |
| $\Gamma_{29}$ | $\bar{D}^0 \bar{K}^*(892)^0$                                 | $(4.4 \pm 0.6) \times 10^{-4}$          |        |
| $\Gamma_{30}$ | $\bar{D}^0 \bar{K}^*(1410)$                                  | $(3.9 \pm 3.5) \times 10^{-4}$          |        |
| $\Gamma_{31}$ | $\bar{D}^0 \bar{K}_0^*(1430)$                                | $(3.0 \pm 0.7) \times 10^{-4}$          |        |
| $\Gamma_{32}$ | $\bar{D}^0 \bar{K}_2^*(1430)$                                | $(1.1 \pm 0.4) \times 10^{-4}$          |        |
| $\Gamma_{33}$ | $\bar{D}^0 \bar{K}^*(1680)$                                  | $< 7.8 \times 10^{-5}$                  | CL=90% |
| $\Gamma_{34}$ | $\bar{D}^0 \bar{K}_0^*(1950)$                                | $< 1.1 \times 10^{-4}$                  | CL=90% |
| $\Gamma_{35}$ | $\bar{D}^0 \bar{K}_3^*(1780)$                                | $< 2.6 \times 10^{-5}$                  | CL=90% |
| $\Gamma_{36}$ | $\bar{D}^0 \bar{K}_4^*(2045)$                                | $< 3.1 \times 10^{-5}$                  | CL=90% |
| $\Gamma_{37}$ | $\bar{D}^0 K^- \pi^+ \text{ (non-resonant)}$                 | $(2.1 \pm 0.8) \times 10^{-4}$          |        |
| $\Gamma_{38}$ | $D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \bar{D}^0 K^-$ | $(2.6 \pm 0.4) \times 10^{-4}$          |        |
| $\Gamma_{39}$ | $D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-$ | $(1.6 \pm 0.8) \times 10^{-5}$          |        |
| $\Gamma_{40}$ | $D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-$ | $(5 \pm 4) \times 10^{-5}$              |        |
| $\Gamma_{41}$ | $D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow \bar{D}^0 K^-$ | $(2.2 \pm 0.6) \times 10^{-5}$          |        |
| $\Gamma_{42}$ | $\bar{D}^0 K^+ K^-$                                          | $(5.5 \pm 0.8) \times 10^{-5}$          |        |
| $\Gamma_{43}$ | $\bar{D}^0 f_0(980)$                                         | $< 3.1 \times 10^{-6}$                  | CL=90% |
| $\Gamma_{44}$ | $\bar{D}^0 \phi$                                             | $(3.0 \pm 0.5) \times 10^{-5}$          |        |
| $\Gamma_{45}$ | $\bar{D}^{*0} \phi$                                          | $(3.7 \pm 0.6) \times 10^{-5}$          |        |
| $\Gamma_{46}$ | $D^{*\mp} \pi^\pm$                                           | $< 6.1 \times 10^{-6}$                  | CL=90% |
| $\Gamma_{47}$ | $\eta_c \phi$                                                | $(5.0 \pm 0.9) \times 10^{-4}$          |        |
| $\Gamma_{48}$ | $\eta_c \pi^+ \pi^-$                                         | $(1.8 \pm 0.7) \times 10^{-4}$          |        |
| $\Gamma_{49}$ | $J/\psi(1S) \phi$                                            | $(1.08 \pm 0.08) \times 10^{-3}$        |        |
| $\Gamma_{50}$ | $J/\psi(1S) \phi \phi$                                       | $(1.24^{+0.17}_{-0.19}) \times 10^{-5}$ |        |
| $\Gamma_{51}$ | $J/\psi(1S) \pi^0$                                           | $< 1.2 \times 10^{-3}$                  | CL=90% |
| $\Gamma_{52}$ | $J/\psi(1S) \eta$                                            | $(4.0 \pm 0.7) \times 10^{-4}$          | S=1.4  |

|               |                                                             |                                            |        |
|---------------|-------------------------------------------------------------|--------------------------------------------|--------|
| $\Gamma_{53}$ | $J/\psi(1S)K_S^0$                                           | $(1.88 \pm 0.15) \times 10^{-5}$           |        |
| $\Gamma_{54}$ | $J/\psi(1S)\bar{K}^{*0}(892)$                               | $(4.1 \pm 0.4) \times 10^{-5}$             |        |
| $\Gamma_{55}$ | $J/\psi(1S)\eta'$                                           | $(3.3 \pm 0.4) \times 10^{-4}$             |        |
| $\Gamma_{56}$ | $J/\psi(1S)\pi^+\pi^-$                                      | $(2.09 \pm 0.23) \times 10^{-4}$           | S=1.3  |
| $\Gamma_{57}$ | $J/\psi(1S)f_0(500), f_0 \rightarrow \pi^+\pi^-$            | $< 4 \times 10^{-6}$                       | CL=90% |
| $\Gamma_{58}$ | $J/\psi(1S)\rho, \rho \rightarrow \pi^+\pi^-$               | $< 4 \times 10^{-6}$                       | CL=90% |
| $\Gamma_{59}$ | $J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-$            | $(1.28 \pm 0.18) \times 10^{-4}$           | S=1.7  |
| $\Gamma_{60}$ | $J/\psi(1S)f_2(1270), f_2 \rightarrow \pi^+\pi^-$           | $(1.1 \pm 0.4) \times 10^{-6}$             |        |
| $\Gamma_{61}$ | $J/\psi(1S)f_2(1270)_0, f_2 \rightarrow \pi^+\pi^-$         | $(7.5 \pm 1.8) \times 10^{-7}$             |        |
| $\Gamma_{62}$ | $J/\psi(1S)f_2(1270)_{  }, f_2 \rightarrow \pi^+\pi^-$      | $(1.09 \pm 0.34) \times 10^{-6}$           |        |
| $\Gamma_{63}$ | $J/\psi(1S)f_2(1270)_{\perp}, f_2 \rightarrow \pi^+\pi^-$   | $(1.3 \pm 0.8) \times 10^{-6}$             |        |
| $\Gamma_{64}$ | $J/\psi(1S)f_0(1370), f_0 \rightarrow \pi^+\pi^-$           | $(4.5 \pm_{-4.0}^{0.7}) \times 10^{-5}$    |        |
| $\Gamma_{65}$ | $J/\psi(1S)f_0(1500), f_0 \rightarrow \pi^+\pi^-$           | $(2.11 \pm_{-0.29}^{0.40}) \times 10^{-5}$ |        |
| $\Gamma_{66}$ | $J/\psi(1S)f_2'(1525)_0, f_2' \rightarrow \pi^+\pi^-$       | $(1.07 \pm 0.24) \times 10^{-6}$           |        |
| $\Gamma_{67}$ | $J/\psi(1S)f_2'(1525)_{  }, f_2' \rightarrow \pi^+\pi^-$    | $(1.3 \pm_{-0.9}^{2.7}) \times 10^{-7}$    |        |
| $\Gamma_{68}$ | $J/\psi(1S)f_2'(1525)_{\perp}, f_2' \rightarrow \pi^+\pi^-$ | $(5 \pm 4) \times 10^{-7}$                 |        |
| $\Gamma_{69}$ | $J/\psi(1S)f_0(1790), f_0 \rightarrow \pi^+\pi^-$           | $(5.0 \pm_{-1.1}^{11.0}) \times 10^{-6}$   |        |
| $\Gamma_{70}$ | $J/\psi(1S)\pi^+\pi^-$ (nonresonant)                        | $(1.8 \pm_{-0.4}^{1.1}) \times 10^{-5}$    |        |
| $\Gamma_{71}$ | $J/\psi(1S)\bar{K}^0\pi^+\pi^-$                             | $< 4.4 \times 10^{-5}$                     | CL=90% |
| $\Gamma_{72}$ | $J/\psi(1S)K^+K^-$                                          | $(7.9 \pm 0.7) \times 10^{-4}$             |        |
| $\Gamma_{73}$ | $J/\psi(1S)K^0K^-\pi^+ + \text{c.c.}$                       | $(9.2 \pm 1.3) \times 10^{-4}$             |        |
| $\Gamma_{74}$ | $J/\psi(1S)\bar{K}^0K^+K^-$                                 | $< 1.2 \times 10^{-5}$                     | CL=90% |
| $\Gamma_{75}$ | $J/\psi(1S)f_2'(1525)$                                      | $(2.6 \pm 0.6) \times 10^{-4}$             |        |
| $\Gamma_{76}$ | $J/\psi(1S)p\bar{p}$                                        | $(3.6 \pm 0.4) \times 10^{-6}$             |        |
| $\Gamma_{77}$ | $J/\psi(1S)\gamma$                                          | $< 7.3 \times 10^{-6}$                     | CL=90% |
| $\Gamma_{78}$ | $J/\psi(1S)\pi^+\pi^-\pi^+\pi^-$                            | $(7.8 \pm 1.0) \times 10^{-5}$             |        |
| $\Gamma_{79}$ | $J/\psi(1S)f_1(1285)$                                       | $(7.2 \pm 1.4) \times 10^{-5}$             |        |
| $\Gamma_{80}$ | $\psi(2S)\eta$                                              | $(3.3 \pm 0.9) \times 10^{-4}$             |        |
| $\Gamma_{81}$ | $\psi(2S)\eta'$                                             | $(1.29 \pm 0.35) \times 10^{-4}$           |        |
| $\Gamma_{82}$ | $\psi(2S)\pi^+\pi^-$                                        | $(7.1 \pm 1.3) \times 10^{-5}$             |        |
| $\Gamma_{83}$ | $\psi(2S)\phi$                                              | $(5.4 \pm 0.6) \times 10^{-4}$             |        |
| $\Gamma_{84}$ | $\psi(2S)K^-\pi^+$                                          | $(3.1 \pm 0.4) \times 10^{-5}$             |        |

|                |                                                    |                                  |        |
|----------------|----------------------------------------------------|----------------------------------|--------|
| $\Gamma_{85}$  | $\psi(2S)\bar{K}^*(892)^0$                         | $(3.3 \pm 0.5) \times 10^{-5}$   |        |
| $\Gamma_{86}$  | $\chi_{c1}\phi$                                    | $(2.04 \pm 0.30) \times 10^{-4}$ |        |
| $\Gamma_{87}$  | $\chi_{c1}K^+K^-$                                  |                                  |        |
| $\Gamma_{88}$  | $\chi_{c2}K^+K^-$                                  |                                  |        |
| $\Gamma_{89}$  | $\pi^+\pi^-$                                       | $(7.0 \pm 1.0) \times 10^{-7}$   |        |
| $\Gamma_{90}$  | $\pi^0\pi^0$                                       | $< 2.1 \times 10^{-4}$           | CL=90% |
| $\Gamma_{91}$  | $\eta\pi^0$                                        | $< 1.0 \times 10^{-3}$           | CL=90% |
| $\Gamma_{92}$  | $\eta\eta$                                         | $< 1.5 \times 10^{-3}$           | CL=90% |
| $\Gamma_{93}$  | $\rho^0\rho^0$                                     | $< 3.20 \times 10^{-4}$          | CL=90% |
| $\Gamma_{94}$  | $\eta'\eta'$                                       | $(3.3 \pm 0.7) \times 10^{-5}$   |        |
| $\Gamma_{95}$  | $\eta'\phi$                                        | $< 8.2 \times 10^{-7}$           | CL=90% |
| $\Gamma_{96}$  | $\phi f_0(980), f_0(980) \rightarrow \pi^+\pi^-$   | $(1.12 \pm 0.21) \times 10^{-6}$ |        |
| $\Gamma_{97}$  | $\phi f_2(1270), f_2(1270) \rightarrow \pi^+\pi^-$ | $(6.1 \pm 1.8) \times 10^{-7}$   |        |
| $\Gamma_{98}$  | $\phi\rho^0$                                       | $(2.7 \pm 0.8) \times 10^{-7}$   |        |
| $\Gamma_{99}$  | $\phi\pi^+\pi^-$                                   | $(3.5 \pm 0.5) \times 10^{-6}$   |        |
| $\Gamma_{100}$ | $\phi\phi$                                         | $(1.87 \pm 0.15) \times 10^{-5}$ |        |
| $\Gamma_{101}$ | $\phi\phi\phi$                                     | $(2.2 \pm 0.7) \times 10^{-6}$   |        |
| $\Gamma_{102}$ | $\pi^+K^-$                                         | $(5.8 \pm 0.7) \times 10^{-6}$   |        |
| $\Gamma_{103}$ | $K^+K^-$                                           | $(2.66 \pm 0.22) \times 10^{-5}$ |        |
| $\Gamma_{104}$ | $K^0\bar{K}^0$                                     | $(2.0 \pm 0.6) \times 10^{-5}$   |        |
| $\Gamma_{105}$ | $K^0\pi^+\pi^-$                                    | $(9.5 \pm 2.1) \times 10^{-6}$   |        |
| $\Gamma_{106}$ | $K^0K^\pm\pi^\mp$                                  | $(8.4 \pm 0.9) \times 10^{-5}$   |        |
| $\Gamma_{107}$ | $K^*(892)^-\pi^+$                                  | $(2.9 \pm 1.1) \times 10^{-6}$   |        |
| $\Gamma_{108}$ | $K^*(892)^\pm K^\mp$                               | $(1.9 \pm 0.5) \times 10^{-5}$   |        |
| $\Gamma_{109}$ | $K_0^*(1430)^\pm K^\mp$                            | $(3.1 \pm 2.5) \times 10^{-5}$   |        |
| $\Gamma_{110}$ | $K_2^*(1430)^\pm K^\mp$                            | $(1.0 \pm 1.7) \times 10^{-5}$   |        |
| $\Gamma_{111}$ | $K^*(892)^0\bar{K}^0 + \text{c.c.}$                | $(2.0 \pm 0.6) \times 10^{-5}$   |        |
| $\Gamma_{112}$ | $K_0^*(1430)\bar{K}^0 + \text{c.c.}$               | $(3.3 \pm 1.0) \times 10^{-5}$   |        |
| $\Gamma_{113}$ | $K_2^*(1430)^0\bar{K}^0 + \text{c.c.}$             | $(1.7 \pm 2.2) \times 10^{-5}$   |        |
| $\Gamma_{114}$ | $K_S^0\bar{K}^*(892)^0 + \text{c.c.}$              | $(1.6 \pm 0.4) \times 10^{-5}$   |        |
| $\Gamma_{115}$ | $K_S^0K^+K^-$                                      | $(1.3 \pm 0.6) \times 10^{-6}$   |        |
| $\Gamma_{116}$ | $\bar{K}^*(892)^0\rho^0$                           | $< 7.67 \times 10^{-4}$          | CL=90% |
| $\Gamma_{117}$ | $\bar{K}^*(892)^0K^*(892)^0$                       | $(1.11 \pm 0.27) \times 10^{-5}$ |        |
| $\Gamma_{118}$ | $K^*(892)^0\bar{K}_2^*(1430)^0$                    |                                  |        |
| $\Gamma_{119}$ | $K_2^*(1430)^0\bar{K}^*(892)^0$                    |                                  |        |
| $\Gamma_{120}$ | $K_2^*(1430)^0\bar{K}_2^*(1430)^0$                 |                                  |        |
| $\Gamma_{121}$ | $\phi K^*(892)^0$                                  | $(1.14 \pm 0.30) \times 10^{-6}$ |        |
| $\Gamma_{122}$ | $p\bar{p}$                                         | $< 1.5 \times 10^{-8}$           | CL=90% |
| $\Gamma_{123}$ | $p\bar{p}K^+K^-$                                   | $(4.5 \pm 0.5) \times 10^{-6}$   |        |
| $\Gamma_{124}$ | $p\bar{p}K^+\pi^-$                                 | $(1.39 \pm 0.26) \times 10^{-6}$ |        |
| $\Gamma_{125}$ | $p\bar{p}\pi^+\pi^-$                               | $(4.3 \pm 2.0) \times 10^{-7}$   |        |

|                |                                   |                                |        |
|----------------|-----------------------------------|--------------------------------|--------|
| $\Gamma_{126}$ | $p\bar{\Lambda}K^- + \text{c.c.}$ | $(5.5 \pm 1.0) \times 10^{-6}$ |        |
| $\Gamma_{127}$ | $\Lambda_c^- \Lambda \pi^+$       | $(3.6 \pm 1.6) \times 10^{-4}$ |        |
| $\Gamma_{128}$ | $\Lambda_c^- \Lambda_c^+$         | $< 8.0 \times 10^{-5}$         | CL=95% |

**Lepton Family number (LF) violating modes or  
 $\Delta B = 1$  weak neutral current (B1) modes**

|                |                                                                 |    |                                |        |
|----------------|-----------------------------------------------------------------|----|--------------------------------|--------|
| $\Gamma_{129}$ | $\gamma\gamma$                                                  | B1 | $< 3.1 \times 10^{-6}$         | CL=90% |
| $\Gamma_{130}$ | $\phi\gamma$                                                    | B1 | $(3.4 \pm 0.4) \times 10^{-5}$ |        |
| $\Gamma_{131}$ | $\mu^+ \mu^-$                                                   | B1 | $(3.0 \pm 0.4) \times 10^{-9}$ |        |
| $\Gamma_{132}$ | $e^+ e^-$                                                       | B1 | $< 2.8 \times 10^{-7}$         | CL=90% |
| $\Gamma_{133}$ | $\tau^+ \tau^-$                                                 | B1 | $< 6.8 \times 10^{-3}$         | CL=95% |
| $\Gamma_{134}$ | $\mu^+ \mu^- \mu^+ \mu^-$                                       | B1 | $< 2.5 \times 10^{-9}$         | CL=95% |
| $\Gamma_{135}$ | $SP, S \rightarrow \mu^+ \mu^-,$<br>$P \rightarrow \mu^+ \mu^-$ | B1 | [b] $< 2.2 \times 10^{-9}$     | CL=95% |
| $\Gamma_{136}$ | $\phi(1020)\mu^+ \mu^-$                                         | B1 | $(8.2 \pm 1.2) \times 10^{-7}$ |        |
| $\Gamma_{137}$ | $K^*(892)^0 \mu^+ \mu^-$                                        |    | $(2.9 \pm 1.1) \times 10^{-8}$ |        |
| $\Gamma_{138}$ | $\pi^+ \pi^- \mu^+ \mu^-$                                       | B1 | $(8.4 \pm 1.7) \times 10^{-8}$ |        |
| $\Gamma_{139}$ | $\phi \nu \bar{\nu}$                                            | B1 | $< 5.4 \times 10^{-3}$         | CL=90% |
| $\Gamma_{140}$ | $e^\pm \mu^\mp$                                                 | LF | [c] $< 5.4 \times 10^{-9}$     | CL=90% |
| $\Gamma_{141}$ | $\mu^\pm \tau^\mp$                                              |    | $< 4.2 \times 10^{-5}$         | CL=95% |

[a] Not a pure measurement. See note at head of  $B_s^0$  Decay Modes.

[b] Here  $S$  and  $P$  are the hypothetical scalar and pseudoscalar particles with masses of  $2.5 \text{ GeV}/c^2$  and  $214.3 \text{ MeV}/c^2$ , respectively.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

### CONSTRAINED FIT INFORMATION

An overall fit to 12 branching ratios uses 20 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 26.7$  for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

|           |          |          |          |          |          |          |
|-----------|----------|----------|----------|----------|----------|----------|
| $x_{12}$  | 28       |          |          |          |          |          |
| $x_{14}$  | 92       | 26       |          |          |          |          |
| $x_{49}$  | 0        | 0        | 0        |          |          |          |
| $x_{56}$  | 0        | 0        | 0        | 72       |          |          |
| $x_{59}$  | 0        | 0        | 0        | 57       | 67       |          |
| $x_{100}$ | 0        | 0        | 0        | 29       | 21       | 17       |
|           | $x_{10}$ | $x_{12}$ | $x_{14}$ | $x_{49}$ | $x_{56}$ | $x_{59}$ |

**$B_s^0$  BRANCHING RATIOS** **$\Gamma(D_s^- \text{ anything})/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

| VALUE                        | EVTs | DOCUMENT ID               | TECN | COMMENT                           |
|------------------------------|------|---------------------------|------|-----------------------------------|
| <b>0.93±0.25 OUR AVERAGE</b> |      |                           |      |                                   |
| 0.91±0.18±0.41               |      | <sup>1</sup> DRUTSKOY 07  | BELL | $e^+e^- \rightarrow \Upsilon(4S)$ |
| 0.81±0.24±0.22               | 90   | <sup>2</sup> BUSKULIC 96E | ALEP | $e^+e^- \rightarrow Z$            |
| 1.56±0.58±0.44               | 147  | <sup>3</sup> ACTON 92N    | OPAL | $e^+e^- \rightarrow Z$            |

<sup>1</sup> The extraction of this result takes into account the correlation between the measurements of  $B(\Upsilon(5S) \rightarrow D_s^- X)$  and  $B(\Upsilon(5S) \rightarrow D^0 X)$ .

<sup>2</sup> BUSKULIC 96E separate  $c\bar{c}$  and  $b\bar{b}$  sources of  $D_s^+$  mesons using a lifetime tag, subtract generic  $\bar{b} \rightarrow W^+ \rightarrow D_s^+$  events, and obtain  $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \text{ anything}) = 0.088 \pm 0.020 \pm 0.020$  assuming  $B(D_s^- \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$  and PDG 1994 values for the relative partial widths to other  $D_s$  channels. We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s^- \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s^- \rightarrow \phi\pi)$ .

<sup>3</sup> ACTON 92N assume that excess of  $147 \pm 48 D_s^0$  events over that expected from  $B^0$ ,  $B^+$ , and  $c\bar{c}$  is all from  $B_s^0$  decay. The product branching fraction is measured to be  $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \text{ anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (5.9 \pm 1.9 \pm 1.1) \times 10^{-3}$ . We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s^- \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s^- \rightarrow \phi\pi)$ .

 **$\Gamma(\ell\nu_\ell X)/\Gamma_{\text{total}}$**  **$\Gamma_2/\Gamma$** 

| VALUE (units $10^{-2}$ )                    | DOCUMENT ID            | TECN | COMMENT                           |
|---------------------------------------------|------------------------|------|-----------------------------------|
| <b>9.6±0.8 OUR AVERAGE</b>                  |                        |      |                                   |
| 9.6±0.4±0.7                                 | <sup>1</sup> OSWALD 13 | BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |
| 9.5 <sup>+2.5+1.1</sup> <sub>-2.0-1.9</sub> | <sup>2</sup> LEES 12A  | BABR | $e^+e^-$                          |

<sup>1</sup> The measurement corresponds to the average of the electron and muon branching fractions.

<sup>2</sup> The measurement corresponds to a branching fraction where the lepton originates from bottom decay and is the average between the electron and muon branching fractions. LEES 12A uses the correlation of the production of  $\phi$  mesons in association with a lepton in  $e^+e^-$  data taken at center-of-mass energies between 10.54 and 11.2 GeV.

 **$\Gamma(e^+ \nu X^-)/\Gamma_{\text{total}}$**  **$\Gamma_3/\Gamma$** 

| VALUE (units $10^{-2}$ ) | DOCUMENT ID | TECN | COMMENT                           |
|--------------------------|-------------|------|-----------------------------------|
| <b>9.1±0.5±0.6</b>       | OSWALD 13   | BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |

 **$\Gamma(\mu^+ \nu X^-)/\Gamma_{\text{total}}$**  **$\Gamma_4/\Gamma$** 

| VALUE (units $10^{-2}$ ) | DOCUMENT ID | TECN | COMMENT                           |
|--------------------------|-------------|------|-----------------------------------|
| <b>10.2±0.6±0.8</b>      | OSWALD 13   | BELL | $e^+e^- \rightarrow \Upsilon(5S)$ |



$\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

The values and averages in this section serve only to show what values result if one assumes our  $B(\bar{b} \rightarrow B_s^0)$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determine  $B(\bar{b} \rightarrow B_s^0)$  as described in the note on “Production and Decay of  $b$ -Flavored Hadrons.”

| VALUE (units $10^{-2}$ )                                                      | EVTs | DOCUMENT ID           | TECN     | COMMENT                           |
|-------------------------------------------------------------------------------|------|-----------------------|----------|-----------------------------------|
| <b>8.1<math>\pm</math>1.3 OUR AVERAGE</b>                                     |      |                       |          |                                   |
| 8.2 $\pm$ 0.2 $\pm$ 1.5                                                       |      | <sup>1</sup> OSWALD   | 15 BELL  | $e^+e^- \rightarrow \Upsilon(5S)$ |
| 7.6 $\pm$ 1.2 $\pm$ 2.1                                                       | 134  | <sup>2</sup> BUSKULIC | 950 ALEP | $e^+e^- \rightarrow Z$            |
| 10.7 $\pm$ 4.3 $\pm$ 2.9                                                      |      | <sup>3</sup> ABREU    | 92M DLPH | $e^+e^- \rightarrow Z$            |
| 10.3 $\pm$ 3.6 $\pm$ 2.8                                                      | 18   | <sup>4</sup> ACTON    | 92N OPAL | $e^+e^- \rightarrow Z$            |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                       |          |                                   |
| 13 $\pm$ 4 $\pm$ 4                                                            | 27   | <sup>5</sup> BUSKULIC | 92E ALEP | $e^+e^- \rightarrow Z$            |

<sup>1</sup> Obtains  $B_s \rightarrow D_s X e \nu$ , and  $D_s X \mu \nu$  separately, then combines them by assuming systematic uncertainties are fully correlated, except for the one on lepton identification. The third uncertainty adds in quadrature systematic uncertainties from external sources (number of  $B_s$  events, and  $D_s^{(*)}$  branching fractions). OSWALD 15 also measures the cross-section  $\sigma(e^+e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = 53.8 \pm 1.4 \pm 5.3$  pb at  $\sqrt{s} = 10.86$  GeV.

<sup>2</sup> BUSKULIC 950 use  $D_s \ell$  correlations. The measured product branching ratio is  $B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) = (0.82 \pm 0.09^{+0.13}_{-0.14})\%$  assuming  $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$  and PDG 1994 values for the relative partial widths to the six other  $D_s$  channels used in this analysis. Combined with results from  $\Upsilon(4S)$  experiments this can be used to extract  $B(\bar{b} \rightarrow B_s) = (11.0 \pm 1.2^{+2.5}_{-2.6})\%$ . We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s \rightarrow \phi\pi)$ .

<sup>3</sup> ABREU 92M measured muons only and obtained product branching ratio  $B(Z \rightarrow b \text{ or } \bar{b}) \times B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s \mu^+ \nu_\mu \text{ anything}) \times B(D_s \rightarrow \phi\pi) = (18 \pm 8) \times 10^{-5}$ . We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s \rightarrow \phi\pi)$ . We use  $B(Z \rightarrow b \text{ or } \bar{b}) = 2B(Z \rightarrow b\bar{b}) = 2 \times (0.2212 \pm 0.0019)$ .

<sup>4</sup> ACTON 92N is measured using  $D_s \rightarrow \phi\pi^+$  and  $K^*(892)^0 K^+$  events. The product branching fraction measured is measured to be  $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$ . We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s \rightarrow \phi\pi)$ .

<sup>5</sup> BUSKULIC 92E is measured using  $D_s \rightarrow \phi\pi^+$  and  $K^*(892)^0 K^+$  events. They use  $2.7 \pm 0.7\%$  for the  $\phi\pi^+$  branching fraction. The average product branching fraction is measured to be  $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) = 0.020 \pm 0.0055^{+0.005}_{-0.006}$ . We evaluate using our current values  $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$  and  $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$ . Our first error is their experiment's and our second error is that due to  $B(\bar{b} \rightarrow B_s^0)$  and  $B(D_s \rightarrow \phi\pi)$ . Superseded by BUSKULIC 950.

$$\Gamma(D_s^{*-} \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_6 / \Gamma$$

| VALUE (units $10^{-2}$ )                | DOCUMENT ID         | TECN | COMMENT                               |
|-----------------------------------------|---------------------|------|---------------------------------------|
| <b><math>5.4 \pm 0.4 \pm 1.0</math></b> | <sup>1</sup> OSWALD | 15   | BELL $e^+ e^- \rightarrow \gamma(5S)$ |

<sup>1</sup> Obtains  $B_s \rightarrow D_s^* X e \nu$ , and  $D_s^* X \mu \nu$  separately, then combines them by assuming systematic uncertainties are fully correlated, except for the one on lepton identification. The third uncertainty adds in quadrature systematic uncertainties from external sources (number of  $B_s$  events, and  $D_s^{(*)}$  branching fractions). OSWALD 15 also measures the cross-section  $\sigma(e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = 53.8 \pm 1.4 \pm 5.3$  pb at  $\sqrt{s} = 10.86$  GeV.

$$\Gamma(D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0) / \Gamma_{\text{total}} \quad \Gamma_7 / \Gamma$$

| VALUE (units $10^{-3}$ )                | DOCUMENT ID         | TECN | COMMENT                   |
|-----------------------------------------|---------------------|------|---------------------------|
| <b><math>2.7 \pm 0.7 \pm 0.2</math></b> | <sup>1</sup> ABAZOV | 09G  | D0 $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> ABAZOV 09G reports  $[\Gamma(B_s^0 \rightarrow D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0) / \Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_s^0)] = (2.66 \pm 0.52 \pm 0.45) \times 10^{-4}$  which we divide by our best value  $B(\bar{b} \rightarrow B_s^0) = (10.0 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+) / \Gamma(D_s^- \ell^+ \nu_\ell \text{ anything}) \quad \Gamma_8 / \Gamma_5$$

| VALUE (units $10^{-2}$ )                | DOCUMENT ID | TECN | COMMENT            |
|-----------------------------------------|-------------|------|--------------------|
| <b><math>5.4 \pm 1.2 \pm 0.5</math></b> | AAIJ        | 11A  | LHCB $pp$ at 7 TeV |

$$\Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+) / \Gamma(D_s^- \ell^+ \nu_\ell \text{ anything}) \quad \Gamma_9 / \Gamma_5$$

| VALUE (units $10^{-2}$ )                | DOCUMENT ID | TECN | COMMENT            |
|-----------------------------------------|-------------|------|--------------------|
| <b><math>3.3 \pm 1.0 \pm 0.4</math></b> | AAIJ        | 11A  | LHCB $pp$ at 7 TeV |

$$\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+) / \Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+) \quad \Gamma_8 / \Gamma_9$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.61 \pm 0.14 \pm 0.05$  <sup>1</sup> AAIJ 11A LHCB  $pp$  at 7 TeV

<sup>1</sup> Not independent of other AAIJ 11A measurements.

$$\Gamma(D_s^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{10} / \Gamma$$

| VALUE (units $10^{-3}$ )                      | EVTS                   | DOCUMENT ID | TECN | COMMENT                          |
|-----------------------------------------------|------------------------|-------------|------|----------------------------------|
| <b><math>3.00 \pm 0.23</math> OUR FIT</b>     |                        |             |      |                                  |
| <b><math>2.99 \pm 0.24</math> OUR AVERAGE</b> |                        |             |      |                                  |
| $2.95 \pm 0.05^{+0.25}_{-0.28}$               | <sup>1</sup> AAIJ      | 12AG        | LHCB | $pp$ at 7 TeV                    |
| $3.6 \pm 0.5 \pm 0.5$                         | <sup>2</sup> LOUVOT    | 09          | BELL | $e^+ e^- \rightarrow \gamma(5S)$ |
| $2.8 \pm 0.6 \pm 0.1$                         | <sup>3</sup> ABULENCIA | 07C         | CDF  | $p\bar{p}$ at 1.96 TeV           |

• • • We do not use the following data for averages, fits, limits, etc. • • •

- |                 |                        |                    |      |                             |
|-----------------|------------------------|--------------------|------|-----------------------------|
| 6.8 ± 2.2 ± 1.6 | DRUTSKOY               | 07A                | BELL | Repl. by LOUVOT 09          |
| 3.3 ± 1.1 ± 0.2 | <sup>4</sup> ABULENCIA | 06J                | CDF  | Repl. by ABULENCIA 07C      |
| <130            | <sup>6</sup>           | <sup>5</sup> AKERS | 94J  | OPAL $e^+e^- \rightarrow Z$ |
| seen            | 1                      | BUSKULIC           | 93G  | ALEP $e^+e^- \rightarrow Z$ |
- <sup>1</sup> AAIJ 12AG reports  $(2.95 \pm 0.05 \pm 0.17^{+0.18}_{-0.22}) \times 10^{-3}$  where the last uncertainty comes from the semileptonic  $f_s/f_d$  measurement. We combined the systematics in quadrature.
- <sup>2</sup> LOUVOT 09 reports  $(3.67^{+0.35+0.65}_{-0.33-0.645}) \times 10^{-3}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] \times [B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$ , which we rescale to our best value  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>3</sup> ABULENCIA 07C reports  $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.13 \pm 0.08 \pm 0.23$  which we multiply by our best value  $B(B^0 \rightarrow D^- \pi^+) = (2.52 \pm 0.13) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>4</sup> ABULENCIA 06J reports  $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.32 \pm 0.18 \pm 0.38$  which we multiply by our best value  $B(B^0 \rightarrow D^- \pi^+) = (2.52 \pm 0.13) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>5</sup> AKERS 94J sees  $\leq 6$  events and measures the limit on the product branching fraction  $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow D_s^- \pi^+) < 1.3\%$  at CL = 90%. We divide by our current value  $B(\bar{b} \rightarrow B_s^0) = 0.105$ .

| $\Gamma(D_s^- \rho^+)/\Gamma(D_s^- \pi^+)$ | $\Gamma_{11}/\Gamma_{10}$ |      |         |                                   |
|--------------------------------------------|---------------------------|------|---------|-----------------------------------|
| VALUE                                      | DOCUMENT ID               | TECN | COMMENT |                                   |
| <b>2.3±0.4±0.2</b>                         | LOUVOT                    | 10   | BELL    | $e^+e^- \rightarrow \Upsilon(5S)$ |

| $\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ | $\Gamma_{12}/\Gamma$   |      |         |                        |
|---------------------------------------------------------|------------------------|------|---------|------------------------|
| VALUE (units $10^{-3}$ )                                | DOCUMENT ID            | TECN | COMMENT |                        |
| <b>6.1±1.0 OUR FIT</b>                                  |                        |      |         |                        |
| <b>6.3±1.5±0.7</b>                                      | <sup>1</sup> ABULENCIA | 07C  | CDF     | $p\bar{p}$ at 1.96 TeV |

- <sup>1</sup> ABULENCIA 07C reports  $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+ \pi^+ \pi^-)] = 1.05 \pm 0.10 \pm 0.22$  which we multiply by our best value  $B(B^0 \rightarrow D^- \pi^+ \pi^+ \pi^-) = (6.0 \pm 0.7) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+)$ | $\Gamma_{12}/\Gamma_{10}$ |      |         |               |
|-------------------------------------------------------|---------------------------|------|---------|---------------|
| VALUE                                                 | DOCUMENT ID               | TECN | COMMENT |               |
| <b>2.05±0.34 OUR FIT</b>                              |                           |      |         |               |
| <b>2.01±0.37±0.20</b>                                 | AAIJ                      | 11E  | LHCB    | $pp$ at 7 TeV |

| $\Gamma(D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$ | $\Gamma_{13}/\Gamma_{12}$ |      |         |               |
|--------------------------------------------------------------------------------------------------------|---------------------------|------|---------|---------------|
| VALUE (units $10^{-3}$ )                                                                               | DOCUMENT ID               | TECN | COMMENT |               |
| <b>4.0±1.0±0.4</b>                                                                                     | AAIJ                      | 12AX | LHCB    | $pp$ at 7 TeV |

$\Gamma(D_s^\mp K^\pm)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

**2.27 $\pm$ 0.19 OUR FIT**

|                                                                    |                     |    |                                        |
|--------------------------------------------------------------------|---------------------|----|----------------------------------------|
| <b>2.3 <math>^{+1.2}_{-1.0}</math> <math>^{+0.4}_{-0.3}</math></b> | <sup>1</sup> LOUVOT | 09 | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
|--------------------------------------------------------------------|---------------------|----|----------------------------------------|

<sup>1</sup> LOUVOT 09 reports  $(2.4^{+1.2}_{-1.0} \pm 0.42) \times 10^{-4}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow D_s^\mp K^\pm)/\Gamma_{\text{total}}] \times [B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$ , which we rescale to our best value  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(D_s^\mp K^\pm)/\Gamma(D_s^- \pi^+)$   $\Gamma_{14}/\Gamma_{10}$ 

| VALUE (units $10^{-2}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

**7.55 $\pm$ 0.24 OUR FIT****7.55 $\pm$ 0.24 OUR AVERAGE**

|                                                                               |          |      |                            |
|-------------------------------------------------------------------------------|----------|------|----------------------------|
| 7.52 $\pm$ 0.15 $\pm$ 0.19                                                    | AAIJ     | 15AC | LHCB $pp$ at 7, 8 TeV      |
| 9.7 $\pm$ 1.8 $\pm$ 0.9                                                       | AALTONEN | 09AQ | CDF $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |      |                            |
| 6.46 $\pm$ 0.43 $\pm$ 0.25                                                    | AAIJ     | 12AG | LHCB Repl. by AAIJ 15AC    |

 $\Gamma(D_s^- K^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$   $\Gamma_{15}/\Gamma_{12}$ 

| VALUE (units $10^{-2}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

|                                                  |      |      |                    |
|--------------------------------------------------|------|------|--------------------|
| <b>5.2<math>\pm</math>0.5<math>\pm</math>0.3</b> | AAIJ | 12AX | LHCB $pp$ at 7 TeV |
|--------------------------------------------------|------|------|--------------------|

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

| VALUE (units $10^{-3}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

**4.4 $\pm$ 0.5 OUR AVERAGE**

|                                                                               |              |          |      |                                        |
|-------------------------------------------------------------------------------|--------------|----------|------|----------------------------------------|
| 4.0 $\pm$ 0.2 $\pm$ 0.5                                                       | <sup>1</sup> | AAIJ     | 13AP | LHCB $pp$ at 7 TeV                     |
| 5.8 $^{+1.1}_{-0.9}$ $\pm$ 1.3                                                | <sup>2</sup> | ESEN     | 13   | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
| 5.4 $\pm$ 0.8 $\pm$ 0.8                                                       | <sup>3</sup> | AALTONEN | 12C  | CDF $p\bar{p}$ at 1.96 TeV             |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |              |          |      |                                        |
| 10.3 $^{+3.9}_{-3.2}$ $^{+2.6}_{-2.5}$                                        | <sup>4</sup> | ESEN     | 10   | BELL Repl. by ESEN 13                  |
| 10.4 $^{+3.5}_{-3.2}$ $\pm$ 1.1                                               | <sup>5</sup> | AALTONEN | 08F  | CDF Repl. by AALTONEN 12C              |
| <67                                                                           | 90           | DRUTSKOY | 07A  | BELL Repl. by ESEN 10                  |

<sup>1</sup> Uses  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ .

<sup>2</sup> Use  $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$  decays assuming  $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$  and  $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$ .

<sup>3</sup> AALTONEN 12C reports  $(f_s/f_d) (B(B_s^0 \rightarrow D_s^+ D_s^-) / B(B^0 \rightarrow D^- D_s^+)) = 0.183 \pm 0.021 \pm 0.017$ . We multiply this result by our best value of  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$  and divide by our best value of  $f_s/f_d$ , where  $1/2 f_s/f_d = 0.1230 \pm 0.0115$ . Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

<sup>4</sup> Uses  $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

<sup>5</sup> AALTONEN 08F reports  $[\Gamma(B_s^0 \rightarrow D_s^+ D_s^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.44^{+0.48}_{-0.44}$  which we multiply by our best value  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(D_s^- D^+)/\Gamma_{\text{total}}$ $\Gamma_{17}/\Gamma$

| VALUE (units $10^{-4}$ )                | DOCUMENT ID       | TECN      | COMMENT            |
|-----------------------------------------|-------------------|-----------|--------------------|
| <b><math>2.8 \pm 0.4 \pm 0.3</math></b> | <sup>1</sup> AAIJ | 14AA LHCB | $pp$ at 7 TeV      |
| $3.6 \pm 0.6 \pm 0.5$                   | <sup>2</sup> AAIJ | 13AP LHCB | Repl. by AAIJ 14AA |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 14AA reports  $[\Gamma(B_s^0 \rightarrow D_s^- D^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 0.038 \pm 0.004 \pm 0.003$  which we multiply by our best value  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

<sup>2</sup> Uses  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ .

### $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma$

| VALUE (units $10^{-4}$ )                | DOCUMENT ID       | TECN      | COMMENT       |
|-----------------------------------------|-------------------|-----------|---------------|
| <b><math>2.2 \pm 0.4 \pm 0.4</math></b> | <sup>1</sup> AAIJ | 13AP LHCB | $pp$ at 7 TeV |

<sup>1</sup> Uses  $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$  and  $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$ .

### $\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$ $\Gamma_{19}/\Gamma$

| VALUE (units $10^{-4}$ )                | DOCUMENT ID       | TECN      | COMMENT       |
|-----------------------------------------|-------------------|-----------|---------------|
| <b><math>1.9 \pm 0.3 \pm 0.4</math></b> | <sup>1</sup> AAIJ | 13AP LHCB | $pp$ at 7 TeV |

<sup>1</sup> Uses  $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$  and  $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$ .

### $\Gamma(D_s^{*-} \pi^+)/\Gamma(D_s^- \pi^+)$ $\Gamma_{20}/\Gamma_{10}$

| VALUE                                             | DOCUMENT ID | TECN    | COMMENT                          |
|---------------------------------------------------|-------------|---------|----------------------------------|
| <b><math>0.65^{+0.15}_{-0.13} \pm 0.07</math></b> | LOUVOT      | 10 BELL | $e^+ e^- \rightarrow \gamma(5S)$ |

### $\Gamma(D_s^{*\mp} K^\pm)/\Gamma(D_s^{*-} \pi^+)$ $\Gamma_{21}/\Gamma_{20}$

| VALUE                                                 | DOCUMENT ID | TECN      | COMMENT          |
|-------------------------------------------------------|-------------|-----------|------------------|
| <b><math>0.068 \pm 0.005^{+0.003}_{-0.002}</math></b> | AAIJ        | 15AD LHCB | $pp$ at 7, 8 TeV |

### $\Gamma(D_s^{*-} \rho^+)/\Gamma(D_s^- \pi^+)$ $\Gamma_{22}/\Gamma_{10}$

| VALUE                                   | DOCUMENT ID | TECN    | COMMENT                          |
|-----------------------------------------|-------------|---------|----------------------------------|
| <b><math>3.2 \pm 0.6 \pm 0.3</math></b> | LOUVOT      | 10 BELL | $e^+ e^- \rightarrow \gamma(5S)$ |

### $\Gamma(D_s^{*-} \rho^+)/\Gamma(D_s^- \rho^+)$ $\Gamma_{22}/\Gamma_{11}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4 \pm 0.3 \pm 0.1$  <sup>1</sup> LOUVOT 10 BELL  $e^+ e^- \rightarrow \gamma(5S)$

<sup>1</sup> Not independent of other LOUVOT 10 measurements.

$$[\Gamma(D_s^{*+} D_s^-) + \Gamma(D_s^{*-} D_s^+)] / \Gamma_{\text{total}} \quad \Gamma_{23} / \Gamma$$

| VALUE (units $10^{-3}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

**13.9 ± 1.7 OUR AVERAGE**

13.6 ± 1.0 ± 1.4

1 AAIJ 16P LHC  $pp$  at 7 TeV17.6<sup>+2.3</sup><sub>-2.2</sub> ± 4.02 ESEN 13 BELL  $e^+e^- \rightarrow \Upsilon(5S)$ 

12.5 ± 1.7 ± 1.8

3 AALTONEN 12C CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

27.5<sup>+8.3</sup><sub>-7.1</sub> ± 6.9

4 ESEN 10 BELL Repl. by ESEN 13

&lt;121

90

DRUTSKOY 07A BELL Repl. by ESEN 10

<sup>1</sup> AAIJ 16P reports  $[\Gamma(B_s^0 \rightarrow D_s^{*+} D_s^-) + \Gamma(D_s^{*-} D_s^+)] / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.88 \pm 0.08 \pm 0.12$  which we multiply by our best value  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Use  $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$  decays assuming  $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$  and  $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$ .

<sup>3</sup> AALTONEN 12C reports  $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^- + D_s^{*-} D_s^+) / B(B^0 \rightarrow D^- D_s^+)) = 0.424 \pm 0.046 \pm 0.035$ . We multiply this result by our best value of  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$  and divide by our best value of  $f_s/f_d$ , where  $1/2 f_s/f_d = 0.1230 \pm 0.0115$ . Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

<sup>4</sup> Uses  $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

$$\Gamma(D_s^{*+} D_s^{*-}) / \Gamma_{\text{total}} \quad \Gamma_{24} / \Gamma$$

| VALUE (units $10^{-3}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

**14.4 ± 2.1 OUR AVERAGE**

Error includes scale factor of 1.1.

12.7 ± 1.3 ± 1.4

1 AAIJ 16P LHC  $pp$  at 7 TeV19.8<sup>+3.3</sup><sub>-3.1</sub> ± 5.22 ESEN 13 BELL  $e^+e^- \rightarrow \Upsilon(5S)$ 

19.2 ± 2.9 ± 2.7

3 AALTONEN 12C CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

30.8<sup>+12.2</sup><sub>-10.4</sub> ± 8.5

4 ESEN 10 BELL Repl. by ESEN 13

&lt;257

90

DRUTSKOY 07A BELL Repl. by ESEN 10

<sup>1</sup> AAIJ 16P reports  $[\Gamma(B_s^0 \rightarrow D_s^{*+} D_s^{*-}) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.76 \pm 0.11 \pm 0.14$  which we multiply by our best value  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Use  $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$  decays assuming  $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$  and  $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$ .

<sup>3</sup> AALTONEN 12C reports  $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^{*-}) / B(B^0 \rightarrow D^- D_s^+)) = 0.654 \pm 0.072 \pm 0.065$ . We multiply this result by our best value of  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$  and divide by our best value of  $f_s/f_d$ , where  $1/2 f_s/f_d = 0.1230 \pm 0.0115$ .

Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.

<sup>4</sup> Uses  $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

$\Gamma(D_s^{(*)+} D_s^{(*)-}) / \Gamma_{\text{total}}$   $\Gamma_{25} / \Gamma$

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) and are described at <https://hflav.web.cern.ch/>. The averaging/rescaling procedure takes into account correlations between the measurements.

| VALUE (%)                                                                     | CL% | DOCUMENT ID           | TECN | COMMENT                                |
|-------------------------------------------------------------------------------|-----|-----------------------|------|----------------------------------------|
| <b>4.5 ± 1.4 OUR EVALUATION</b>                                               |     |                       |      |                                        |
| <b>3.4 ± 0.4 OUR AVERAGE</b>                                                  |     |                       |      |                                        |
| 3.07 ± 0.22 ± 0.33                                                            |     | <sup>1</sup> AAIJ     | 16P  | LHCB $pp$ at 7 TeV                     |
| 4.32 <sup>+0.42+1.04</sup> <sub>-0.39-1.03</sub>                              |     | <sup>2</sup> ESEN     | 13   | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |
| 3.7 ± 0.4 ± 0.5                                                               |     | <sup>3</sup> AALTONEN | 12C  | CDF $p\bar{p}$ at 1.96 TeV             |
| 3.5 ± 1.0 ± 1.1                                                               |     | <sup>4</sup> ABAZOV   | 09I  | D0 $p\bar{p}$ at 1.96 TeV              |
| 14 ± 6 ± 3                                                                    |     | <sup>5,6</sup> BARATE | 00K  | ALEP $e^+e^- \rightarrow Z$            |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                       |      |                                        |
| 6.85 <sup>+1.53+1.79</sup> <sub>-1.30-1.80</sub>                              |     | <sup>7,8</sup> ESEN   | 10   | BELL Repl. by ESEN 13                  |
| 3.9 <sup>+1.9+1.6</sup> <sub>-1.7-1.5</sub>                                   |     | <sup>4</sup> ABAZOV   | 07Y  | D0 Repl. by ABAZOV 09I                 |
| <0.218                                                                        | 90  | BARATE                | 98Q  | ALEP $e^+e^- \rightarrow Z$            |

<sup>1</sup> AAIJ 16P reports  $[\Gamma(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 4.24 \pm 0.14 \pm 0.27$  which we multiply by our best value  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Use  $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$  decays assuming  $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$  and  $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$ .

<sup>3</sup> AALTONEN 12C reports  $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) / B(B^0 \rightarrow D^- D_s^+)) = 1.261 \pm 0.095 \pm 0.112$ . We multiply this result by our best value of  $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$  and divide by our best value of  $f_s/f_d$ , where  $1/2 f_s/f_d = 0.1230 \pm 0.0115$ . Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.

<sup>4</sup> Uses the final states where  $D_s^+ \rightarrow \phi \pi^+$  and  $D_s^- \rightarrow \phi \mu^- \bar{\nu}_\mu$ .

<sup>5</sup> Reports  $B(B_s^0(\text{short}) \rightarrow D_s^{(*)+} D_s^{(*)-}) = (0.23 \pm 0.10 \pm 0.05) \cdot [0.17/B(D_s \rightarrow \phi \chi)]^2$  assuming  $B(B_s^0 \rightarrow B_s^0(\text{short})) = 50\%$ . We use our best value of  $B(D_s \rightarrow \phi \chi) = 15.7 \pm 1.0\%$  to obtain the quoted result.

<sup>6</sup> Uses  $\phi\phi$  correlations from  $B_s^0(\text{short}) \rightarrow D_s^{(*)+} D_s^{(*)-}$ .

<sup>7</sup> Sum of exclusive  $B_s \rightarrow D_s^+ D_s^-$ ,  $B_s \rightarrow D_s^{*\pm} D_s^\mp$  and  $B_s \rightarrow D_s^{*+} D_s^{*-}$ .

<sup>8</sup> Uses  $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$  assuming  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

$\Gamma(\bar{D}^{*0}\bar{K}^0)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$ 

| VALUE (units $10^{-4}$ )                | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------------------------|-------------------|----------|------------------|
| <b><math>2.8 \pm 1.0 \pm 0.5</math></b> | <sup>1</sup> AAIJ | 16C LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured and normalized to the  $B_S^0 \rightarrow \bar{D}^{*0} K_S^0$  decay with  $f_s/f_d = 0.259 \pm 0.015$ . Signal significance is 4.4 standard deviations.

 $\Gamma(\bar{D}^0\bar{K}^0)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

| VALUE (units $10^{-4}$ )                | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------------------------|-------------------|----------|------------------|
| <b><math>4.3 \pm 0.5 \pm 0.7</math></b> | <sup>1</sup> AAIJ | 16C LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured and normalized to the  $B^0 \rightarrow \bar{D}^0 K_S^0$  decay with  $f_s/f_d = 0.259 \pm 0.015$ .

 $\Gamma(\bar{D}^0 K^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

| VALUE (units $10^{-4}$ )                 | DOCUMENT ID       | TECN      | COMMENT       |
|------------------------------------------|-------------------|-----------|---------------|
| <b><math>10.4 \pm 1.1 \pm 0.5</math></b> | <sup>1</sup> AAIJ | 13AQ LHCb | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 13AQ reports  $[\Gamma(B_S^0 \rightarrow \bar{D}^0 K^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)] = 1.18 \pm 0.05 \pm 0.12$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.8 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

| VALUE (units $10^{-4}$ )                    | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------------------|-------------------|-----------|------------------|
| <b><math>4.4 \pm 0.6</math> OUR AVERAGE</b> |                   |           |                  |
| $4.29 \pm 0.09 \pm 0.65$                    | <sup>1</sup> AAIJ | 14BH LHCb | $pp$ at 7, 8 TeV |
| $4.7 \pm 1.2 \pm 0.3$                       | <sup>2</sup> AAIJ | 11D LHCb  | $pp$ at 7 TeV    |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |                   |           |                    |
|-----------------------|-------------------|-----------|--------------------|
| $3.5 \pm 0.4 \pm 0.4$ | <sup>3</sup> AAIJ | 13BX LHCb | Repl. by AAIJ 14BH |
|-----------------------|-------------------|-----------|--------------------|

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

<sup>2</sup> AAIJ 11D reports  $[\Gamma(B_S^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \rho^0)] = 1.48 \pm 0.34 \pm 0.19$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 \rho^0) = (3.21 \pm 0.21) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> AAIJ 13BX reports  $[\Gamma(B_S^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^*(892)^0)] = 7.8 \pm 0.7 \pm 0.3 \pm 0.6$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 K^*(892)^0) = (4.5 \pm 0.6) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^0 \bar{K}^*(1410))/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID       | TECN      | COMMENT          |
|--------------------------------------------|-------------------|-----------|------------------|
| <b><math>38.6 \pm 11.4 \pm 33.3</math></b> | <sup>1</sup> AAIJ | 14BH LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 \bar{K}_0^*(1430))/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

| VALUE (units $10^{-5}$ )                 | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------|-------------------|-----------|------------------|
| <b><math>30.0 \pm 2.4 \pm 6.8</math></b> | <sup>1</sup> AAIJ | 14BH LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays. Corresponds to the resonant  $K_0^*(1430)$  part of LASS parametrization.



$\Gamma(\bar{D}^0 K_2^*(1430))/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

| VALUE (units $10^{-5}$ )                 | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>11.1 \pm 1.8 \pm 3.8</math></b> |     | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 K^*(1680))/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$ 

| VALUE (units $10^{-5}$ )     | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 7.8</math></b> | 90  | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 \bar{K}_0^*(1950))/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$ 

| VALUE (units $10^{-5}$ )    | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 11</math></b> | 90  | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 \bar{K}_3^*(1780))/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

| VALUE (units $10^{-5}$ )     | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 2.6</math></b> | 90  | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 \bar{K}_4^*(2045))/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$ 

| VALUE (units $10^{-5}$ )     | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 3.1</math></b> | 90  | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 K^- \pi^+ (\text{non-resonant}))/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

| VALUE (units $10^{-5}$ )                 | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>20.6 \pm 3.8 \pm 7.3</math></b> |     | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays. Corresponds to the non-resonant part of the LASS parametrization.

 $\Gamma(D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \bar{D}^0 K^-)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$ 

| VALUE (units $10^{-5}$ )                 | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>25.7 \pm 0.7 \pm 4.0</math></b> |     | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>1.6 \pm 0.4 \pm 0.7</math></b> |     | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>5.0 \pm 1.2 \pm 3.4</math></b> |     | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

$\Gamma(D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------|-------------------|-----------|------------------|
| <b><math>2.2 \pm 0.1 \pm 0.6</math></b> | <sup>1</sup> AAIJ | 14BH LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses Dalitz plot analysis of  $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$  decays.

 $\Gamma(\bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------|-------------------|-----------|------------------|
| <b><math>5.5 \pm 0.7 \pm 0.5</math></b> | <sup>1</sup> AAIJ | 18AZ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |                     |          |                    |
|-----------------------|---------------------|----------|--------------------|
| $5.3 \pm 2.0 \pm 0.5$ | <sup>2,3</sup> AAIJ | 12AMLHCB | Repl. by AAIJ 18AZ |
|-----------------------|---------------------|----------|--------------------|

<sup>1</sup> AAIJ 18AZ reports  $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^+ K^-)] = 0.930 \pm 0.089 \pm 0.069$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 K^+ K^-) = (5.9 \pm 0.5) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AAIJ 12AM reports  $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^+ K^-)] = 0.90 \pm 0.27 \pm 0.20$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 K^+ K^-) = (5.9 \pm 0.5) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Uses  $B(b \rightarrow B_s^0) / B(b \rightarrow B^0) = 0.267^{+0.023}_{-0.020}$  measured by the same authors.

 $\Gamma(\bar{D}^0 f_0(980)) / \Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$ 

| VALUE                                       | CL% | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------------------|-----|-------------|-----------|------------------|
| <b><math>&lt; 3.1 \times 10^{-6}</math></b> | 90  | AAIJ        | 15AG LHCB | $pp$ at 7, 8 TeV |

 $\Gamma(\bar{D}^0 \phi) / \Gamma(\bar{D}^0 \bar{K}^*(892)^0)$   $\Gamma_{44}/\Gamma_{29}$ 

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |      |           |                    |
|-----------------------------|------|-----------|--------------------|
| $0.069 \pm 0.013 \pm 0.007$ | AAIJ | 13BX LHCB | Repl. by AAIJ 18AY |
|-----------------------------|------|-----------|--------------------|

 $\Gamma(\bar{D}^0 \phi) / \Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | DOCUMENT ID       | TECN      | COMMENT             |
|-----------------------------------------|-------------------|-----------|---------------------|
| <b><math>3.0 \pm 0.4 \pm 0.2</math></b> | <sup>1</sup> AAIJ | 18AY LHCB | $pp$ at 7 and 8 TeV |

<sup>1</sup> AAIJ 18AY reports  $[\Gamma(B_s^0 \rightarrow \bar{D}^0 \phi) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)] = (3.4 \pm 0.4 \pm 0.3) \times 10^{-2}$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.8 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^{*0} \phi) / \Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$ 

| VALUE (units $10^{-5}$ )                | DOCUMENT ID       | TECN      | COMMENT             |
|-----------------------------------------|-------------------|-----------|---------------------|
| <b><math>3.7 \pm 0.6 \pm 0.2</math></b> | <sup>1</sup> AAIJ | 18AY LHCB | $pp$ at 7 and 8 TeV |

<sup>1</sup> AAIJ 18AY reports  $[\Gamma(B_s^0 \rightarrow \bar{D}^{*0} \phi) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)] = (4.2 \pm 0.5 \pm 0.4) \times 10^{-2}$  which we multiply by our best value  $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.8 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

## $\Gamma(D^{*\mp}\pi^\pm)/\Gamma_{\text{total}}$ $\Gamma_{46}/\Gamma$

| VALUE                            | CL% | DOCUMENT ID       | TECN      | COMMENT       |
|----------------------------------|-----|-------------------|-----------|---------------|
| <b>&lt;6.1 × 10<sup>-6</sup></b> | 90  | <sup>1</sup> AAIJ | 13AL LHCb | $pp$ at 7 TeV |

<sup>1</sup> Uses  $f_s/f_d = 0.256 \pm 0.020$  and  $B(B^0 \rightarrow D^{*-}\pi^+) = (2.76 \pm 0.13) \times 10^{-3}$ .

## $\Gamma(\eta_c\phi)/\Gamma_{\text{total}}$ $\Gamma_{47}/\Gamma$

| VALUE (units 10 <sup>-4</sup> ) | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------|-------------------|----------|------------------|
| <b>5.01±0.53±0.68</b>           | <sup>1</sup> AAIJ | 17U LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> The last uncertainty includes the limited knowledge of the external branching fractions where the  $\eta_c$  is reconstructed in the  $p\bar{p}, K^+K^-\pi^+\pi^-$ ,  $\pi^+\pi^-\pi^+\pi^-$ , and  $K^+K^-K^+K^-$  decays and  $\phi(1020) \rightarrow K^+K^-$ .

## $\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{48}/\Gamma$

| VALUE (units 10 <sup>-4</sup> ) | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------|-------------------|----------|------------------|
| <b>1.76±0.59±0.31</b>           | <sup>1</sup> AAIJ | 17U LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> The last uncertainty includes the limited knowledge of the external branching fractions where the  $\eta_c$  is reconstructed in the  $p\bar{p}, K^+K^-\pi^+\pi^-$ ,  $\pi^+\pi^-\pi^+\pi^-$ , and  $K^+K^-K^+K^-$  decays. The significance of the signal, including systematic uncertainties, is 4.6 standard deviations.

## $\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}}$ $\Gamma_{49}/\Gamma$

| VALUE (units 10 <sup>-3</sup> ) | EVTS         | DOCUMENT ID | TECN      | COMMENT                           |
|---------------------------------|--------------|-------------|-----------|-----------------------------------|
| <b>1.08 ± 0.08 OUR FIT</b>      |              |             |           |                                   |
| <b>1.10 ± 0.09 OUR AVERAGE</b>  |              |             |           |                                   |
| 1.050±0.013±0.104               | <sup>1</sup> | AAIJ        | 13AN LHCb | $pp$ at 7 TeV                     |
| 1.25 ± 0.07 ± 0.23              | <sup>2</sup> | THORNE      | 13 BELL   | $e^+e^- \rightarrow \Upsilon(5S)$ |
| 1.5 ± 0.5 ± 0.1                 | <sup>3</sup> | ABE         | 96Q CDF   | $p\bar{p}$                        |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|      |              |                    |          |                        |
|------|--------------|--------------------|----------|------------------------|
| <6   | <sup>1</sup> | 4 AKERS            | 94J OPAL | $e^+e^- \rightarrow Z$ |
| seen | 14           | <sup>5</sup> ABE   | 93F CDF  | $p\bar{p}$ at 1.8 TeV  |
| seen | 1            | <sup>6</sup> ACTON | 92N OPAL | Sup. by AKERS 94J      |

<sup>1</sup> Uses  $f_s/f_d = 0.256 \pm 0.020$  and  $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$ .

<sup>2</sup> Uses  $f_s = (17.2 \pm 3.0)\%$  as the fraction of  $\Upsilon(5S)$  decaying to  $B_s^{(*)}\bar{B}_s^{(*)}$ .

<sup>3</sup> ABE 96Q reports  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}] \times [\Gamma(\bar{b} \rightarrow B_s^0)/[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)]] = (0.185 \pm 0.055 \pm 0.020) \times 10^{-3}$  which we divide by our best value  $\Gamma(\bar{b} \rightarrow B_s^0)/[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)] = 0.1230 \pm 0.0115$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> AKERS 94J sees one event and measures the limit on the product branching fraction  $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) < 7 \times 10^{-4}$  at CL = 90%. We divide by  $B(\bar{b} \rightarrow B_s^0) = 0.112$ .

<sup>5</sup> ABE 93F measured using  $J/\psi(1S) \rightarrow \mu^+\mu^-$  and  $\phi \rightarrow K^+K^-$ .

<sup>6</sup> In ACTON 92N a limit on the product branching fraction is measured to be  $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) \leq 0.22 \times 10^{-2}$ .

$\Gamma(J/\psi(1S)\phi\phi)/\Gamma(J/\psi(1S)\phi)$  $\Gamma_{50}/\Gamma_{49}$ 

| VALUE (units $10^{-2}$ )                          | EVTS | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------------------------|------|-------------------|----------|------------------|
| <b><math>1.15 \pm 0.12^{+0.05}_{-0.09}</math></b> | 128  | <sup>1</sup> AAIJ | 16U LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $J/\psi \rightarrow \mu^+\mu^-$ ,  $\phi \rightarrow K^+K^-$  decays, and observes  $128 \pm 13$  events of  $B_S^0 \rightarrow J/\psi\phi\phi$ .

 $\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{51}/\Gamma$ 

| VALUE                                      | CL% | DOCUMENT ID           | TECN   |
|--------------------------------------------|-----|-----------------------|--------|
| <b><math>&lt;1.2 \times 10^{-3}</math></b> | 90  | <sup>1</sup> ACCIARRI | 97C L3 |

<sup>1</sup> ACCIARRI 97C assumes  $B^0$  production fraction ( $39.5 \pm 4.0\%$ ) and  $B_S$  ( $12.0 \pm 3.0\%$ ).

 $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$  $\Gamma_{52}/\Gamma$ 

| VALUE (units $10^{-4}$ )                    | CL% | DOCUMENT ID                         | TECN     | COMMENT                         |
|---------------------------------------------|-----|-------------------------------------|----------|---------------------------------|
| <b><math>4.0 \pm 0.7</math> OUR AVERAGE</b> |     | Error includes scale factor of 1.4. |          |                                 |
| 3.6 $^{+0.5}_{-0.6}$ $^{+0.3}_{-0.2}$       |     | <sup>1</sup> AAIJ                   | 13A LHCb | $pp$ at 7 TeV                   |
| 5.10 $\pm 0.50^{+1.17}_{-0.83}$             |     | <sup>2</sup> LI                     | 12 BELL  | $e^+e^- \rightarrow \gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<38$  90 <sup>3</sup> ACCIARRI 97C L3

<sup>1</sup> AAIJ 13A reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\eta)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 14.0 \pm 1.2^{+1.1+1.1}_{-1.5-1.0}$  which we multiply by our best value  $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.55^{+0.18}_{-0.16}) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Observed for the first time with significances over  $10\sigma$ . The second error are total systematic uncertainties including the error on  $N(B_S^{(*)}\bar{B}_S^{(*)})$ .

<sup>3</sup> ACCIARRI 97C assumes  $B^0$  production fraction ( $39.5 \pm 4.0\%$ ) and  $B_S$  ( $12.0 \pm 3.0\%$ ).

 $\Gamma(J/\psi(1S)K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{53}/\Gamma$ 

| VALUE (units $10^{-5}$ )                      | DOCUMENT ID           | TECN      | COMMENT                |
|-----------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>1.88 \pm 0.15</math> OUR AVERAGE</b> |                       |           |                        |
| 1.87 $\pm 0.14 \pm 0.06$                      | <sup>1</sup> AAIJ     | 15AL LHCb | $pp$ at 7, 8 TeV       |
| 1.9 $\pm 0.4 \pm 0.2$                         | <sup>2</sup> AALTONEN | 11A CDF   | $p\bar{p}$ at 1.96 TeV |
| 1.98 $\pm 0.16 \pm 0.20$                      | <sup>3</sup> AAIJ     | 13AB LHCb | Repl. by AAIJ 15AL     |
| 1.98 $\pm 0.25 \pm 0.20$                      | <sup>4</sup> AAIJ     | 120 LHCb  | Repl. by AAIJ 13AB     |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 15AL reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K_S^0)] = (4.31 \pm 0.17 \pm 0.12 \pm 0.25) \times 10^{-2}$  which we multiply by our best value  $B(B^0 \rightarrow J/\psi(1S)K_S^0) = (4.34 \pm 0.15) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AALTONEN 11A reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_S^0)] / [B(\bar{b} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K_S^0)] = (1.09 \pm 0.19 \pm 0.11) \times 10^{-2}$  which we multiply or divide by our best values  $B(\bar{b} \rightarrow B_S^0) = (10.0 \pm 0.8) \times 10^{-2}$ ,  $B(\bar{b} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ ,  $B(B^0 \rightarrow J/\psi(1S)K_S^0) = 1/2 \times B(B^0 \rightarrow J/\psi(1S)K^0) = 1/2 \times (8.68 \pm 0.30) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>3</sup> AAIJ 13AB reports  $(1.97 \pm 0.14 \pm 0.07 \pm 0.15 \pm 0.08) \times 10^{-5}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0)]$  assuming  $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.98 \pm 0.35) \times 10^{-4}$ ,  $\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0) = 0.256 \pm 0.020$ , which we rescale to our best values  $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.68 \pm 0.30) \times 10^{-4}$ ,  $\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>4</sup> AAIJ 120 reports  $(1.83 \pm 0.21 \pm 0.10 \pm 0.14 \pm 0.07) \times 10^{-5}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0)]$  assuming  $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.71 \pm 0.32) \times 10^{-4}$ ,  $\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0) = 0.267^{+0.021}_{-0.02}$ , which we rescale to our best values  $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.68 \pm 0.30) \times 10^{-4}$ ,  $\Gamma(\bar{B} \rightarrow B_S^0)/\Gamma(\bar{B} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

### $\Gamma(J/\psi(1S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}$

$\Gamma_{54}/\Gamma$

| VALUE (units $10^{-5}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b>4.14 ± 0.18 ± 0.35</b>                                                     | <sup>1</sup> AAIJ     | 15AV LHCb | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| 4.4 $^{+0.5}_{-0.4} \pm 0.8$                                                  | <sup>2</sup> AAIJ     | 12AP LHCb | Repl. by AAIJ 15AV     |
| 9 $\pm 4 \pm 1$                                                               | <sup>3</sup> AALTONEN | 11A CDF   | $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> AAIJ 15AV result combines two measurements with different normalizing modes of  $B^0 \rightarrow J/\psi K^*(892)^0$  and  $B_S^0 \rightarrow J/\psi \phi$ .

<sup>2</sup> AAIJ 12AP reports  $B(B_S^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0)/B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (3.43^{+0.34}_{-0.36} \pm 0.50) \times 10^{-2}$  and  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.29 \pm 0.05 \pm 0.13) \times 10^{-3}$  after correcting for the contribution from  $K\pi$  S-wave beneath the  $K^*$  peak.

<sup>3</sup> AALTONEN 11A reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\bar{B} \rightarrow B_S^0)] / [B(\bar{B} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] = 0.0168 \pm 0.0024 \pm 0.0068$  which we multiply or divide by our best values  $B(\bar{B} \rightarrow B_S^0) = (10.0 \pm 0.8) \times 10^{-2}$ ,  $B(\bar{B} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ ,  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

### $\Gamma(J/\psi(1S)\eta')/\Gamma_{\text{total}}$

$\Gamma_{55}/\Gamma$

| VALUE (units $10^{-4}$ )       | DOCUMENT ID       | TECN     | COMMENT                           |
|--------------------------------|-------------------|----------|-----------------------------------|
| <b>3.3 ± 0.4 OUR AVERAGE</b>   |                   |          |                                   |
| 3.2 $^{+0.4}_{-0.5} \pm 0.2$   | <sup>1</sup> AAIJ | 13A LHCb | $pp$ at 7 TeV                     |
| 3.71 ± 0.61 $^{+0.85}_{-0.60}$ | <sup>2</sup> LI   | 12 BELL  | $e^+e^- \rightarrow \Upsilon(4S)$ |

<sup>1</sup> AAIJ 13A reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\eta')/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 12.7 \pm 1.1^{+0.5+1.0}_{-1.3-0.9}$  which we multiply by our best value  $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.55^{+0.18}_{-0.16}) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Observed for the first time with significances over  $10\sigma$ . The second error are total systematic uncertainties including the error on  $N(B_S^{(*)}\bar{B}_S^{(*)})$ .

$\Gamma(J/\psi(1S)\eta')/\Gamma(J/\psi(1S)\eta)$  $\Gamma_{55}/\Gamma_{52}$ 

| VALUE                          | DOCUMENT ID       | TECN     | COMMENT                         |
|--------------------------------|-------------------|----------|---------------------------------|
| <b>0.87 ± 0.06 OUR AVERAGE</b> |                   |          |                                 |
| 0.902 ± 0.072 ± 0.045          | <sup>1</sup> AAIJ | 15D LHCb | $pp$ at 7, 8 TeV                |
| 0.90 ± 0.09 $^{+0.06}_{-0.02}$ | <sup>2</sup> AAIJ | 13A LHCb | $pp$ at 7 TeV                   |
| 0.73 ± 0.14 ± 0.02             | <sup>2</sup> LI   | 12 BELL  | $e^+e^- \rightarrow \gamma(4S)$ |

<sup>1</sup> Uses  $J/\psi \rightarrow \mu^+\mu^-$ ,  $\eta' \rightarrow \rho^0\gamma$ , and  $\eta' \rightarrow \eta\pi^+\pi^-$  decays.<sup>2</sup> Strongly correlated with measurements of  $\Gamma(J/\psi(1S)\eta)/\Gamma$  and  $\Gamma(J/\psi(1S)\eta')/\Gamma$  reported in the same reference. $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\phi)$  $\Gamma_{56}/\Gamma_{49}$ 

| VALUE (units $10^{-2}$ )                                      | DOCUMENT ID       | TECN      | COMMENT       |
|---------------------------------------------------------------|-------------------|-----------|---------------|
| <b>19.4 ± 1.5 OUR FIT</b> Error includes scale factor of 2.2. |                   |           |               |
| <b>19.9 ± 0.7 ± 0.2</b>                                       | <sup>1</sup> AAIJ | 12A0 LHCb | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 12A0 reports  $(19.79 \pm 0.47 \pm 0.52) \times 10^{-2}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{59}/\Gamma$ 

| VALUE (units $10^{-4}$ )                                       | DOCUMENT ID     | TECN    | COMMENT                         |
|----------------------------------------------------------------|-----------------|---------|---------------------------------|
| <b>1.28 ± 0.18 OUR FIT</b> Error includes scale factor of 1.7. |                 |         |                                 |
| <b>1.16 <math>^{+0.31}_{-0.19}</math> ± 0.30</b>               | <sup>1</sup> LI | 11 BELL | $e^+e^- \rightarrow \gamma(5S)$ |

<sup>1</sup> The second error includes both the detector systematic and the uncertainty in the number of produced  $Y(5S) \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$  pairs. $\Gamma(J/\psi(1S)f_0(500), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)$  $\Gamma_{57}/\Gamma_{59}$ 

| VALUE            | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------|-----|-------------------|-----------|------------------|
| <b>&lt;0.034</b> | 90  | <sup>1</sup> AAIJ | 14BR LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis. $\Gamma(J/\psi(1S)\rho, \rho \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_{58}/\Gamma_{56}$ 

| VALUE            | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------|-----|-------------------|-----------|------------------|
| <b>&lt;0.017</b> | 90  | <sup>1</sup> AAIJ | 14BR LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis. $\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_{59}/\Gamma_{56}$ 

| VALUE                                                                                 | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------------------------------------------------------------|-------------------|-----------|------------------|
| <b>0.61 <math>^{+0.06}_{-0.07}</math> OUR FIT</b> Error includes scale factor of 2.1. |                   |           |                  |
| <b>0.703 ± 0.015 <math>^{+0.004}_{-0.051}</math></b>                                  | <sup>1</sup> AAIJ | 14BR LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_0, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{61}/\Gamma_{56}$$

| VALUE (%)                | DOCUMENT ID       | TECN      | COMMENT          |
|--------------------------|-------------------|-----------|------------------|
| $0.36 \pm 0.07 \pm 0.03$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_{||}, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{62}/\Gamma_{56}$$

| VALUE (%)                       | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $0.52 \pm 0.15^{+0.05}_{-0.02}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_{\perp}, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{63}/\Gamma_{56}$$

| VALUE (%)                       | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $0.63 \pm 0.34^{+0.16}_{-0.08}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_0(1500), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{65}/\Gamma_{56}$$

| VALUE                               | DOCUMENT ID       | TECN      | COMMENT          |
|-------------------------------------|-------------------|-----------|------------------|
| $0.101 \pm 0.008^{+0.011}_{-0.003}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f'_2(1525)_0, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{66}/\Gamma_{56}$$

| VALUE (%)                       | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $0.51 \pm 0.09^{+0.05}_{-0.04}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f'_2(1525)_{||}, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{67}/\Gamma_{56}$$

| VALUE (%)                       | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $0.06^{+0.13}_{-0.04} \pm 0.01$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f'_2(1525)_{\perp}, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{68}/\Gamma_{56}$$

| VALUE (%)                       | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $0.26 \pm 0.18^{+0.06}_{-0.04}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_0(1790), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{69}/\Gamma_{56}$$

| VALUE                               | DOCUMENT ID       | TECN      | COMMENT          |
|-------------------------------------|-------------------|-----------|------------------|
| $0.024 \pm 0.004^{+0.050}_{-0.002}$ | <sup>1</sup> AAIJ | 14BR LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\phi) \quad \Gamma_{59}/\Gamma_{49}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

**$0.119^{+0.013}_{-0.014}$  OUR FIT** Error includes scale factor of 2.4.

**$0.111^{+0.020}_{-0.018}$  OUR AVERAGE** Error includes scale factor of 2.5. See the ideogram below.

$0.069 \pm 0.012 \pm 0.001$  <sup>1</sup> KHACHATRYAN 16Q CMS  $pp$  at 7 TeV

$0.140^{+0.026}_{-0.013} \pm 0.002$  <sup>2,3</sup> AAIJ 12AO LHCb  $pp$  at 7 TeV

$0.135 \pm 0.036 \pm 0.001$  <sup>4</sup> ABAZOV 12C D0  $p\bar{p}$  at 1.96 TeV

$0.126 \pm 0.012 \pm 0.001$  <sup>5</sup> AALTONEN 11AB CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.124^{+0.026}_{-0.023} \pm 0.001$  <sup>6</sup> AAIJ 11 LHCb Repl. by AAIJ 12AO

<sup>1</sup> KHACHATRYAN 16Q reports  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.140 \pm 0.008 \pm 0.023$  which we multiply by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AAIJ 12AO reports  $(13.9 \pm 0.6^{+2.5}_{-1.2}) \times 10^{-2}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Measured in Dalitz plot like analysis of  $B_s \rightarrow J/\psi\pi^+\pi^-$  decays.


<sup>4</sup> ABAZOV 12C reports  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.275 \pm 0.041 \pm 0.061$  which we multiply by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> AALTONEN 11AB reports  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.257 \pm 0.020 \pm 0.014$  which we multiply by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup> AAIJ 11 reports  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.252^{+0.046+0.027}_{-0.032-0.033}$  which we multiply by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.



WEIGHTED AVERAGE  
0.111±0.020-0.018 (Error scaled by 2.5)



### $\Gamma(J/\psi(1S)f_0(1370), f_0 \rightarrow \pi^+\pi^-) / \Gamma_{\text{total}}$ $\Gamma_{64}/\Gamma$

| VALUE (units $10^{-4}$ )                                                      | DOCUMENT ID     | TECN | COMMENT                                |
|-------------------------------------------------------------------------------|-----------------|------|----------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |      |                                        |
| $0.34^{+0.11+0.085}_{-0.14-0.054}$                                            | <sup>1</sup> LI | 11   | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |

<sup>1</sup> The second error includes both the detector systematic and the uncertainty in the number of produced  $\Upsilon(5S) \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$  pairs.

### $\Gamma(J/\psi(1S)f_0(1370), f_0 \rightarrow \pi^+\pi^-) / \Gamma(J/\psi(1S)\phi)$ $\Gamma_{64}/\Gamma_{49}$

| VALUE (units $10^{-2}$ )        | DOCUMENT ID         | TECN | COMMENT            |
|---------------------------------|---------------------|------|--------------------|
| $4.22^{+0.55}_{-3.76} \pm 0.05$ | <sup>1,2</sup> AAIJ | 12AO | LHCB $pp$ at 7 TeV |

<sup>1</sup> AAIJ 12AO reports  $(4.19 \pm 0.53^{+0.12}_{-3.7}) \times 10^{-2}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(1370), f_0 \rightarrow \pi^+\pi^-) / \Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Measured in Dalitz plot like analysis of  $B_s \rightarrow J/\psi\pi^+\pi^-$  decays.

$$\Gamma(J/\psi(1S)f_2(1270), f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\phi) \quad \Gamma_{60}/\Gamma_{49}$$

| VALUE (units $10^{-4}$ )    | DOCUMENT ID | TECN      | COMMENT       |
|-----------------------------|-------------|-----------|---------------|
| $9.9^{+3.4}_{-3.6} \pm 0.1$ | 1,2 AAIJ    | 12AO LHCB | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 12AO reports  $(0.098 \pm 0.033^{+0.006}_{-0.015}) \times 10^{-2}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)f_2(1270), f_2 \rightarrow \pi^+\pi^-)/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Measured in Dalitz plot like analysis of  $B_S \rightarrow J/\psi\pi^+\pi^-$  decays for the  $f_2$  helicity state  $\lambda = 0$ .

$$\Gamma(J/\psi(1S)\pi^+\pi^-(\text{nonresonant}))/\Gamma(J/\psi(1S)\phi) \quad \Gamma_{70}/\Gamma_{49}$$

| VALUE (units $10^{-2}$ )        | DOCUMENT ID | TECN      | COMMENT       |
|---------------------------------|-------------|-----------|---------------|
| $1.67^{+1.02}_{-0.32} \pm 0.02$ | 1,2 AAIJ    | 12AO LHCB | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 12AO reports  $(1.66 \pm 0.31^{+0.96}_{-0.08}) \times 10^{-2}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)\pi^+\pi^-(\text{nonresonant}))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Measured in Dalitz plot like analysis of  $B_S \rightarrow J/\psi\pi^+\pi^-$  decays.

$$\Gamma(J/\psi(1S)\bar{K}^0\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma$$

| VALUE                 | CL% | DOCUMENT ID | TECN     | COMMENT       |
|-----------------------|-----|-------------|----------|---------------|
| $<4.4 \times 10^{-5}$ | 90  | 1 AAIJ      | 14L LHCB | $pp$ at 7 TeV |

<sup>1</sup> Measured with  $B(B_S^0 \rightarrow J/\psi K_S^0\pi^+\pi^-) / B(B^0 \rightarrow J/\psi K_S^0\pi^+\pi^-)$  using PDG 12 values for the involved branching fractions.

$$\Gamma(J/\psi(1S)K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{72}/\Gamma$$

| VALUE (units $10^{-4}$ )                    | DOCUMENT ID | TECN      | COMMENT                           |
|---------------------------------------------|-------------|-----------|-----------------------------------|
| <b><math>7.9 \pm 0.7</math> OUR AVERAGE</b> |             |           |                                   |
| $7.70 \pm 0.08 \pm 0.72$                    | 1 AAIJ      | 13AN LHCB | $pp$ at 7 TeV                     |
| $10.1 \pm 0.9 \pm 2.1$                      | 2 THORNE    | 13 BELL   | $e^+e^- \rightarrow \Upsilon(5S)$ |

<sup>1</sup> Uses  $f_s/f_d = 0.256 \pm 0.020$  and  $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$ .

<sup>2</sup> Uses  $f_s = (17.2 \pm 3.0)\%$  as the fraction of  $\Upsilon(5S)$  decaying to  $B_S^{(*)}\bar{B}_S^{(*)}$ .

$$\Gamma(J/\psi(1S)K^0K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{73}/\Gamma$$

| VALUE (units $10^{-4}$ )                | DOCUMENT ID | TECN     | COMMENT       |
|-----------------------------------------|-------------|----------|---------------|
| <b><math>9.2 \pm 1.0 \pm 0.9</math></b> | 1 AAIJ      | 14L LHCB | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 14L reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)K^0K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0\pi^+\pi^-)] = 2.12 \pm 0.15 \pm 0.18$  which we multiply by our best value  $B(B^0 \rightarrow J/\psi(1S)K^0\pi^+\pi^-) = (4.3 \pm 0.4) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. This is an observation of  $B_S^0 \rightarrow J/\psi K_S^0 K^\pm \pi^\mp$  with more than 10 standard deviations.

$\Gamma(J/\psi(1S)\bar{K}^0 K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$

| VALUE                           | CL% | DOCUMENT ID       | TECN     | COMMENT       |
|---------------------------------|-----|-------------------|----------|---------------|
| <b>&lt;12 × 10<sup>-6</sup></b> | 90  | <sup>1</sup> AAIJ | 14L LHCb | $pp$ at 7 TeV |

<sup>1</sup> Measured with  $B(B_S^0 \rightarrow J/\psi K_S^0 K^+ K^-)/B(B^0 \rightarrow J/\psi K_S^0 \pi^+ \pi^-)$  using PDG 12 values for the involved branching fractions.

$\Gamma(J/\psi(1S)f_2'(1525))/\Gamma(J/\psi(1S)\phi)$   $\Gamma_{75}/\Gamma_{49}$

| VALUE (units 10 <sup>-2</sup> )                                               | DOCUMENT ID           | TECN     | COMMENT                            |
|-------------------------------------------------------------------------------|-----------------------|----------|------------------------------------|
| <b>21 ± 4 OUR AVERAGE</b>                                                     |                       |          |                                    |
| 21.5 ± 4.9 ± 2.6                                                              | <sup>1</sup> THORNE   | 13 BELL  | $e^+ e^- \rightarrow \Upsilon(5S)$ |
| 21 ± 7 ± 1                                                                    | <sup>2,3</sup> ABAZOV | 12AF D0  | $p\bar{p}$ at 1.96 TeV             |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |          |                                    |
| 27 ± 4 ± 1                                                                    | <sup>4</sup> AAIJ     | 12S LHCb | Repl. by AAIJ 13AN                 |

<sup>1</sup> Uses  $B(f_2'(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1)\%$ .

<sup>2</sup> ABAZOV 12AF reports  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)f_2'(1525))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] \times B(f_2'(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-) = 0.19 \pm 0.05 \pm 0.04$  which we divide and multiply by our best values  $B(f_2'(1525) \rightarrow K^+ K^-) = \frac{1}{2} (87.6 \pm 2.2) \times 10^{-2}$ ,  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>3</sup> ABAZOV 12AF fits the invariant masses of the  $K^+ K^-$  pair in the range  $1.35 < M(K^+ K^-) < 2$  GeV.

<sup>4</sup> AAIJ 12S reports  $[(26.4 \pm 2.7 \pm 2.4) \times 10^{-2}]$  from a measurement of  $\Gamma(B_S^0 \rightarrow J/\psi(1S)f_2'(1525))/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi) \times B(f_2'(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-)$  assuming  $B(f_2'(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1) \times 10^{-2}$ ,  $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$ , which we rescale to our best values  $B(f_2'(1525) \rightarrow K^+ K^-) = \frac{1}{2} (87.6 \pm 2.2) \times 10^{-2}$ ,  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(J/\psi(1S)f_2'(1525))/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$

| VALUE (units 10 <sup>-4</sup> )                    | DOCUMENT ID       | TECN      | COMMENT       |
|----------------------------------------------------|-------------------|-----------|---------------|
| <b>2.61 ± 0.20<sup>+0.56</sup><sub>-0.50</sub></b> | <sup>1</sup> AAIJ | 13AN LHCb | $pp$ at 7 TeV |

<sup>1</sup> Uses  $f_s/f_d = 0.256 \pm 0.020$  and  $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$ .

$\Gamma(\psi(2S)\eta)/\Gamma(J/\psi(1S)\eta)$   $\Gamma_{80}/\Gamma_{52}$

| VALUE                     | DOCUMENT ID       | TECN      | COMMENT       |
|---------------------------|-------------------|-----------|---------------|
| <b>0.83 ± 0.14 ± 0.12</b> | <sup>1</sup> AAIJ | 13AA LHCb | $pp$ at 7 TeV |

<sup>1</sup> Assuming lepton universality for dimuon decay modes of  $J/\psi$  and  $\psi(2S)$  mesons, the ratio  $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$  was used.

$\Gamma(\psi(2S)\eta')/\Gamma(J/\psi(1S)\eta')$   $\Gamma_{81}/\Gamma_{55}$

| VALUE (units 10 <sup>-2</sup> ) | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------|-------------------|----------|------------------|
| <b>38.7 ± 9.0 ± 1.6</b>         | <sup>1</sup> AAIJ | 15D LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $J/\psi \rightarrow \mu^+ \mu^-$ ,  $\eta' \rightarrow \rho^0 \gamma$ , and  $\eta' \rightarrow \eta \pi^+ \pi^-$  decays.

$\Gamma(J/\psi(1S)p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$ 

| VALUE (units $10^{-6}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

|                                            |  |                   |          |                      |
|--------------------------------------------|--|-------------------|----------|----------------------|
| <b><math>3.58 \pm 0.19 \pm 0.39</math></b> |  | <sup>1</sup> AAIJ | 19U LHCb | $pp$ at 7, 8, 13 TeV |
|--------------------------------------------|--|-------------------|----------|----------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|      |    |                   |          |                     |
|------|----|-------------------|----------|---------------------|
| <4.8 | 90 | <sup>2</sup> AAIJ | 13Z LHCb | Repl. by AAIJ 2019U |
|------|----|-------------------|----------|---------------------|

<sup>1</sup> Measured relative to  $B_S^0 \rightarrow J/\psi\phi$  assuming  $B(B_S^0 \rightarrow J/\psi\phi) = (10.5 \pm 0.13 \pm 0.64) \times 10^{-4}$  and taking into account small  $K^+K^-$   $S$ -wave contribution.

<sup>2</sup> Uses  $B(B_S^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (1.98 \pm 0.20) \times 10^{-4}$ .

 $\Gamma(J/\psi(1S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$ 

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                                            |    |                   |           |                  |
|--------------------------------------------|----|-------------------|-----------|------------------|
| <b><math>&lt;7.3 \times 10^{-6}</math></b> | 90 | <sup>1</sup> AAIJ | 15BB LHCb | $pp$ at 7, 8 TeV |
|--------------------------------------------|----|-------------------|-----------|------------------|

<sup>1</sup> Branching fractions of normalization modes  $B_S^0 \rightarrow J/\psi\gamma X$  taken from PDG 14. Uses  $f_s/f_d = 0.259 \pm 0.015$ .

 $\Gamma(J/\psi(1S)\pi^+\pi^-\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_{78}/\Gamma_{56}$ 

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

|                                               |                   |          |                 |
|-----------------------------------------------|-------------------|----------|-----------------|
| <b><math>0.371 \pm 0.015 \pm 0.022</math></b> | <sup>1</sup> AAIJ | 14Y LHCb | $pp$ at 7,8 TeV |
|-----------------------------------------------|-------------------|----------|-----------------|

<sup>1</sup> Excludes contributions from  $\psi(2S)$  and  $\chi_{c1}(3872)$  decaying to  $J/\psi(1S)\pi^+\pi^-$ .

 $\Gamma(J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$ 

| VALUE (units $10^{-5}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

|                                         |                   |          |                  |
|-----------------------------------------|-------------------|----------|------------------|
| <b><math>7.2 \pm 1.3 \pm 0.4</math></b> | <sup>1</sup> AAIJ | 14Y LHCb | $pp$ at 7, 8 TeV |
|-----------------------------------------|-------------------|----------|------------------|

<sup>1</sup> AAIJ 14Y reports  $(7.14 \pm 0.99^{+0.83}_{-0.91} \pm 0.41) \times 10^{-5}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow 2\pi^+2\pi^-)]$  assuming  $B(f_1(1285) \rightarrow 2\pi^+2\pi^-) = 0.11^{+0.007}_{-0.006}$ , which we rescale to our best value  $B(f_1(1285) \rightarrow 2\pi^+2\pi^-) = (10.9 \pm 0.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\psi(2S)\phi)/\Gamma_{\text{total}}$   $\Gamma_{83}/\Gamma$ 

| VALUE (units $10^{-4}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|      |   |          |          |                        |
|------|---|----------|----------|------------------------|
| seen | 1 | BUSKULIC | 93G ALEP | $e^+e^- \rightarrow Z$ |
|------|---|----------|----------|------------------------|

 $\Gamma(\psi(2S)\phi)/\Gamma(J/\psi(1S)\phi)$   $\Gamma_{83}/\Gamma_{49}$ 

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

|                                                 |  |  |  |
|-------------------------------------------------|--|--|--|
| <b><math>0.503 \pm 0.034</math> OUR AVERAGE</b> |  |  |  |
|-------------------------------------------------|--|--|--|

|                             |                     |          |               |
|-----------------------------|---------------------|----------|---------------|
| $0.500 \pm 0.034 \pm 0.011$ | <sup>1,2</sup> AAIJ | 12L LHCb | $pp$ at 7 TeV |
|-----------------------------|---------------------|----------|---------------|

|                          |        |        |                        |
|--------------------------|--------|--------|------------------------|
| $0.53 \pm 0.10 \pm 0.09$ | ABAZOV | 09Y D0 | $p\bar{p}$ at 1.96 TeV |
|--------------------------|--------|--------|------------------------|

|                          |           |         |                        |
|--------------------------|-----------|---------|------------------------|
| $0.52 \pm 0.13 \pm 0.07$ | ABULENCIA | 06N CDF | $p\bar{p}$ at 1.96 TeV |
|--------------------------|-----------|---------|------------------------|

<sup>1</sup> AAIJ 12L reports  $0.489 \pm 0.026 \pm 0.021 \pm 0.012$  from a measurement of  $[\Gamma(B_S^0 \rightarrow \psi(2S)\phi)/\Gamma(B_S^0 \rightarrow J/\psi(1S)\phi)] \times [B(J/\psi(1S) \rightarrow e^+e^-)] / [B(\psi(2S) \rightarrow e^+e^-)]$  assuming  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow e^+e^-) = (7.72 \pm 0.17) \times 10^{-3}$ , which we rescale to our best values  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow e^+e^-) = (7.93 \pm 0.17) \times 10^{-3}$ . Our first error

is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> Assumes  $B(J/\psi \rightarrow \mu^+ \mu^-) / B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-) / B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$ .

### $\Gamma(\psi(2S)K^-\pi^+)/\Gamma_{\text{total}}$

$\Gamma_{84}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID       | TECN     | COMMENT          |
|--------------------------------------------|-------------------|----------|------------------|
| <b><math>3.12 \pm 0.30 \pm 0.21</math></b> | <sup>1</sup> AAIJ | 15U LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 15U reports  $[\Gamma(B_S^0 \rightarrow \psi(2S)K^-\pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \psi(2S)K^+\pi^-)] = (5.38 \pm 0.36 \pm 0.22 \pm 0.31) \times 10^{-2}$  which we multiply by our best value  $B(B^0 \rightarrow \psi(2S)K^+\pi^-) = (5.8 \pm 0.4) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\psi(2S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}$

$\Gamma_{85}/\Gamma$

| VALUE (units $10^{-5}$ )                      | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------------------------------|-------------------|----------|------------------|
| <b><math>3.3 \pm 0.5^{+0.2}_{-0.3}</math></b> | <sup>1</sup> AAIJ | 15U LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 15U reports  $[\Gamma(B_S^0 \rightarrow \psi(2S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \psi(2S)K^*(892)^0)] = (5.58 \pm 0.57 \pm 0.40 \pm 0.32) \times 10^{-2}$  which we multiply by our best value  $B(B^0 \rightarrow \psi(2S)K^*(892)^0) = (5.9 \pm 0.4) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\chi_{c1}\phi)/\Gamma(J/\psi(1S)\phi)$

$\Gamma_{86}/\Gamma_{49}$

| VALUE (units $10^{-2}$ )                 | DOCUMENT ID       | TECN      | COMMENT       |
|------------------------------------------|-------------------|-----------|---------------|
| <b><math>18.9 \pm 1.8 \pm 1.5</math></b> | <sup>1</sup> AAIJ | 13AC LHCb | $pp$ at 7 TeV |

<sup>1</sup> Uses  $B(\chi_{c1} \rightarrow J/\psi\gamma) = (34.4 \pm 1.5)\%$ .

### $\Gamma(\chi_{c2}K^+K^-)/\Gamma(\chi_{c1}K^+K^-)$

$\Gamma_{88}/\Gamma_{87}$

| VALUE (units $10^{-2}$ )                 | DOCUMENT ID       | TECN      | COMMENT              |
|------------------------------------------|-------------------|-----------|----------------------|
| <b><math>17.1 \pm 3.1 \pm 1.0</math></b> | <sup>1</sup> AAIJ | 18AC LHCb | $pp$ at 7, 8, 13 TeV |

<sup>1</sup> Measures the ratio for  $\pm 15$  MeV window around  $\phi$  mass.

### $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{82}/\Gamma_{56}$

| VALUE                                      | DOCUMENT ID       | TECN      | COMMENT       |
|--------------------------------------------|-------------------|-----------|---------------|
| <b><math>0.34 \pm 0.04 \pm 0.03</math></b> | <sup>1</sup> AAIJ | 13AA LHCb | $pp$ at 7 TeV |

<sup>1</sup> Assuming lepton universality for dimuon decay modes of  $J/\psi$  and  $\psi(2S)$  mesons, the ratio  $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$  was used.

### $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{89}/\Gamma$

| VALUE (units $10^{-7}$ )                    | CL% | DOCUMENT ID           | TECN     | COMMENT                |
|---------------------------------------------|-----|-----------------------|----------|------------------------|
| <b><math>7.0 \pm 1.0</math> OUR AVERAGE</b> |     |                       |          |                        |
| $7.3 \pm 0.9 \pm 0.7$                       |     | <sup>1</sup> AAIJ     | 17G LHCb | $pp$ at 7 and 8 TeV    |
| $6.4 \pm 1.8 \pm 0.6$                       |     | <sup>2</sup> AALTONEN | 12L CDF  | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                              |    |                          |           |                                 |
|------------------------------|----|--------------------------|-----------|---------------------------------|
| $10.4^{+2.4}_{-2.1} \pm 1.0$ |    | <sup>3</sup> AAIJ        | 12AR LHCb | Repl. by AAIJ 17G               |
| < 120                        | 90 | <sup>4</sup> PENG        | 10 BELL   | $e^+e^- \rightarrow \gamma(5S)$ |
| < 12                         | 90 | <sup>5</sup> AALTONEN    | 09C CDF   | Repl. by AALTONEN 12L           |
| < 17                         | 90 | <sup>6</sup> ABULENCIA,A | 06D CDF   | Repl. by AALTONEN 09C           |
| <2320                        | 90 | <sup>7</sup> ABE         | 00C SLD   | $e^+e^- \rightarrow Z$          |
| <1700                        | 90 | <sup>8</sup> BUSKULIC    | 96V ALEP  | $e^+e^- \rightarrow Z$          |

<sup>1</sup> AAIJ 17G reports  $[\Gamma(B_S^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0)] = (9.15 \pm 0.71 \pm 0.83) \times 10^{-3}$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ ,  $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> AALTONEN 12L reports  $[\Gamma(B_S^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.008 \pm 0.002 \pm 0.001$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ ,  $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>3</sup> AAIJ 12AR reports  $[\Gamma(B_S^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \pi^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.050^{+0.011}_{-0.009} \pm 0.004$  which we multiply or divide by our best values  $B(B^0 \rightarrow \pi^+\pi^-) = (5.12 \pm 0.19) \times 10^{-6}$ ,  $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>4</sup> Uses  $\gamma(10860) \rightarrow B_S^* \bar{B}_S^*$  and assumes  $B(\gamma(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\gamma(10860) \rightarrow B_S^* \bar{B}_S^*) / \Gamma(\gamma(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

<sup>5</sup> Obtains this result from  $(f_s/f_d) \cdot B(B_S \rightarrow \pi^+\pi^-)/B(B^0 \rightarrow K^+\pi^-) = 0.007 \pm 0.004 \pm 0.005$ , assuming  $f_s/f_d = 0.276 \pm 0.034$  and  $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$ .

<sup>6</sup> ABULENCIA,A 06D obtains this from  $B(B_S \rightarrow \pi^+\pi^-) / B(B_S \rightarrow K^+K^-) < 0.05$  at 90% CL, assuming  $B(B_S \rightarrow K^+K^-) = (33 \pm 6 \pm 7) \times 10^{-6}$ .

<sup>7</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_S} = (10.5^{+1.8}_{-2.2})\%$ .

<sup>8</sup> BUSKULIC 96V assumes PDG 96 production fractions for  $B^0$ ,  $B^+$ ,  $B_S$ ,  $b$  baryons.

| $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ |     |                       |        |                        | $\Gamma_{90}/\Gamma$ |
|--------------------------------------------|-----|-----------------------|--------|------------------------|----------------------|
| VALUE                                      | CL% | DOCUMENT ID           | TECN   | COMMENT                |                      |
| $<2.1 \times 10^{-4}$                      | 90  | <sup>1</sup> ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |                      |

<sup>1</sup> ACCIARRI 95H assumes  $f_{B^0} = 39.5 \pm 4.0$  and  $f_{B_S} = 12.0 \pm 3.0\%$ .

| $\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$ |     |                       |        |                        | $\Gamma_{91}/\Gamma$ |
|-------------------------------------------|-----|-----------------------|--------|------------------------|----------------------|
| VALUE                                     | CL% | DOCUMENT ID           | TECN   | COMMENT                |                      |
| $<1.0 \times 10^{-3}$                     | 90  | <sup>1</sup> ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |                      |

<sup>1</sup> ACCIARRI 95H assumes  $f_{B^0} = 39.5 \pm 4.0$  and  $f_{B_S} = 12.0 \pm 3.0\%$ .

### $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ $\Gamma_{92}/\Gamma$

| VALUE                                      | CL% | DOCUMENT ID           | TECN   | COMMENT                |
|--------------------------------------------|-----|-----------------------|--------|------------------------|
| <b><math>&lt;1.5 \times 10^{-3}</math></b> | 90  | <sup>1</sup> ACCIARRI | 95H L3 | $e^+e^- \rightarrow Z$ |

<sup>1</sup> ACCIARRI 95H assumes  $f_{B^0} = 39.5 \pm 4.0$  and  $f_{B_s} = 12.0 \pm 3.0\%$ .

### $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$ $\Gamma_{93}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID      | TECN    | COMMENT                |
|---------------------------------------------|-----|------------------|---------|------------------------|
| <b><math>&lt;3.20 \times 10^{-4}</math></b> | 90  | <sup>1</sup> ABE | 00C SLD | $e^+e^- \rightarrow Z$ |

<sup>1</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

### $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ $\Gamma_{94}/\Gamma$

| VALUE (units $10^{-5}$ )                | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------------------------|-------------------|----------|------------------|
| <b><math>3.3 \pm 0.7 \pm 0.1</math></b> | <sup>1</sup> AAIJ | 150 LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 150 reports  $[\Gamma(B_S^0 \rightarrow \eta'\eta')/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \eta'K^+)] = 0.47 \pm 0.09 \pm 0.04$  which we multiply by our best value  $B(B^+ \rightarrow \eta'K^+) = (7.04 \pm 0.25) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\eta'\phi)/\Gamma_{\text{total}}$ $\Gamma_{95}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt;0.82 \times 10^{-6}</math></b> | 90  | <sup>1</sup> AAIJ | 17BA LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Corresponds to the 95% CL upper limit  $1.01 \times 10^{-6}$ . Uses the normalization mode  $B^+ \rightarrow \eta'K^+$  with branching fraction  $(70.6 \pm 2.5) \times 10^{-6}$  and the ratio of hadronisation fractions  $f_s/f_d = 0.259 \pm 0.015$ , which is assumed equal to  $f_s/f_u$ .

### $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{99}/\Gamma$

| VALUE (units $10^{-6}$ )                   | DOCUMENT ID       | TECN     | COMMENT          |
|--------------------------------------------|-------------------|----------|------------------|
| <b><math>3.48 \pm 0.23 \pm 0.39</math></b> | <sup>1</sup> AAIJ | 17A LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Inclusive decays in mass range  $400 < m(\pi^+\pi^-) < 1600 \text{ MeV}/c^2$ .

### $\Gamma(\phi\rho^0)/\Gamma_{\text{total}}$ $\Gamma_{98}/\Gamma$

| VALUE (units $10^{-7}$ )                | CL% | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------------------------|-----|-------------------|----------|------------------|
| <b><math>2.7 \pm 0.7 \pm 0.3</math></b> |     | <sup>1</sup> AAIJ | 17A LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|         |    |                  |         |                        |
|---------|----|------------------|---------|------------------------|
| $<6170$ | 90 | <sup>2</sup> ABE | 00C SLD | $e^+e^- \rightarrow Z$ |
|---------|----|------------------|---------|------------------------|

<sup>1</sup> Signal evidence is 4 standard deviations.  
<sup>2</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

### $\Gamma(\phi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{96}/\Gamma$

| VALUE (units $10^{-6}$ )                   | DOCUMENT ID       | TECN     | COMMENT          |
|--------------------------------------------|-------------------|----------|------------------|
| <b><math>1.12 \pm 0.16 \pm 0.14</math></b> | <sup>1</sup> AAIJ | 17A LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Signal is observed with 8 standard deviations significance.

$\Gamma(\phi f_2(1270), f_2(1270) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ 
 $\Gamma_{97} / \Gamma$ 

| VALUE (units $10^{-6}$ )                          | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------------------------|-------------------|----------|------------------|
| <b><math>0.61 \pm 0.13^{+0.13}_{-0.08}</math></b> | <sup>1</sup> AAIJ | 17A LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Signal is observed with 5 standard deviations significance.

 $\Gamma(\phi\phi) / \Gamma_{\text{total}}$ 
 $\Gamma_{100} / \Gamma$ 

| VALUE (units $10^{-6}$ )                 | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>18.7 \pm 1.5</math> OUR FIT</b> |     |                   |           |                  |
| <b><math>18.5 \pm 1.4 \pm 1.0</math></b> |     | <sup>1</sup> AAIJ | 15AS LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                                           |                  |         |                         |
|-----------------------------------------------------------|------------------|---------|-------------------------|
| 14 $\begin{smallmatrix} +6 \\ -5 \end{smallmatrix} \pm 6$ | 2 ACOSTA         | 05J CDF | Repl. by AALTONEN 11AN  |
| <1183 90                                                  | <sup>3</sup> ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |

<sup>1</sup> AAIJ 15AS reports  $[\Gamma(B_s^0 \rightarrow \phi\phi) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \phi)] = 1.84 \pm 0.05 \pm 0.13$  which we multiply by our best value  $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Uses  $B(B^0 \rightarrow J/\psi\phi) = (1.38 \pm 0.49) \times 10^{-3}$  and production cross-section ratio of  $\sigma(B_s) / \sigma(B^0) = 0.26 \pm 0.04$ .

<sup>3</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

 $\Gamma(\phi\phi) / \Gamma(J/\psi(1S)\phi)$ 
 $\Gamma_{100} / \Gamma_{49}$ 

| VALUE (units $10^{-2}$ )                   | DOCUMENT ID | TECN     | COMMENT                |
|--------------------------------------------|-------------|----------|------------------------|
| <b><math>1.73 \pm 0.16</math> OUR FIT</b>  |             |          |                        |
| <b><math>1.78 \pm 0.14 \pm 0.20</math></b> | AALTONEN    | 11AN CDF | $p\bar{p}$ at 1.96 TeV |

 $\Gamma(\phi\phi\phi) / \Gamma(\phi\phi)$ 
 $\Gamma_{101} / \Gamma_{100}$ 

| VALUE                                         | DOCUMENT ID | TECN      | COMMENT          |
|-----------------------------------------------|-------------|-----------|------------------|
| <b><math>0.117 \pm 0.030 \pm 0.015</math></b> | AAIJ        | 17BB LHCb | $pp$ at 7, 8 TeV |

 $\Gamma(\pi^+ K^-) / \Gamma_{\text{total}}$ 
 $\Gamma_{102} / \Gamma$ 

| VALUE (units $10^{-6}$ )                    | CL% | DOCUMENT ID           | TECN      | COMMENT                |
|---------------------------------------------|-----|-----------------------|-----------|------------------------|
| <b><math>5.8 \pm 0.7</math> OUR AVERAGE</b> |     |                       |           |                        |
| $5.9 \pm 0.7 \pm 0.6$                       |     | <sup>1</sup> AAIJ     | 12AR LHCb | $pp$ at 7 TeV          |
| $5.7 \pm 1.0 \pm 0.5$                       |     | <sup>2</sup> AALTONEN | 09C CDF   | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|          |                          |          |                                    |
|----------|--------------------------|----------|------------------------------------|
| < 26 90  | <sup>3</sup> PENG        | 10 BELL  | $e^+ e^- \rightarrow \Upsilon(5S)$ |
| < 5.6 90 | <sup>4</sup> ABULENCIA,A | 06D CDF  | Repl. by AALTONEN 09C              |
| <261 90  | <sup>5</sup> ABE         | 00C SLD  | $e^+ e^- \rightarrow Z$            |
| <210 90  | <sup>6</sup> BUSKULIC    | 96V ALEP | $e^+ e^- \rightarrow Z$            |
| <260 90  | <sup>7</sup> AKERS       | 94L OPAL | $e^+ e^- \rightarrow Z$            |

<sup>1</sup> AAIJ 12AR reports  $[\Gamma(B_s^0 \rightarrow \pi^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0) / \Gamma(\bar{b} \rightarrow B^0)] = 0.074 \pm 0.006 \pm 0.006$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ ,  $\Gamma(\bar{b} \rightarrow B_s^0) / \Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.



<sup>2</sup> AALTONEN 09C reports  $[\Gamma(B_S^0 \rightarrow \pi^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [B(\bar{b} \rightarrow B_S^0)] / [B(\bar{b} \rightarrow B^0)] = 0.071 \pm 0.010 \pm 0.007$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ ,  $B(\bar{b} \rightarrow B_S^0) = (10.0 \pm 0.8) \times 10^{-2}$ ,  $B(\bar{b} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>3</sup> Uses  $\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*$  and assumes  $B(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*) / \Gamma(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

<sup>4</sup> ABULENCIA,A 06D obtains this from  $(f_s/f_d) (B(B_S \rightarrow \pi^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) < 0.08$  at 90% CL, assuming  $f_s/f_d = 0.260 \pm 0.039$  and  $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$ .

<sup>5</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_S} = (10.5^{+1.8}_{-2.2})\%$ .

<sup>6</sup> BUSKULIC 96V assumes PDG 96 production fractions for  $B^0$ ,  $B^+$ ,  $B_S$ ,  $b$  baryons.

<sup>7</sup> Assumes  $B(Z \rightarrow b\bar{b}) = 0.217$  and  $B_d^0$  ( $B_S^0$ ) fraction 39.5% (12%).

## $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$

$\Gamma_{103}/\Gamma$

| VALUE (units $10^{-6}$ )      | CL% | DOCUMENT ID           | TECN      | COMMENT                            |
|-------------------------------|-----|-----------------------|-----------|------------------------------------|
| <b>26.6 ± 2.2 OUR AVERAGE</b> |     |                       |           |                                    |
| 25.2 ± 1.7 ± 2.4              |     | <sup>1</sup> AAIJ     | 12AR LHCb | $p\bar{p}$ at 7 TeV                |
| 27.6 ± 2.3 ± 2.7              |     | <sup>2</sup> AALTONEN | 11N CDF   | $p\bar{p}$ at 1.96 TeV             |
| 38 $^{+10}_{-9}$ ± 7          |     | <sup>3</sup> PENG     | 10 BELL   | $e^+ e^- \rightarrow \Upsilon(5S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|            |    |                              |          |                                    |
|------------|----|------------------------------|----------|------------------------------------|
| <310       | 90 | DRUTSKOY                     | 07A BELL | $e^+ e^- \rightarrow \Upsilon(5S)$ |
| 33 ± 6 ± 7 |    | <sup>4</sup> ABULENCIA,A 06D | CDF      | Repl. by AALTONEN 11N              |
| <283       | 90 | <sup>5</sup> ABE             | 00C SLD  | $e^+ e^- \rightarrow Z$            |
| < 59       | 90 | <sup>6</sup> BUSKULIC        | 96V ALEP | $e^+ e^- \rightarrow Z$            |
| <140       | 90 | <sup>7</sup> AKERS           | 94L OPAL | $e^+ e^- \rightarrow Z$            |

<sup>1</sup> AAIJ 12AR reports  $[\Gamma(B_S^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_S^0)] / [\Gamma(\bar{b} \rightarrow B^0)] = 0.316 \pm 0.009 \pm 0.019$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ ,  $\Gamma(\bar{b} \rightarrow B_S^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> AALTONEN 11N reports  $(f_s/f_d) (B(B_S^0 \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.347 \pm 0.020 \pm 0.021$ . We multiply this result by our best value of  $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$  and divide by our best value of  $f_s/f_d$ , where  $1/2 f_s/f_d = 0.1230 \pm 0.0115$ . Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

<sup>3</sup> Uses  $\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*$  and assumes  $B(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_S^* \bar{B}_S^*) / \Gamma(\Upsilon(10860) \rightarrow B_S^{(*)} \bar{B}_S^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

<sup>4</sup> ABULENCIA,A 06D obtains this from  $(f_s/f_d) (B(B_S \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.46 \pm 0.08 \pm 0.07$ , assuming  $f_s/f_d = 0.260 \pm 0.039$  and  $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$ .

<sup>5</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_S} = (10.5^{+1.8}_{-2.2})\%$ .

<sup>6</sup> BUSKULIC 96V assumes PDG 96 production fractions for  $B^0$ ,  $B^+$ ,  $B_S$ ,  $b$  baryons.

<sup>7</sup> Assumes  $B(Z \rightarrow b\bar{b}) = 0.217$  and  $B_d^0$  ( $B_S^0$ ) fraction 39.5% (12%).

# $\Gamma(K^0 \bar{K}^0)/\Gamma_{\text{total}}$ $\Gamma_{104}/\Gamma$

| VALUE (units $10^{-5}$ )                 | CL% | DOCUMENT ID      | TECN | COMMENT                                |
|------------------------------------------|-----|------------------|------|----------------------------------------|
| $1.96^{+0.58}_{-0.51} \pm 0.10 \pm 0.20$ |     | <sup>1</sup> PAL | 16   | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.6 90 <sup>2</sup> PENG 10 BELL Repl. by PAL 16

<sup>1</sup> Observed in  $B_s^0 \rightarrow K_S^0 K_S^0$  with significance of  $5.1 \sigma$ . The last uncertainty is due to the uncertainty of the total number of  $B_s^0 \bar{B}_s^0$  pairs.

<sup>2</sup> Uses  $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$  and assumes  $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$  and  $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$ .

# $\Gamma(K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ $\Gamma_{105}/\Gamma$

| VALUE (units $10^{-6}$ ) | DOCUMENT ID         | TECN      | COMMENT          |
|--------------------------|---------------------|-----------|------------------|
| $9.5 \pm 2.1 \pm 0.3$    | <sup>1,2</sup> AAIJ | 17BP LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

14  $\pm 4 \pm 1$  <sup>3</sup> AAIJ 13BP LHCB Repl. by AAIJ 17BP

<sup>1</sup> AAIJ 17BP reports  $[\Gamma(B_s^0 \rightarrow K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.191 \pm 0.027 \pm 0.033$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.97 \pm 0.18) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Used  $f_s/f_d = 0.259 \pm 0.015$ .

<sup>3</sup> AAIJ 13BP reports  $[\Gamma(B_s^0 \rightarrow K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.29 \pm 0.06 \pm 0.04$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.97 \pm 0.18) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

# $\Gamma(K^*(892)^- \pi^+)/\Gamma_{\text{total}}$ $\Gamma_{107}/\Gamma$

| VALUE (units $10^{-6}$ ) | DOCUMENT ID         | TECN     | COMMENT       |
|--------------------------|---------------------|----------|---------------|
| $2.9 \pm 1.0 \pm 0.2$    | <sup>1,2</sup> AAIJ | 14BMLHCB | $pp$ at 7 TeV |

<sup>1</sup> AAIJ 14BM reports  $[\Gamma(B_s^0 \rightarrow K^*(892)^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+ \pi^-)] = 0.39 \pm 0.13 \pm 0.05$  which we multiply by our best value  $B(B^0 \rightarrow K^*(892)^+ \pi^-) = (7.5 \pm 0.4) \times 10^{-6}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Uses  $f_s/f_d = 0.259 \pm 0.015$ .

# $\Gamma(K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ $\Gamma_{106}/\Gamma$

| VALUE (units $10^{-5}$ ) | DOCUMENT ID         | TECN      | COMMENT          |
|--------------------------|---------------------|-----------|------------------|
| $8.4 \pm 0.8 \pm 0.3$    | <sup>1,2</sup> AAIJ | 17BP LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.4  $\pm 0.9 \pm 0.3$  <sup>3</sup> AAIJ 13BP LHCB Repl. by AAIJ 17BP

<sup>1</sup> AAIJ 17BP reports  $[\Gamma(B_s^0 \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 1.70 \pm 0.07 \pm 0.15$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.97 \pm 0.18) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Used  $f_s/f_d = 0.259 \pm 0.015$ .

<sup>3</sup> AAIJ 13BP reports  $[\Gamma(B_S^0 \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 1.48 \pm 0.12 \pm 0.14$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.97 \pm 0.18) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$ $\Gamma_{108}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID | TECN     | COMMENT           |
|--------------------------------------------|-------------|----------|-------------------|
| <b><math>1.86 \pm 0.12 \pm 0.45</math></b> | 1,2 AAIJ    | 19K LHCB | $pp$ at 7, 8 TeV  |
| $1.12 \pm 0.21^{+0.07}_{-0.06}$            | 3,4 AAIJ    | 14BMLHCB | Repl. by AAIJ 19K |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 19K reports  $(18.6 \pm 1.2 \pm 0.8 \pm 4.0 \pm 2.0) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

<sup>3</sup> AAIJ 14BM reports  $[\Gamma(B_S^0 \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+ \pi^-)] = 1.49 \pm 0.22 \pm 0.18$  which we multiply by our best value  $B(B^0 \rightarrow K^*(892)^+ \pi^-) = (7.5 \pm 0.4) \times 10^{-6}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Uses  $f_s/f_d = 0.259 \pm 0.015$ .

### $\Gamma(K_0^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ $\Gamma_{109}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID | TECN     | COMMENT          |
|--------------------------------------------|-------------|----------|------------------|
| <b><math>3.13 \pm 0.23 \pm 2.53</math></b> | 1,2 AAIJ    | 19K LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 19K reports  $(31.3 \pm 2.3 \pm 0.7 \pm 25.1 \pm 3.3) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

### $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ $\Gamma_{110}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID | TECN     | COMMENT          |
|--------------------------------------------|-------------|----------|------------------|
| <b><math>1.03 \pm 0.25 \pm 1.64</math></b> | 1,2 AAIJ    | 19K LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 19K reports  $(10.3 \pm 2.5 \pm 1.1 \pm 16.3 \pm 1.1) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

### $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_{111}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID | TECN     | COMMENT          |
|--------------------------------------------|-------------|----------|------------------|
| <b><math>1.98 \pm 0.28 \pm 0.50</math></b> | 1,2 AAIJ    | 19K LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 19K reports  $(19.8 \pm 2.8 \pm 1.2 \pm 4.4 \pm 2.1) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

### $\Gamma(K_0^*(1430) \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_{112}/\Gamma$

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID | TECN     | COMMENT          |
|--------------------------------------------|-------------|----------|------------------|
| <b><math>3.30 \pm 0.25 \pm 0.98</math></b> | 1,2 AAIJ    | 19K LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 19K reports  $(33.0 \pm 2.5 \pm 0.9 \pm 9.1 \pm 3.5) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{113}/\Gamma$ 

| VALUE (units $10^{-5}$ )                   | DOCUMENT ID         | TECN     | COMMENT          |
|--------------------------------------------|---------------------|----------|------------------|
| <b><math>1.68 \pm 0.45 \pm 2.13</math></b> | <sup>1,2</sup> AAIJ | 19K LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 19K reports  $(16.8 \pm 4.5 \pm 1.7 \pm 21.2 \pm 1.8) \times 10^{-6}$  as the measured value. We have combined in quadrature all systematic uncertainties into a single one.

<sup>2</sup> Measured in Dalitz plot analysis of  $B_S^0 \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

 $\Gamma(K_S^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$ 

| VALUE (units $10^{-6}$ )                 | DOCUMENT ID       | TECN    | COMMENT       |
|------------------------------------------|-------------------|---------|---------------|
| <b><math>16.4 \pm 3.4 \pm 2.3</math></b> | <sup>1</sup> AAIJ | 16 LHCb | $pp$ at 7 TeV |

<sup>1</sup> Measured relative to  $B^0 \rightarrow K_S^0 \pi^+ \pi^-$  using the value of  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.2) \times 10^{-5}$ .

 $\Gamma(K^0 K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{115}/\Gamma$ 

| VALUE (units $10^{-7}$ )                 | CL%   | DOCUMENT ID | TECN      | COMMENT          |
|------------------------------------------|-------|-------------|-----------|------------------|
| <b><math>12.9 \pm 6.5 \pm 0.5</math></b> | 1,2,3 | AAIJ        | 17BP LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<34 90 <sup>4</sup> AAIJ 13BP LHCb Repl. by AAIJ 17BP

<sup>1</sup> AAIJ 17BP reports  $[\Gamma(B_S^0 \rightarrow K^0 K^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.026 \pm 0.011 \pm 0.007$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.97 \pm 0.18) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AAIJ 17BP also set the limit range  $4-25 \times 10^{-7}$  at 90% CL using the world average value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ .

<sup>3</sup> Used  $f_s/f_d = 0.259 \pm 0.015$ .

<sup>4</sup> AAIJ 13BP reports  $[\Gamma(B_S^0 \rightarrow K^0 K^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] < 0.068$  which we multiply by our best value  $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = 4.97 \times 10^{-5}$ .

 $\Gamma(\bar{K}^*(892)^0 \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$ 

| VALUE                                       | CL% | DOCUMENT ID      | TECN    | COMMENT                 |
|---------------------------------------------|-----|------------------|---------|-------------------------|
| <b><math>&lt;7.67 \times 10^{-4}</math></b> | 90  | <sup>1</sup> ABE | 00C SLD | $e^+ e^- \rightarrow Z$ |

<sup>1</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

 $\Gamma(\bar{K}^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{117}/\Gamma$ 

| VALUE (units $10^{-5}$ )                   | CL% | DOCUMENT ID       | TECN      | COMMENT       |
|--------------------------------------------|-----|-------------------|-----------|---------------|
| <b><math>1.11 \pm 0.26 \pm 0.06</math></b> |     | <sup>1</sup> AAIJ | 15AF LHCb | $pp$ at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.81 \pm 0.46 \pm 0.56$  <sup>2</sup> AAIJ 12F LHCb Repl. by AAIJ 15AF

<168.1 90 <sup>3</sup> ABE 00C SLD  $e^+ e^- \rightarrow Z$

<sup>1</sup> AAIJ 15AF reports  $[\Gamma(B_S^0 \rightarrow \bar{K}^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \phi)] = 1.11 \pm 0.22 \pm 0.12 \pm 0.06$  which we multiply by our best value  $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

- <sup>2</sup> Uses  $B^0 \rightarrow J/\psi K^{*0}$  for normalization and assumes  $B(B^0 \rightarrow J/\psi K^{*0}) B(J/\psi \rightarrow \mu^+ \mu^-) B(K^{*0} \rightarrow K^+ \pi^-) = (1.33 \pm 0.06) \times 10^{-3}$  and  $f_s/f_d = 0.253 \pm 0.031$ . The second quoted error is total uncertainty including the error of 0.34 on  $f_s/f_d$ .
- <sup>3</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

## $\Gamma(\phi K^*(892)^0)/\Gamma_{\text{total}}$ $\Gamma_{121}/\Gamma$

| VALUE (units $10^{-6}$ )                   | CL% | DOCUMENT ID       | TECN      | COMMENT       |
|--------------------------------------------|-----|-------------------|-----------|---------------|
| <b><math>1.14 \pm 0.29 \pm 0.06</math></b> |     | <sup>1</sup> AAIJ | 13BW LHCB | $pp$ at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

- <1013      90      <sup>2</sup> ABE      00C      SLD       $e^+e^- \rightarrow Z$
- <sup>1</sup> AAIJ 13BW reports  $[\Gamma(B_s^0 \rightarrow \phi K^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \phi)] = 0.113 \pm 0.024 \pm 0.016$  which we multiply by our best value  $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>2</sup> ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

## $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ $\Gamma_{122}/\Gamma$

Test for  $\Delta B=1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE (units $10^{-8}$ ) | CL% | DOCUMENT ID       | TECN      | COMMENT             |
|--------------------------|-----|-------------------|-----------|---------------------|
| <b>&lt; 1.5</b>          | 90  | <sup>1</sup> AAIJ | 17BJ LHCB | $pp$ at 7 and 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

- $2.84^{+2.03+0.85}_{-1.68-0.18}$       <sup>2</sup> AAIJ      13BQ LHCB      Repl. by AAIJ 17BJ
- <5900      90      <sup>3</sup> BUSKULIC      96V ALEP       $e^+e^- \rightarrow Z$
- <sup>1</sup> Uses normalization mode  $B(B^0 \rightarrow K^+ \pi^-) = (19.6 \pm 0.5) \times 10^{-6}$  and  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$ .
- <sup>2</sup> Uses normalization mode  $B(B^0 \rightarrow K^+ \pi^-) = (19.55 \pm 0.54) \times 10^{-6}$  and  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ .
- <sup>3</sup> BUSKULIC 96V assumes PDG 96 production fractions for  $B^0$ ,  $B^+$ ,  $B_s$ ,  $b$  baryons.

## $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$ $\Gamma_{123}/\Gamma$

| VALUE (units $10^{-6}$ )                | DOCUMENT ID         | TECN      | COMMENT          |
|-----------------------------------------|---------------------|-----------|------------------|
| <b><math>4.5 \pm 0.4 \pm 0.2</math></b> | <sup>1,2</sup> AAIJ | 17BD LHCB | $pp$ at 7, 8 TeV |

- <sup>1</sup> AAIJ 17BD reports  $[\Gamma(B_s^0 \rightarrow p\bar{p}K^+K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [B(J/\psi(1S) \rightarrow p\bar{p})] / [B(K^*(892) \rightarrow (K\pi)^\pm)] = 1.67 \pm 0.12 \pm 0.11$  which we multiply by our best values  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$ ,  $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$ ,  $B(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes  $f_s/f_d = 0.259 \pm 0.015$ .
- <sup>2</sup> The branching ratio is given for  $m_{p\bar{p}} < 2.85$  GeV.

$$\Gamma(p\bar{p}K^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{124}/\Gamma$$

| VALUE (units $10^{-7}$ )                 | DOCUMENT ID | TECN      | COMMENT          |
|------------------------------------------|-------------|-----------|------------------|
| <b><math>13.9 \pm 2.5 \pm 0.5</math></b> | 1,2 AAIJ    | 17BD LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17BD reports  $[\Gamma(B_s^0 \rightarrow p\bar{p}K^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [B(J/\psi(1S) \rightarrow p\bar{p})] / [B(K^*(892) \rightarrow (K\pi)^\pm)] = 0.52 \pm 0.08 \pm 0.05$  which we multiply by our best values  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$ ,  $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$ ,  $B(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes  $f_s/f_d = 0.259 \pm 0.015$ .

<sup>2</sup> The branching ratio is given for  $m_{p\bar{p}} < 2.85$  GeV.

$$\Gamma(p\bar{p}K^+\pi^-)/\Gamma(p\bar{p}K^+K^-) \quad \Gamma_{124}/\Gamma_{123}$$

| VALUE                                      | DOCUMENT ID | TECN      | COMMENT          |
|--------------------------------------------|-------------|-----------|------------------|
| <b><math>0.31 \pm 0.05 \pm 0.02</math></b> | 1,2 AAIJ    | 17BD LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Reports  $B(B_s^0 \rightarrow p\bar{p}K^+\pi^-) / B(B^0 \rightarrow p\bar{p}K^+\pi^-) = 0.22 \pm 0.04 \pm 0.02 \pm 0.01$ , where the third error is due to  $f_s/f_d$ .

<sup>2</sup> The ratio is given for  $m_{p\bar{p}} < 2.85$  GeV and assuming  $f_s/f_d = 0.259 \pm 0.015$ .

$$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{125}/\Gamma$$

| VALUE (units $10^{-7}$ )                | DOCUMENT ID | TECN      | COMMENT          |
|-----------------------------------------|-------------|-----------|------------------|
| <b><math>4.3 \pm 2.0 \pm 0.2</math></b> | 1,2 AAIJ    | 17BD LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17BD reports  $[\Gamma(B_s^0 \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [B(J/\psi(1S) \rightarrow p\bar{p})] / [B(K^*(892) \rightarrow (K\pi)^\pm)] = 0.16 \pm 0.07 \pm 0.02$  which we multiply by our best values  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$ ,  $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$ ,  $B(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes  $f_s/f_d = 0.259 \pm 0.015$ .

<sup>2</sup> The branching ratio is given for  $m_{p\bar{p}} < 2.85$  GeV.

$$\Gamma(p\bar{\Lambda}K^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{126}/\Gamma$$

| VALUE (units $10^{-6}$ )                | DOCUMENT ID | TECN      | COMMENT          |
|-----------------------------------------|-------------|-----------|------------------|
| <b><math>5.5 \pm 0.6 \pm 0.8</math></b> | 1,2 AAIJ    | 17AL LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17AL reports  $(5.46 \pm 0.61 \pm 0.82) \times 10^{-6}$  from a measurement of  $[\Gamma(B_s^0 \rightarrow p\bar{\Lambda}K^- + \text{c.c.})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow p\bar{\Lambda}\pi^-)]$  assuming  $B(B^0 \rightarrow p\bar{\Lambda}\pi^-) = (3.14 \pm 0.29) \times 10^{-6}$ .

<sup>2</sup> AAIJ 17AL value represents the sum of  $B_s^0 \rightarrow p\bar{\Lambda}K^-$  and  $B_s^0 \rightarrow \bar{p}\Lambda K^+$  and assumes the fraction  $f_s/f_d = 0.259 \pm 0.015$ .

$$\Gamma(\Lambda_c^- \Lambda \pi^+)/\Gamma_{\text{total}} \quad \Gamma_{127}/\Gamma$$

| VALUE (units $10^{-4}$ )                | DOCUMENT ID            | TECN    | COMMENT                         |
|-----------------------------------------|------------------------|---------|---------------------------------|
| <b><math>3.6 \pm 1.1 \pm 1.2</math></b> | <sup>1</sup> SOLOVIEVA | 13 BELL | $e^+e^- \rightarrow \gamma(4S)$ |

<sup>1</sup> The second error is the total systematic uncertainty including the  $\Lambda_c$  absolute branching fractions and the normalization number of  $B_s$  events.

| $\Gamma(\Lambda_c^- \Lambda_c^+)/\Gamma_{\text{total}}$                                 |     |                   |      |         | $\Gamma_{128}/\Gamma$ |
|-----------------------------------------------------------------------------------------|-----|-------------------|------|---------|-----------------------|
| VALUE                                                                                   | CL% | DOCUMENT ID       | TECN | COMMENT |                       |
| $< 8.0 \times 10^{-5}$                                                                  | 95  | <sup>1</sup> AAIJ | 14AA | LHCB    | $pp$ at 7 TeV         |
| <sup>1</sup> Uses $B(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$ . |     |                   |      |         |                       |

| $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$                                                                        |     |                       |      |         | $\Gamma_{129}/\Gamma$           |
|---------------------------------------------------------------------------------------------------------------------|-----|-----------------------|------|---------|---------------------------------|
| Test for $\Delta B=1$ weak neutral current.                                                                         |     |                       |      |         |                                 |
| VALUE (units $10^{-6}$ )                                                                                            | CL% | DOCUMENT ID           | TECN | COMMENT |                                 |
| $< 3.1$                                                                                                             | 90  | <sup>1</sup> DUTTA    | 15   | BELL    | $e^+e^- \rightarrow \gamma(5S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                       |     |                       |      |         |                                 |
| $< 8.7$                                                                                                             | 90  | <sup>2</sup> WICHT    | 08A  | BELL    | Repl. by DUTTA 15               |
| $< 53$                                                                                                              | 90  | DRUTSKOY              | 07A  | BELL    | Repl. by WICHT 08A              |
| $< 148$                                                                                                             | 90  | <sup>3</sup> ACCIARRI | 95I  | L3      | $e^+e^- \rightarrow Z$          |
| <sup>1</sup> Assumes the fraction of $B_s^{(*)} \bar{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$ . |     |                       |      |         |                                 |
| <sup>2</sup> Assumes $\gamma(5S) \rightarrow B_s^* \bar{B}_s^* = (19.5^{+3.0}_{-2.3})\%$ .                          |     |                       |      |         |                                 |
| <sup>3</sup> ACCIARRI 95I assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = (12.0 \pm 3.0)\%$ .                       |     |                       |      |         |                                 |

| $\Gamma(\phi\gamma)/\Gamma_{\text{total}}$                                                                                                                                                                                                                                                                                                                                                      |     |                    |      |         | $\Gamma_{130}/\Gamma$           |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|--------------------|------|---------|---------------------------------|
| VALUE (units $10^{-6}$ )                                                                                                                                                                                                                                                                                                                                                                        | CL% | DOCUMENT ID        | TECN | COMMENT |                                 |
| <b><math>34 \pm 4</math> OUR AVERAGE</b>                                                                                                                                                                                                                                                                                                                                                        |     |                    |      |         |                                 |
| $36 \pm 5 \pm 7$                                                                                                                                                                                                                                                                                                                                                                                |     | <sup>1</sup> DUTTA | 15   | BELL    | $e^+e^- \rightarrow \gamma(5S)$ |
| $33.8 \pm 3.4 \pm 2.0$                                                                                                                                                                                                                                                                                                                                                                          |     | <sup>2</sup> AAIJ  | 13   | LHCB    | $pp$ at 7 TeV                   |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                                                                                                                                                                                                                                                                                   |     |                    |      |         |                                 |
| $39 \pm 5$                                                                                                                                                                                                                                                                                                                                                                                      |     | <sup>3</sup> AAIJ  | 12AE | LHCB    | Repl. by AAIJ 13                |
| $57^{+18}_{-15} \pm 12_{-11}$                                                                                                                                                                                                                                                                                                                                                                   |     | <sup>4</sup> WICHT | 08A  | BELL    | Repl. by DUTTA 15               |
| $< 390$                                                                                                                                                                                                                                                                                                                                                                                         | 90  | DRUTSKOY           | 07A  | BELL    | $e^+e^- \rightarrow \gamma(5S)$ |
| $< 120$                                                                                                                                                                                                                                                                                                                                                                                         | 90  | ACOSTA             | 02G  | CDF     | $p\bar{p}$ at 1.8 TeV           |
| $< 700$                                                                                                                                                                                                                                                                                                                                                                                         | 90  | <sup>5</sup> ADAM  | 96D  | DLPH    | $e^+e^- \rightarrow Z$          |
| <sup>1</sup> Assumes the fraction of $B_s^{(*)} \bar{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$ . The systematic uncertainty from $f_s$ is $0.6 \times 10^{-5}$ .                                                                                                                                                                                                             |     |                    |      |         |                                 |
| <sup>2</sup> AAIJ 13 reports $[\Gamma(B_s^0 \rightarrow \phi\gamma)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \gamma)] = 0.81 \pm 0.04 \pm 0.07$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0 \gamma) = (4.18 \pm 0.25) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. |     |                    |      |         |                                 |
| <sup>3</sup> Measures $B(B^0 \rightarrow K^{*0} \gamma)/B(B_s \rightarrow \phi\gamma) = 1.12 \pm 0.08(\text{stat})^{+0.06}_{-0.04}(\text{sys})^{+0.09}_{-0.08}(f_s/f_d)$ and uses current world-average value of $B(B^0 \rightarrow K^{*0} \gamma) = (4.33 \pm 0.15) \times 10^{-5}$ .                                                                                                          |     |                    |      |         |                                 |
| <sup>4</sup> Assumes $\gamma(5S) \rightarrow B_s^* \bar{B}_s^* = (19.5^{+3.0}_{-2.3})\%$ .                                                                                                                                                                                                                                                                                                      |     |                    |      |         |                                 |
| <sup>5</sup> ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$ .                                                                                                                                                                                                                                                                                                                 |     |                    |      |         |                                 |

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_{131}/\Gamma$ Test for  $\Delta B = 1$  weak neutral current.

| VALUE (units $10^{-9}$ )                                                      | CL% | DOCUMENT ID               | TECN      | COMMENT                               |
|-------------------------------------------------------------------------------|-----|---------------------------|-----------|---------------------------------------|
| <b><math>3.0 \pm 0.4</math> OUR AVERAGE</b>                                   |     |                           |           |                                       |
| $2.8^{+0.8}_{-0.7}$                                                           |     | <sup>1</sup> AABOUD       | 19L ATLS  | $pp$ at 7, 8, 13 TeV                  |
| $3.0 \pm 0.6^{+0.3}_{-0.2}$                                                   |     | AAIJ                      | 17Al LHCb | $pp$ at 7, 8, 13 TeV                  |
| $13^{+9}_{-7}$                                                                |     | <sup>2</sup> AALTONEN     | 13F CDF   | $p\bar{p}$ at 1.96 TeV                |
| $3.0^{+1.0}_{-0.9}$                                                           |     | <sup>3</sup> CHATRCHYAN   | 13AW CMS  | $pp$ at 7, 8 TeV                      |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                           |           |                                       |
| $0.9^{+1.1}_{-0.8}$                                                           |     | <sup>4</sup> AABOUD       | 16L ATLS  | $pp$ at 7, 8 TeV, Repl. by AABOUD 19L |
| $2.8^{+0.7}_{-0.6}$                                                           |     | <sup>5</sup> KHACHATRY... | 15BE LHC  | $pp$ at 7, 8 TeV                      |
| $3.2^{+1.4+0.5}_{-1.2-0.3}$                                                   |     | <sup>6</sup> AAIJ         | 13B LHCb  | Repl. by AAIJ 13BA                    |
| $2.9^{+1.1+0.3}_{-1.0-0.1}$                                                   |     | <sup>7</sup> AAIJ         | 13BA LHCb | Repl. by KHACHA-TRYAN 15BE            |
| <12                                                                           | 90  | <sup>8</sup> ABAZOV       | 13C D0    | $p\bar{p}$ at 1.96 TeV                |
| <19                                                                           | 90  | <sup>9</sup> AAD          | 12AE ATLS | $pp$ at 7 TeV                         |
| <12                                                                           | 90  | <sup>10</sup> AAIJ        | 12A LHCb  | Repl. by AAIJ 12W                     |
| < 3.8                                                                         | 90  | <sup>11</sup> AAIJ        | 12W LHCb  | Repl. by AAIJ 13B                     |
| < 6.4                                                                         | 90  | <sup>12</sup> CHATRCHYAN  | 12A CMS   | $pp$ at 7 TeV                         |
| <43                                                                           | 90  | <sup>13</sup> AAIJ        | 11B LHCb  | Repl. by AAIJ 12A                     |
| <35                                                                           | 90  | <sup>14</sup> AALTONEN    | 11AG CDF  | $p\bar{p}$ at 1.96 TeV                |
| <16                                                                           | 90  | <sup>15</sup> CHATRCHYAN  | 11T CMS   | Repl. by CHATRCHYAN 12A               |
| <42                                                                           | 90  | <sup>16</sup> ABAZOV      | 10S D0    | $p\bar{p}$ at 1.96 TeV                |

<sup>1</sup> Uses normalization mode  $B(B^+ \rightarrow J/\psi K^+) = (1.010 \pm 0.029) \times 10^{-3}$  and  $B$  production ratio  $f(b \rightarrow B_s^0)/f(b \rightarrow B_d^0) = 0.256 \pm 0.013$ .

<sup>2</sup> Uses normalization mode  $B(B^+ \rightarrow J/\psi K^+) = (10.22 \pm 0.35) \times 10^{-4}$  and  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.28 \pm 0.04$ .

<sup>3</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$  and  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$  for normalization.

<sup>4</sup> This value corresponds to an upper limit of  $< 3.0 \times 10^{-9}$  at 95% C.L. It uses  $f_s/f_d = 0.24 \pm 0.02$ .

<sup>5</sup> Determined from the joint fit to CMS and LHCb data. Uncertainty includes both statistical and systematic component.

<sup>6</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$  and two normalization modes:  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$  and  $B(B^0 \rightarrow K^+\pi^-) = (1.94 \pm 0.06) \times 10^{-5}$ .

<sup>7</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$  and normalization modes  $B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+$  and  $B^0 \rightarrow K^+\pi^-$ .

<sup>8</sup> Uses normalization mode  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$  and  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.263 \pm 0.017$ .

<sup>9</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.75 \pm 0.29$  and  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+\mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$ .



- <sup>10</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$  and three normalization modes  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ ,  $B(B^0 \rightarrow K^+ \pi^-) = (1.94 \pm 0.06) \times 10^{-5}$ , and  $B(B_s^0 \rightarrow J/\psi \phi \rightarrow \mu^+ \mu^- K^+ K^-) = (3.4 \pm 0.9) \times 10^{-5}$ .
- <sup>11</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$  and three normalization modes of  $B^+ \rightarrow J/\psi K^+$ ,  $B^0 \rightarrow K^+ \pi^-$ , and  $B_s^0 \rightarrow J/\psi \phi$ .
- <sup>12</sup> Uses  $f_s/f_u = 0.267 \pm 0.021$  and  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$ .
- <sup>13</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.71 \pm 0.47$  and three normalization modes.
- <sup>14</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.55 \pm 0.47$  and  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ .
- <sup>15</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.55 \pm 0.42$  and  $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$ .
- <sup>16</sup> Uses  $B$  production ratio  $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.86 \pm 0.59$ , and the number of  $B^+ \rightarrow J/\psi K^+$  decays.

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{132}/\Gamma$ Test for  $\Delta B = 1$  weak neutral current.

| VALUE                                                                                                      | CL% | DOCUMENT ID           | TECN    | COMMENT                |
|------------------------------------------------------------------------------------------------------------|-----|-----------------------|---------|------------------------|
| $<2.8 \times 10^{-7}$                                                                                      | 90  | AALTONEN              | 09P CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                              |     |                       |         |                        |
| $<5.4 \times 10^{-5}$                                                                                      | 90  | <sup>1</sup> ACCIARRI | 97B L3  | $e^+e^- \rightarrow Z$ |
| <sup>1</sup> ACCIARRI 97B assume PDG 96 production fractions for $B^+$ , $B^0$ , $B_s$ , and $\Lambda_b$ . |     |                       |         |                        |

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$  $\Gamma_{133}/\Gamma$ 

| VALUE                                                                       | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------------------------------------------|-----|-------------------|-----------|------------------|
| $<6.8 \times 10^{-3}$                                                       | 95  | <sup>1</sup> AAIJ | 17AJ LHCB | $pp$ at 7, 8 TeV |
| <sup>1</sup> Assuming no contribution from $B^0 \rightarrow \tau^+\tau^-$ . |     |                   |           |                  |

 $\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_{134}/\Gamma$ 

| VALUE                                                                         | CL% | DOCUMENT ID       | TECN      | COMMENT           |
|-------------------------------------------------------------------------------|-----|-------------------|-----------|-------------------|
| $<2.5 \times 10^{-9}$                                                         | 95  | AAIJ              | 17N LHCB  | $pp$ at 7, 8 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                   |           |                   |
| $<1.2 \times 10^{-8}$                                                         | 90  | <sup>1</sup> AAIJ | 13AW LHCB | Repl. by AAIJ 17N |
| <sup>1</sup> Also reports a limit of $<1.6 \times 10^{-8}$ at 95% CL.         |     |                   |           |                   |

 $\Gamma(SP, S \rightarrow \mu^+\mu^-, P \rightarrow \mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_{135}/\Gamma$ Here  $S$  and  $P$  are the hypothetical scalar and pseudoscalar particles with masses of 2.5 GeV/ $c^2$  and 214.3 MeV/ $c^2$ , respectively.

| VALUE                                                                         | CL% | DOCUMENT ID       | TECN      | COMMENT           |
|-------------------------------------------------------------------------------|-----|-------------------|-----------|-------------------|
| $<2.2 \times 10^{-9}$                                                         | 95  | AAIJ              | 17N LHCB  | $pp$ at 7, 8 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |                   |           |                   |
| $<1.2 \times 10^{-8}$                                                         | 90  | <sup>1</sup> AAIJ | 13AW LHCB | Repl. by AAIJ 17N |
| <sup>1</sup> Also reports a limit of $<1.6 \times 10^{-8}$ at 95% CL.         |     |                   |           |                   |

$\Gamma(\phi(1020)\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{136}/\Gamma$ 
Test for  $\Delta B = 1$  weak neutral current.

| VALUE (units $10^{-7}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                    |    |                     |         |                        |
|--------------------|----|---------------------|---------|------------------------|
| $<32$              | 90 | <sup>1</sup> ABAZOV | 06G D0  | $p\bar{p}$ at 1.96 TeV |
| $<4.7 \times 10^2$ | 90 | ACOSTA              | 02D CDF | $p\bar{p}$ at 1.8 TeV  |

<sup>1</sup> Uses  $B(B_S^0 \rightarrow J/\psi\phi) = 9.3 \times 10^{-4}$ .
 $\Gamma(\phi(1020)\mu^+\mu^-)/\Gamma(J/\psi(1S)\phi)$   $\Gamma_{136}/\Gamma_{49}$ 

| VALUE (units $10^{-3}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

**0.76 ± 0.09 OUR AVERAGE** Error includes scale factor of 1.9.

|                                     |  |      |           |                  |
|-------------------------------------|--|------|-----------|------------------|
| $0.741^{+0.042}_{-0.040} \pm 0.029$ |  | AAIJ | 15AQ LHCb | $pp$ at 7, 8 TeV |
|-------------------------------------|--|------|-----------|------------------|

|                          |  |          |          |                        |
|--------------------------|--|----------|----------|------------------------|
| $1.13 \pm 0.19 \pm 0.07$ |  | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
|--------------------------|--|----------|----------|------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                     |  |                   |          |                    |
|-------------------------------------|--|-------------------|----------|--------------------|
| $0.674^{+0.061}_{-0.056} \pm 0.016$ |  | <sup>1</sup> AAIJ | 13X LHCb | Repl. by AAIJ 15AQ |
|-------------------------------------|--|-------------------|----------|--------------------|

|                          |  |          |         |                        |
|--------------------------|--|----------|---------|------------------------|
| $1.11 \pm 0.25 \pm 0.09$ |  | AALTONEN | 11L CDF | Repl. by AALTONEN 11AI |
|--------------------------|--|----------|---------|------------------------|

|        |    |          |         |                       |
|--------|----|----------|---------|-----------------------|
| $<2.3$ | 90 | AALTONEN | 09B CDF | Repl. by AALTONEN 11L |
|--------|----|----------|---------|-----------------------|

<sup>1</sup> Replaced by AAIJ 15AQ.
 $\Gamma(\bar{K}^*(892)^0\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{137}/\Gamma$ 

| VALUE (units $10^{-8}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

|                        |                   |           |                      |
|------------------------|-------------------|-----------|----------------------|
| <b>2.9 ± 1.0 ± 0.4</b> | <sup>1</sup> AAIJ | 18AB LHCb | $pp$ at 7, 8, 13 TeV |
|------------------------|-------------------|-----------|----------------------|

<sup>1</sup> Normalizes to  $B(B^0 \rightarrow J/\psi K^{*0}) = 1.19 \pm 0.01 \pm 0.08\%$  and  $B(J/\psi \rightarrow \mu^+\mu^-) = 5.96 \pm 0.03\%$ , and uses  $f_S/f_d = 0.259 \pm 0.015$ .
 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{138}/\Gamma$ 

| VALUE (units $10^{-8}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

|                        |                   |          |                  |
|------------------------|-------------------|----------|------------------|
| <b>8.4 ± 1.6 ± 0.3</b> | <sup>1</sup> AAIJ | 15S LHCb | $pp$ at 7, 8 TeV |
|------------------------|-------------------|----------|------------------|

<sup>1</sup> AAIJ 15S reports  $(8.6 \pm 1.5 \pm 0.7 \pm 0.7) \times 10^{-8}$  from a measurement of  $[\Gamma(B_S^0 \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)]$  assuming  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.3 \pm 0.1) \times 10^{-3}$ , which we rescale to our best value  $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\phi\nu\bar{\nu})/\Gamma_{\text{total}}$   $\Gamma_{139}/\Gamma$ 
Test for  $\Delta B = 1$  weak neutral current.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                                  |    |                   |          |                        |
|----------------------------------|----|-------------------|----------|------------------------|
| <b>&lt;5.4 × 10<sup>-3</sup></b> | 90 | <sup>1</sup> ADAM | 96D DLPH | $e^+e^- \rightarrow Z$ |
|----------------------------------|----|-------------------|----------|------------------------|

<sup>1</sup> ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .
 $\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$   $\Gamma_{140}/\Gamma$ 

Test of lepton family number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                                  |    |                   |          |                  |
|----------------------------------|----|-------------------|----------|------------------|
| <b>&lt;5.4 × 10<sup>-9</sup></b> | 90 | <sup>1</sup> AAIJ | 18T LHCb | $pp$ at 7, 8 TeV |
|----------------------------------|----|-------------------|----------|------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |    |                       |          |                        |
|-----------------------|----|-----------------------|----------|------------------------|
| $<1.1 \times 10^{-8}$ | 90 | <sup>2</sup> AAIJ     | 13BMLHCB | Repl. by AAIJ 18T      |
| $<2.0 \times 10^{-7}$ | 90 | AALTONEN              | 09P CDF  | $p\bar{p}$ at 1.96 TeV |
| $<6.1 \times 10^{-6}$ | 90 | ABE                   | 98V CDF  | Repl. by AALTONEN 09P  |
| $<4.1 \times 10^{-5}$ | 90 | <sup>3</sup> ACCIARRI | 97B L3   | $e^+e^- \rightarrow Z$ |

<sup>1</sup> AAIJ 18T uses normalization modes  $B(B^0 \rightarrow K^+\pi^-) = (19.6 \pm 0.5) \times 10^{-6}$  and  $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$  with  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$ . The upper limit increases to  $6 \times 10^{-9}$  with the assumption of  $B_L$ -dominated decay amplitude.

<sup>2</sup> Uses normalization mode  $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$  and  $B$  production ratio  $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ .

<sup>3</sup> ACCIARRI 97B assume PDG 96 production fractions for  $B^+$ ,  $B^0$ ,  $B_s$ , and  $\Lambda_b$ .

### $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ $\Gamma_{141}/\Gamma$

| VALUE                                      | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|--------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt;4.2 \times 10^{-5}</math></b> | 95  | <sup>1</sup> AAIJ | 19AK LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Assuming no contribution from  $B^0 \rightarrow \mu^\pm \tau^\mp$ .

## POLARIZATION IN $B_s^0$ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal ( $L$ ), or both are transverse and parallel ( $\parallel$ ), or perpendicular ( $\perp$ ) to each other with the parameters  $\Gamma_L/\Gamma$ ,  $\Gamma_\perp/\Gamma$ , and the relative phases  $\phi_\parallel$  and  $\phi_\perp$ . In decays involving two tensor mesons, the transverse polarization states are described by parameters  $\Gamma_{\parallel 1}$ ,  $\Gamma_{\parallel 2}$ ,  $\Gamma_{\perp 1}$ ,  $\Gamma_{\perp 2}$  and their relative phases  $\phi_{\parallel 1}$ ,  $\phi_{\parallel 2}$ ,  $\phi_{\perp 1}$ ,  $\phi_{\perp 2}$ . See also the review on "Polarization in  $B$  Decays."

### $\Gamma_L/\Gamma$ in $B_s^0 \rightarrow D_s^* \rho^+$

| VALUE                                              | DOCUMENT ID | TECN | COMMENT                                |
|----------------------------------------------------|-------------|------|----------------------------------------|
| <b><math>1.05^{+0.08+0.03}_{-0.10-0.04}</math></b> | LOUVOT      | 10   | BELL $e^+e^- \rightarrow \Upsilon(5S)$ |

### $\Gamma_L/\Gamma$ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE                                             | DOCUMENT ID                         | TECN      | COMMENT                |
|---------------------------------------------------|-------------------------------------|-----------|------------------------|
| <b><math>0.5199 \pm 0.0035</math> OUR AVERAGE</b> | Error includes scale factor of 1.1. |           |                        |
| $0.5186 \pm 0.0029 \pm 0.0024$                    | AAIJ                                | 19Q LHCB  | $pp$ at 13 TeV         |
| $0.522 \pm 0.003 \pm 0.007$                       | <sup>1</sup> AAD                    | 16AP ATLS | $pp$ at 7, 8 TeV       |
| $0.510 \pm 0.005 \pm 0.011$                       | KHACHATRY...                        | 16S CMS   | $pp$ at 8 TeV          |
| $0.524 \pm 0.013 \pm 0.015$                       | <sup>2</sup> AALTONEN               | 12D CDF   | $p\bar{p}$ at 1.96 TeV |
| $0.558^{+0.017}_{-0.019}$                         | <sup>2,3</sup> ABAZOV               | 12D D0    | $p\bar{p}$ at 1.96 TeV |
| $0.61 \pm 0.14 \pm 0.02$                          | <sup>4</sup> AFFOLDER               | 00N CDF   | $p\bar{p}$ at 1.8 TeV  |
| $0.56 \pm 0.21^{+0.02}_{-0.04}$                   | ABE                                 | 95Z CDF   | $p\bar{p}$ at 1.8 TeV  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                |                       |      |      |                       |
|--------------------------------|-----------------------|------|------|-----------------------|
| $0.5241 \pm 0.0034 \pm 0.0067$ | AAIJ                  | 15I  | LHCB | Repl. by AAIJ 19Q     |
| $0.529 \pm 0.006 \pm 0.012$    | <sup>1</sup> AAD      | 14U  | ATLS | Repl. by AAD 16AP     |
| $0.539 \pm 0.014 \pm 0.016$    | <sup>2</sup> AAD      | 12CV | ATLS | Repl. by AAD 14U      |
| $0.555 \pm 0.027 \pm 0.006$    | <sup>5</sup> ABAZOV   | 09E  | D0   | Repl. by ABAZOV 12D   |
| $0.531 \pm 0.020 \pm 0.007$    | <sup>2</sup> AALTONEN | 08J  | CDF  | Repl. by AALTONEN 12D |
| $0.62 \pm 0.06 \pm 0.01$       | ACOSTA                | 05   | CDF  | Repl. by AALTONEN 08J |

<sup>1</sup> Measured using the flavor tagged, time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>2</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>3</sup> The error includes both statistical and systematic uncertainties.

<sup>4</sup> AFFOLDER 00N measurements are based on 40  $B_s^0$  candidates obtained from a data sample of  $89 \text{ pb}^{-1}$ . The  $P$ -wave fraction is found to be  $0.23 \pm 0.19 \pm 0.04$ .

<sup>5</sup> Measured the angular and lifetime parameters for the time-dependent angular untagged decays  $B_d^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi \phi$ .

### $\Gamma_L/\Gamma$ in $B_s^0 \rightarrow D_s^{*+} D_s^{*-}$

| VALUE                           | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------|-------------|------|---------------------------------------|
| $0.06^{+0.18}_{-0.17} \pm 0.03$ | ESEN        | 13   | BELL $e^+ e^- \rightarrow \gamma(5S)$ |

### $\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE                                           | DOCUMENT ID           | TECN | COMMENT                    |
|-------------------------------------------------|-----------------------|------|----------------------------|
| <b><math>0.228 \pm 0.007</math> OUR AVERAGE</b> |                       |      |                            |
| $0.227 \pm 0.004 \pm 0.006$                     | <sup>1</sup> AAD      | 16AP | ATLS $pp$ at 7, 8 TeV      |
| $0.231 \pm 0.014 \pm 0.015$                     | <sup>2</sup> AALTONEN | 12D  | CDF $p\bar{p}$ at 1.96 TeV |
| $0.231^{+0.024}_{-0.030}$                       | <sup>2,3</sup> ABAZOV | 12D  | D0 $p\bar{p}$ at 1.96 TeV  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |                       |      |      |                       |
|-----------------------------|-----------------------|------|------|-----------------------|
| $0.220 \pm 0.008 \pm 0.009$ | <sup>1</sup> AAD      | 14U  | ATLS | Repl. by AAD 16AP     |
| $0.224 \pm 0.010 \pm 0.009$ | <sup>2</sup> AAD      | 12CV | ATLS | Repl. by AAD 14U      |
| $0.244 \pm 0.032 \pm 0.014$ | <sup>4</sup> ABAZOV   | 09E  | D0   | Repl. by ABAZOV 12D   |
| $0.230 \pm 0.029 \pm 0.011$ | <sup>2</sup> AALTONEN | 08J  | CDF  | Repl. by AALTONEN 12D |
| $0.260 \pm 0.084 \pm 0.013$ | ACOSTA                | 05   | CDF  | Repl. by AALTONEN 08J |

<sup>1</sup> Measured using a tagged, time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>2</sup> Measured using the time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi \phi$  decays.

<sup>3</sup> The error includes both statistical and systematic uncertainties.

<sup>4</sup> Measured the angular and lifetime parameters for the time-dependent angular untagged decays  $B_d^0 \rightarrow J/\psi K^{*0}$  and  $B_s^0 \rightarrow J/\psi \phi$ .

### $\Gamma_{\perp}/\Gamma$ in $B_s^0 \rightarrow J/\psi(1S)\phi$

| VALUE                                           | DOCUMENT ID  | TECN | COMMENT             |
|-------------------------------------------------|--------------|------|---------------------|
| <b><math>0.245 \pm 0.004</math> OUR AVERAGE</b> |              |      |                     |
| $0.2456 \pm 0.0040 \pm 0.0019$                  | AAIJ         | 19Q  | LHCB $pp$ at 13 TeV |
| $0.243 \pm 0.008 \pm 0.012$                     | KHACHATRY... | 16S  | CMS $pp$ at 8 TeV   |

• • • We do not use the following data for averages, fits, limits, etc. • • •


|                                |      |     |      |                   |
|--------------------------------|------|-----|------|-------------------|
| $0.2504 \pm 0.0049 \pm 0.0036$ | AAIJ | 15I | LHCB | Repl. by AAIJ 19Q |
|--------------------------------|------|-----|------|-------------------|

$\phi_{\parallel}$  in  $B_s^0 \rightarrow J/\psi(1S)\phi$ 

| VALUE (rad)                                                                   | DOCUMENT ID         | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|---------------------|-----------|------------------------|
| <b><math>3.10 \pm 0.06</math> OUR AVERAGE</b>                                 |                     |           |                        |
| $3.06^{+0.08}_{-0.07} \pm 0.04$                                               | AAIJ                | 19Q LHCb  | $pp$ at 13 TeV         |
| $3.15 \pm 0.10 \pm 0.05$                                                      | AAD                 | 16AP ATLS | $pp$ at 7, 8 TeV       |
| $3.48^{+0.07}_{-0.09} \pm 0.68$                                               | KHACHATRY...16S     | CMS       | $pp$ at 8 TeV          |
| $3.15 \pm 0.22$                                                               | <sup>1</sup> ABAZOV | 12D D0    | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |           |                        |
| $3.26^{+0.10+0.06}_{-0.17-0.07}$                                              | AAIJ                | 15I LHCb  | Repl. by AAIJ 19Q      |
| $2.72^{+1.12}_{-0.27} \pm 0.26$                                               | ABAZOV              | 09E D0    | Repl. by ABAZOV 12D    |

<sup>1</sup> The error includes both statistical and systematic uncertainties. $\phi_{\perp}$  in  $B_s^0 \rightarrow J/\psi(1S)\phi$ 

| VALUE (rad)                                                                   | DOCUMENT ID                                                 | TECN      | COMMENT           |
|-------------------------------------------------------------------------------|-------------------------------------------------------------|-----------|-------------------|
| <b><math>2.9 \pm 0.4</math> OUR AVERAGE</b>                                   | Error includes scale factor of 2.7. See the ideogram below. |           |                   |
| $2.64 \pm 0.13 \pm 0.10$                                                      | AAIJ                                                        | 19Q LHCb  | $pp$ at 13 TeV    |
| $4.15 \pm 0.32 \pm 0.16$                                                      | <sup>1</sup> AAD                                            | 16AP ATLS | $pp$ at 7, 8 TeV  |
| $2.98 \pm 0.36 \pm 0.66$                                                      | KHACHATRY...16S                                             | CMS       | $pp$ at 8 TeV     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                                             |           |                   |
| $3.08^{+0.14}_{-0.15} \pm 0.06$                                               | AAIJ                                                        | 15I LHCb  | Repl. by AAIJ 19Q |
| $3.89 \pm 0.47 \pm 0.11$                                                      | <sup>1</sup> AAD                                            | 14U ATLS  | Repl. by AAD 16AP |

<sup>1</sup> Measured using a tagged, time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays.

$\Gamma_{\perp}/\Gamma$  in  $B_s^0 \rightarrow \psi(2S)\phi$ 

| VALUE                               | DOCUMENT ID       | TECN      | COMMENT          |
|-------------------------------------|-------------------|-----------|------------------|
| $0.264^{+0.024}_{-0.023} \pm 0.002$ | <sup>1</sup> AAIJ | 16AK LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S)\phi$  decays.

 $\phi_{\parallel}$  in  $B_s^0 \rightarrow \psi(2S)\phi$ 

| VALUE (rad)                     | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $3.67^{+0.13}_{-0.18} \pm 0.03$ | <sup>1</sup> AAIJ | 16AK LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S)\phi$  decays.

 $\phi_{\perp}$  in  $B_s^0 \rightarrow \psi(2S)\phi$ 

| VALUE (rad)                     | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------|-------------------|-----------|------------------|
| $3.29^{+0.43}_{-0.39} \pm 0.04$ | <sup>1</sup> AAIJ | 16AK LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S)\phi$  decays.

 $\Gamma_L/\Gamma$  for  $B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0$ 

Longitudinal polarization fraction, equals to  $f_L$  using notation of “Polarization in  $B$  decays” review.

| VALUE                                                                         | DOCUMENT ID       | TECN      | COMMENT            |
|-------------------------------------------------------------------------------|-------------------|-----------|--------------------|
| $0.497 \pm 0.025 \pm 0.025$                                                   | AAIJ              | 15AV LHCB | $pp$ at 7, 8 TeV   |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                   |           |                    |
| $0.50 \pm 0.08 \pm 0.02$                                                      | <sup>1</sup> AAIJ | 12AP LHCB | Repl. by AAIJ 15AV |

<sup>1</sup> The non-resonant  $K\pi$  background contributions are subtracted. Also reports an  $S$ -wave amplitude  $|A_S|^2 = 0.07^{+0.15}_{-0.07}$ .

 $\Gamma_{\parallel}/\Gamma$  for  $B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0$ 

Parallel polarization fraction, equals to  $1 - f_L - f_{\perp}$  using notation of “Polarization in  $B$  decays” review.

| VALUE                                                                         | DOCUMENT ID       | TECN      | COMMENT            |
|-------------------------------------------------------------------------------|-------------------|-----------|--------------------|
| $0.179 \pm 0.027 \pm 0.013$                                                   | AAIJ              | 15AV LHCB | $pp$ at 7, 8 TeV   |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                   |           |                    |
| $0.19^{+0.10}_{-0.08} \pm 0.02$                                               | <sup>1</sup> AAIJ | 12AP LHCB | Repl. by AAIJ 15AV |

<sup>1</sup> The non-resonant  $K\pi$  background contributions are subtracted. Also reports an  $S$ -wave amplitude  $|A_S|^2 = 0.07^{+0.15}_{-0.07}$ .

 $\Gamma_{\parallel}/\Gamma$  of  $K^*(892)^0$  in  $B_s^0 \rightarrow \psi(2S)\bar{K}^*(892)^0$ 

| VALUE                       | DOCUMENT ID | TECN     | COMMENT          |
|-----------------------------|-------------|----------|------------------|
| $0.524 \pm 0.056 \pm 0.029$ | AAIJ        | 15U LHCB | $pp$ at 7, 8 TeV |

$\Gamma_L/\Gamma$  in  $B_s^0 \rightarrow \phi\phi$ 

| VALUE                                                                         | DOCUMENT ID | TECN      | COMMENT                 |
|-------------------------------------------------------------------------------|-------------|-----------|-------------------------|
| <b><math>0.378 \pm 0.013</math> OUR AVERAGE</b>                               |             |           |                         |
| $0.381 \pm 0.007 \pm 0.012$                                                   | AAIJ        | 19AP LHCb | $pp$ at 7, 8 and 13 TeV |
| $0.348 \pm 0.041 \pm 0.021$                                                   | AALTONEN    | 11AN CDF  | $p\bar{p}$ at 1.96 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |           |                         |
| $0.364 \pm 0.012 \pm 0.009$                                                   | AAIJ        | 14AE LHCb | Repl. by AAIJ 19AP      |
| $0.365 \pm 0.022 \pm 0.012$                                                   | AAIJ        | 12P LHCb  | Repl. by AAIJ 14AE      |

 $\Gamma_{\perp}/\Gamma$  in  $B_s^0 \rightarrow \phi\phi$ 

| VALUE                                                                         | DOCUMENT ID       | TECN      | COMMENT                 |
|-------------------------------------------------------------------------------|-------------------|-----------|-------------------------|
| <b><math>0.292 \pm 0.009</math> OUR AVERAGE</b>                               |                   |           |                         |
| $0.290 \pm 0.008 \pm 0.005$                                                   | <sup>1</sup> AAIJ | 19AP LHCb | $pp$ at 7, 8 and 13 TeV |
| $0.365 \pm 0.044 \pm 0.027$                                                   | AALTONEN          | 11AN CDF  | $p\bar{p}$ at 1.96 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                   |           |                         |
| $0.305 \pm 0.013 \pm 0.005$                                                   | AAIJ              | 14AE LHCb | Repl. by AAIJ 19AP      |
| $0.291 \pm 0.024 \pm 0.010$                                                   | AAIJ              | 12P LHCb  | Repl. by AAIJ 14AE      |

<sup>1</sup> Note: in the summary of AAIJ 19AP the systematic uncertainty is 0.007. We take the systematic uncertainty as given in Table 5 in the paper.

 $\phi_{\parallel}$  in  $B_s^0 \rightarrow \phi\phi$ 

| VALUE (rad)                                                                   | DOCUMENT ID           | TECN      | COMMENT                 |
|-------------------------------------------------------------------------------|-----------------------|-----------|-------------------------|
| <b><math>2.56 \pm 0.06</math> OUR AVERAGE</b>                                 |                       |           |                         |
| $2.559 \pm 0.045 \pm 0.033$                                                   | AAIJ                  | 19AP LHCb | $pp$ at 7, 8 and 13 TeV |
| $2.71^{+0.31}_{-0.36} \pm 0.22$                                               | <sup>1</sup> AALTONEN | 11AN CDF  | $p\bar{p}$ at 1.96 TeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                         |
| $2.54 \pm 0.07 \pm 0.09$                                                      | <sup>2</sup> AAIJ     | 14AE LHCb | Repl. by AAIJ 19AP      |
| $2.57 \pm 0.15 \pm 0.06$                                                      | <sup>3</sup> AAIJ     | 12P LHCb  | Repl. by AAIJ 14AE      |

<sup>1</sup> AALTONEN 11AN quotes  $\cos\phi_{\parallel} = -0.91^{+0.15}_{-0.13} \pm 0.09$  which we convert to  $\phi_{\parallel}$  taking the smaller solution.

<sup>2</sup> AAIJ 14AE reports measurement of  $\phi_{\perp}$  and  $\phi_{\perp} - \phi_{\parallel}$ , which we convert into  $\phi_{\parallel}$ . Statistical uncertainty includes correlation between measured parameters, while systematic uncertainties are assumed uncorrelated.

<sup>3</sup> AAIJ 12P quotes  $\cos\phi_{\parallel} = -0.844 \pm 0.068 \pm 0.029$  which we convert to  $\phi_{\parallel}$ , taking the smaller solution.

 $\phi_{\perp}$  in  $B_s^0 \rightarrow \phi\phi$ 

| VALUE (rad)                                                                   | DOCUMENT ID | TECN      | COMMENT                 |
|-------------------------------------------------------------------------------|-------------|-----------|-------------------------|
| <b><math>2.818 \pm 0.178 \pm 0.073</math></b>                                 | AAIJ        | 19AP LHCb | $pp$ at 7, 8 and 13 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |           |                         |
| $2.67 \pm 0.23 \pm 0.07$                                                      | AAIJ        | 14AE LHCb | Repl. by AAIJ 19AP      |

 $\Gamma_L/\Gamma$  in  $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$ 

| VALUE                                         | DOCUMENT ID       | TECN     | COMMENT             |
|-----------------------------------------------|-------------------|----------|---------------------|
| <b><math>0.240 \pm 0.031 \pm 0.025</math></b> | <sup>1</sup> AAIJ | 19L LHCb | $pp$ at 7 and 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |                   |      |      |                    |
|-----------------------------|-------------------|------|------|--------------------|
| $0.208 \pm 0.032 \pm 0.046$ | <sup>2</sup> AAIJ | 18S  | LHCB | Repl. by AAIJ 19L  |
| $0.201 \pm 0.057 \pm 0.040$ | <sup>3</sup> AAIJ | 15AF | LHCB | Repl. by AAIJ 18S  |
| $0.31 \pm 0.12 \pm 0.04$    | AAIJ              | 12F  | LHCB | Repl. by AAIJ 15AF |

<sup>1</sup> Untagged and time-integrated analysis within 150 MeV of the  $K^{*0}$  mass.

<sup>2</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions.

<sup>3</sup> Measured in angular analysis, which takes into account  $S$ -wave contributions.

### $\Gamma_{\perp}/\Gamma$ in $B_s^0 \rightarrow K^{*0} \bar{K}^{*0}$

| VALUE                                      | DOCUMENT ID | TECN | COMMENT            |
|--------------------------------------------|-------------|------|--------------------|
| <b><math>0.38 \pm 0.11 \pm 0.04</math></b> | AAIJ        | 12F  | LHCB $pp$ at 7 TeV |

### $\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow K^{*}(892)^0 \bar{K}^{*}(892)^0$

| VALUE                                         | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------------------------------|-------------------|------|-----------------------|
| <b><math>0.297 \pm 0.029 \pm 0.042</math></b> | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |      |      |      |                   |
|-----------------------------|------|------|------|-------------------|
| $0.215 \pm 0.046 \pm 0.015$ | AAIJ | 15AF | LHCB | Repl. by AAIJ 18S |
|-----------------------------|------|------|------|-------------------|

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions.

### $\Phi_{\parallel}$ in $B_s^0 \rightarrow K^{*}(892)^0 \bar{K}^{*}(892)^0$

| VALUE                                      | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------------------------|-------------------|------|-----------------------|
| <b><math>2.40 \pm 0.11 \pm 0.33</math></b> | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                          |      |      |      |                   |
|--------------------------|------|------|------|-------------------|
| $5.31 \pm 0.24 \pm 0.14$ | AAIJ | 15AF | LHCB | Repl. by AAIJ 18S |
|--------------------------|------|------|------|-------------------|

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions.

### $\Phi_{\perp}$ in $B_s^0 \rightarrow K^{*}(892)^0 \bar{K}^{*}(892)^0$

| VALUE (rad)                                | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------------------------|-------------------|------|-----------------------|
| <b><math>2.62 \pm 0.26 \pm 0.64</math></b> | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions.

### $\Gamma_L/\Gamma$ in $B_s^0 \rightarrow \phi \bar{K}^{*0}$

| VALUE                                      | DOCUMENT ID | TECN | COMMENT            |
|--------------------------------------------|-------------|------|--------------------|
| <b><math>0.51 \pm 0.15 \pm 0.07</math></b> | AAIJ        | 13BW | LHCB $pp$ at 7 TeV |

### $\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow \phi \bar{K}^{*0}$

| VALUE                                      | DOCUMENT ID | TECN | COMMENT            |
|--------------------------------------------|-------------|------|--------------------|
| <b><math>0.21 \pm 0.11 \pm 0.02</math></b> | AAIJ        | 13BW | LHCB $pp$ at 7 TeV |

### $\phi_{\parallel}$ in $B_s^0 \rightarrow \phi \bar{K}^{*0}$

| VALUE (rad)                                | DOCUMENT ID       | TECN | COMMENT            |
|--------------------------------------------|-------------------|------|--------------------|
| <b><math>1.75 \pm 0.53 \pm 0.29</math></b> | <sup>1</sup> AAIJ | 13BW | LHCB $pp$ at 7 TeV |

<sup>1</sup> Measures  $\cos(\phi_{\parallel}) = -0.18 \pm 0.52 \pm 0.29$ , which we convert to  $\phi_{\parallel}$  by taking the smaller solution.

### $\Gamma_L/\Gamma$ in $B_s^0 \rightarrow \bar{D}^{*0} \phi$

| VALUE                                      | DOCUMENT ID | TECN | COMMENT                  |
|--------------------------------------------|-------------|------|--------------------------|
| <b><math>0.73 \pm 0.15 \pm 0.04</math></b> | AAIJ        | 18AY | LHCB $pp$ at 7 and 8 TeV |



$\Gamma_L/\Gamma$  in  $B_s^0 \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                       | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------------|-------------------|------|-----------------------|
| $0.911 \pm 0.020 \pm 0.165$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_{\parallel}/\Gamma$  in  $B_s^0 \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                       | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------------|-------------------|------|-----------------------|
| $0.012 \pm 0.008 \pm 0.053$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_L/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}^*(892)^0$ 

| VALUE                    | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------|-------------------|------|-----------------------|
| $0.62 \pm 0.16 \pm 0.25$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_{\parallel}/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}^*(892)^0$ 

| VALUE                    | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------|-------------------|------|-----------------------|
| $0.24 \pm 0.10 \pm 0.14$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_L/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                    | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------|-------------------|------|-----------------------|
| $0.25 \pm 0.14 \pm 0.18$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_{\parallel 1}/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                    | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------|-------------------|------|-----------------------|
| $0.17 \pm 0.11 \pm 0.14$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_{\perp 1}/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                    | DOCUMENT ID       | TECN | COMMENT               |
|--------------------------|-------------------|------|-----------------------|
| $0.30 \pm 0.18 \pm 0.21$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $\Gamma_{\parallel 2}/\Gamma$  in  $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$ 

| VALUE                       | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------------|-------------------|------|-----------------------|
| $0.015 \pm 0.033 \pm 0.107$ | <sup>1</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in angular analysis, which takes into account  $S$ -,  $P$ - and  $D$ -wave. contributions. $F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$  ( $0.10 < q^2 < 2.00 \text{ GeV}^2/c^4$ )

| VALUE                           | DOCUMENT ID | TECN | COMMENT               |
|---------------------------------|-------------|------|-----------------------|
| $0.20^{+0.08}_{-0.09} \pm 0.02$ | AAIJ        | 15AQ | LHCB $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |      |     |                         |
|---------------------------------|------|-----|-------------------------|
| $0.37^{+0.19}_{-0.17} \pm 0.07$ | AAIJ | 13X | LHCB Repl. by AAIJ 15AQ |
|---------------------------------|------|-----|-------------------------|

**$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (2.00 < q^2 < 5.0 \text{ GeV}^2/c^4)$** 

| VALUE                           | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------|-------------|-----------|------------------|
| $0.68^{+0.16}_{-0.13} \pm 0.03$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |                   |          |                    |
|---------------------------------|-------------------|----------|--------------------|
| $0.53^{+0.25}_{-0.23} \pm 0.10$ | <sup>1</sup> AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in  $2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$ .

 **$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (5.0 < q^2 < 8.0 \text{ GeV}^2/c^4)$** 

| VALUE                           | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------|-------------|-----------|------------------|
| $0.54^{+0.10}_{-0.09} \pm 0.02$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |                   |          |                    |
|---------------------------------|-------------------|----------|--------------------|
| $0.81^{+0.11}_{-0.13} \pm 0.05$ | <sup>1</sup> AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in  $4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$ .

 **$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (11.0 < q^2 < 12.5 \text{ GeV}^2/c^4)$** 

| VALUE                    | DOCUMENT ID | TECN      | COMMENT          |
|--------------------------|-------------|-----------|------------------|
| $0.29 \pm 0.11 \pm 0.04$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |                   |          |                    |
|---------------------------------|-------------------|----------|--------------------|
| $0.33^{+0.14}_{-0.12} \pm 0.06$ | <sup>1</sup> AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in  $10.09 < q^2 < 12.90 \text{ GeV}^2/c^4$ .

 **$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (15.0 < q^2 < 17.0 \text{ GeV}^2/c^4)$** 

| VALUE                           | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------|-------------|-----------|------------------|
| $0.23^{+0.09}_{-0.08} \pm 0.02$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |                   |          |                    |
|---------------------------------|-------------------|----------|--------------------|
| $0.34^{+0.18}_{-0.17} \pm 0.07$ | <sup>1</sup> AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in  $14.18 < q^2 < 16 \text{ GeV}^2/c^4$ .

 **$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (17.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$** 

| VALUE                           | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------|-------------|-----------|------------------|
| $0.40^{+0.13}_{-0.15} \pm 0.02$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |                   |          |                    |
|---------------------------------|-------------------|----------|--------------------|
| $0.16^{+0.17}_{-0.10} \pm 0.07$ | <sup>1</sup> AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in  $16.0 < q^2 < 19.0 \text{ GeV}^2/c^4$ .

 **$F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-) (1.00 < q^2 < 6.00 \text{ GeV}^2/c^4)$** 

| VALUE                           | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------|-------------|-----------|------------------|
| $0.63^{+0.09}_{-0.09} \pm 0.03$ | AAIJ        | 15AQ LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                 |      |          |                    |
|---------------------------------|------|----------|--------------------|
| $0.56^{+0.17}_{-0.16} \pm 0.09$ | AAIJ | 13X LHCB | Repl. by AAIJ 15AQ |
|---------------------------------|------|----------|--------------------|

## $B_s^0\text{-}\overline{B}_s^0$ MIXING

For a discussion of  $B_s^0\text{-}\overline{B}_s^0$  mixing see the note on “ $B^0\text{-}\overline{B}^0$  Mixing” in the  $B^0$  Particle Listings above.

$\chi_s$  is a measure of the time-integrated  $B_s^0\text{-}\overline{B}_s^0$  mixing probability that produced  $B_s^0(\overline{B}_s^0)$  decays as a  $\overline{B}_s^0(B_s^0)$ . Mixing violates  $\Delta B \neq 2$  rule.

$$\chi_s = \frac{x_s^2}{2(1+x_s^2)}$$

$$x_s = \frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = (m_{B_{sH}^0} - m_{B_{sL}^0}) \tau_{B_s^0},$$

where  $H, L$  stand for heavy and light states of two  $B_s^0$   $CP$  eigenstates and

$$\tau_{B_s^0} = \frac{1}{0.5(\Gamma_{B_{sH}^0} + \Gamma_{B_{sL}^0})}.$$

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$$

$\Delta m_{B_s^0}$  is a measure of  $2\pi$  times the  $B_s^0\text{-}\overline{B}_s^0$  oscillation frequency in time-dependent mixing experiments.

“OUR EVALUATION” is provided by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) by taking into account correlations between measurements.

| VALUE ( $10^{12} \hbar s^{-1}$ )                                              | CL% | DOCUMENT ID                  | TECN      | COMMENT                     |
|-------------------------------------------------------------------------------|-----|------------------------------|-----------|-----------------------------|
| <b>17.749±0.020 OUR EVALUATION</b>                                            |     |                              |           |                             |
| <b>17.756±0.021 OUR AVERAGE</b>                                               |     |                              |           |                             |
| 17.703±0.059±0.018                                                            |     | <sup>1</sup> AAIJ            | 19Q LHCb  | $pp$ at 13 TeV              |
| 17.768±0.023±0.006                                                            |     | <sup>2</sup> AAIJ            | 13BI LHCb | $pp$ at 7 TeV               |
| 17.93 ±0.22 ±0.15                                                             |     | <sup>3</sup> AAIJ            | 13CF LHCb | $pp$ at 7 TeV               |
| 17.63 ±0.11 ±0.02                                                             |     | <sup>4</sup> AAIJ            | 12I LHCb  | $pp$ at 7 TeV               |
| 17.77 ±0.10 ±0.07                                                             |     | <sup>5</sup> ABULENCIA,A 06G | CDF       | $p\overline{p}$ at 1.96 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |                              |           |                             |
| 17.711 <sup>+0.055</sup> <sub>−0.057</sub> ±0.011                             |     | <sup>1</sup> AAIJ            | 15I LHCb  | Repl. by AAIJ 19Q           |
| 17–21                                                                         | 90  | <sup>6</sup> ABAZOV          | 06B D0    | $p\overline{p}$ at 1.96 TeV |
| 17.31 <sup>+0.33</sup> <sub>−0.18</sub> ±0.07                                 |     | <sup>7</sup> ABULENCIA       | 06Q CDF   | Repl. by ABULENCIA,A 06G    |
| > 8.0                                                                         | 95  | <sup>8</sup> ABDALLAH        | 04J DLPH  | $e^+e^- \rightarrow Z^0$    |
| > 4.9                                                                         | 95  | <sup>9</sup> ABDALLAH        | 04J DLPH  | $e^+e^- \rightarrow Z^0$    |
| > 8.5                                                                         | 95  | <sup>10</sup> ABDALLAH       | 04J DLPH  | $e^+e^- \rightarrow Z^0$    |
| > 5.0                                                                         | 95  | <sup>11</sup> ABDALLAH       | 03B DLPH  | $e^+e^- \rightarrow Z$      |
| >10.3                                                                         | 95  | <sup>12</sup> ABE            | 03 SLD    | $e^+e^- \rightarrow Z$      |
| >10.9                                                                         | 95  | <sup>13</sup> HEISTER        | 03E ALEP  | $e^+e^- \rightarrow Z$      |
| > 5.3                                                                         | 95  | <sup>14</sup> ABE            | 02V SLD   | $e^+e^- \rightarrow Z$      |
| > 1.0                                                                         | 95  | <sup>15</sup> ABBIENDI       | 01D OPAL  | $e^+e^- \rightarrow Z$      |
| > 7.4                                                                         | 95  | <sup>16</sup> ABREU          | 00Y DLPH  | Repl. by ABDALLAH 04J       |
| > 4.0                                                                         | 95  | <sup>17</sup> ABREU,P        | 00G DLPH  | $e^+e^- \rightarrow Z$      |

|       |    |    |            |     |      |                        |
|-------|----|----|------------|-----|------|------------------------|
| > 5.2 | 95 | 18 | ABBIENDI   | 99S | OPAL | $e^+e^- \rightarrow Z$ |
| <96   | 95 | 19 | ABE        | 99D | CDF  | $p\bar{p}$ at 1.8 TeV  |
| > 5.8 | 95 | 20 | ABE        | 99J | CDF  | $p\bar{p}$ at 1.8 TeV  |
| > 9.6 | 95 | 21 | BARATE     | 99J | ALEP | $e^+e^- \rightarrow Z$ |
| > 7.9 | 95 | 22 | BARATE     | 98C | ALEP | Repl. by BARATE 99J    |
| > 3.1 | 95 | 23 | ACKERSTAFF | 97U | OPAL | Repl. by ABBIENDI 99S  |
| > 2.2 | 95 | 24 | ACKERSTAFF | 97V | OPAL | Repl. by ABBIENDI 99S  |
| > 6.5 | 95 | 25 | ADAM       | 97  | DLPH | Repl. by ABREU 00Y     |
| > 6.6 | 95 | 26 | BUSKULIC   | 96M | ALEP | Repl. by BARATE 98C    |
| > 2.2 | 95 | 24 | AKERS      | 95J | OPAL | Sup. by ACKERSTAFF 97V |
| > 5.7 | 95 | 27 | BUSKULIC   | 95J | ALEP | $e^+e^- \rightarrow Z$ |
| > 1.8 | 95 | 24 | BUSKULIC   | 94B | ALEP | $e^+e^- \rightarrow Z$ |

<sup>1</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>2</sup> Measured using  $B_s^0 \rightarrow D_s^- \pi^+$  decays.

<sup>3</sup> Measured using  $B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu X$  decays.

<sup>4</sup> Measured using  $B_s^0 \rightarrow D_s^- \pi^+$  and  $D_s^- \pi^+ \pi^- \pi^+$  decays.

<sup>5</sup> Significance of oscillation signal is  $5.4 \sigma$ . Also reports  $|V_{td} / V_{ts}| = 0.2060 \pm 0.0007^{+0.0081}_{-0.0060}$ .

<sup>6</sup> A likelihood scan over the oscillation frequency,  $\Delta m_s$ , gives a most probable value of  $19 \text{ ps}^{-1}$  and a range of  $17 < \Delta m_s < 21 (\text{ps}^{-1})$  at 90% C.L. assuming Gaussian uncertainties. Also excludes  $\Delta m_s < 14.8 \text{ ps}^{-1}$  at 95% C.L.

<sup>7</sup> Significance of oscillation signal is 0.2%. Also reported the value  $|V_{td} / V_{ts}| = 0.208^{+0.001+0.008}_{-0.002-0.006}$ .

<sup>8</sup> Uses leptons emitted with large momentum transverse to a jet and improved techniques for vertexing and flavor-tagging.

<sup>9</sup> Updates of  $D_s$ -lepton analysis.

<sup>10</sup> Combined results from all Delphi analyses.

<sup>11</sup> Events with a high transverse momentum lepton were removed and an inclusively reconstructed vertex was required.

<sup>12</sup> ABE 03 uses the novel "charge dipole" technique to reconstruct separate secondary and tertiary vertices originating from the  $B \rightarrow D$  decay chain. The analysis excludes  $\Delta m_s < 4.9 \text{ ps}^{-1}$  and  $7.9 < \Delta m_s < 10.3 \text{ ps}^{-1}$ .

<sup>13</sup> Three analyses based on complementary event selections: (1) fully-reconstructed hadronic decays; (2) semileptonic decays with  $D_s$  exclusively reconstructed; (3) inclusive semileptonic decays.

<sup>14</sup> ABE 02V uses exclusively reconstructed  $D_s^-$  mesons and excludes  $\Delta m_s < 1.4 \text{ ps}^{-1}$  and  $2.4 < \Delta m_s < 5.3 \text{ ps}^{-1}$  at 95%CL.

<sup>15</sup> Uses fully or partially reconstructed  $D_s \ell$  vertices and a mixing tag as a flavor tagging.

<sup>16</sup> Replaced by ABDALLAH 04A. Uses  $D_s^- \ell^+$ , and  $\phi \ell^+$  vertices, and a multi-variable discriminant as a flavor tagging.

<sup>17</sup> Uses inclusive  $D_s$  vertices and fully reconstructed  $B_s$  decays and a multi-variable discriminant as a flavor tagging.

<sup>18</sup> Uses  $\ell$ - $Q_{\text{hem}}$  and  $\ell$ - $\ell$ .

<sup>19</sup> ABE 99D assumes  $\tau_{B_s^0} = 1.55 \pm 0.05 \text{ ps}$  and  $\Delta\Gamma/\Delta m = (5.6 \pm 2.6) \times 10^{-3}$ .

<sup>20</sup> ABE 99J uses  $\phi$   $\ell$ - $\ell$  correlation.

<sup>21</sup> BARATE 99J uses combination of an inclusive lepton and  $D_s^-$ -based analyses.

- <sup>22</sup> BARATE 98C combines results from  $D_s h\text{-}\ell/Q_{\text{hem}}$ ,  $D_s h\text{-}K$  in the same side,  $D_s \ell\text{-}\ell/Q_{\text{hem}}$  and  $D_s \ell\text{-}K$  in the same side.
- <sup>23</sup> Uses  $\ell\text{-}Q_{\text{hem}}$ .
- <sup>24</sup> Uses  $\ell\text{-}\ell$ .
- <sup>25</sup> ADAM 97 combines results from  $D_s \ell\text{-}Q_{\text{hem}}$ ,  $\ell\text{-}Q_{\text{hem}}$ , and  $\ell\text{-}\ell$ .
- <sup>26</sup> BUSKULIC 96M uses  $D_s$  lepton correlations and lepton, kaon, and jet charge tags.
- <sup>27</sup> BUSKULIC 95J uses  $\ell\text{-}Q_{\text{hem}}$ . They find  $\Delta m_s > 5.6$  [ $> 6.1$ ] for  $f_s=10\%$  [12%]. We interpolate to our central value  $f_s=10.5\%$ .

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0}$$

This is derived by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) from the results on  $\Delta m_{B_s^0}$  and “OUR EVALUATION” of the  $B_s^0$  mean lifetime.

VALUE  
**26.89±0.07 OUR EVALUATION**

DOCUMENT ID

$\chi_s$

This is a  $B_s^0\text{-}\bar{B}_s^0$  integrated mixing parameter derived from  $x_s$  above and OUR EVALUATION of  $\Delta \Gamma_{B_s^0} / \Gamma_{B_s^0}$ .

VALUE  
**0.499312±0.000004 OUR EVALUATION**

DOCUMENT ID

## CP VIOLATION PARAMETERS in $B_s^0$


$$\text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$$

CP impurity in  $B_s^0$  system.

“OUR EVALUATION” is an average obtained by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) and described at <https://hflav.web.cern.ch/>. It is the result of a fit to  $B_d$  and  $B_s$  CP asymmetries, which includes the  $B_s$  measurements listed below and the  $B_d$  measurements listed in the  $B_d$  section, and takes into account correlations between those measurements.

| <u>VALUE (units <math>10^{-3}</math>)</u>                                                                                                                                                                                                                                             | <u>DOCUMENT ID</u>                                          | <u>TECN</u> | <u>COMMENT</u>            |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------|-------------|---------------------------|
| <b>−0.15±0.70 OUR EVALUATION</b>                                                                                                                                                                                                                                                      |                                                             |             |                           |
| <b>0.0 ±1.1 OUR AVERAGE</b>                                                                                                                                                                                                                                                           | Error includes scale factor of 1.6. See the ideogram below. |             |                           |
| 0.98±0.65±0.5                                                                                                                                                                                                                                                                         | <sup>1</sup> AAIJ                                           | 16G         | LHCB $pp$ at 7, 8 TeV     |
| −2.15±1.85                                                                                                                                                                                                                                                                            | <sup>2</sup> ABAZOV                                         | 14          | D0 $p\bar{p}$ at 1.96 TeV |
| −2.8 ±1.9 ±0.4                                                                                                                                                                                                                                                                        | <sup>3</sup> ABAZOV                                         | 13          | D0 $p\bar{p}$ at 1.96 TeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●                                                                                                                                                                                                         |                                                             |             |                           |
| −0.15±1.25±0.90                                                                                                                                                                                                                                                                       | <sup>4</sup> AAIJ                                           | 14D         | LHCB Repl. by AAIJ 16G    |
| −4.5 ±2.7                                                                                                                                                                                                                                                                             | <sup>5</sup> ABAZOV                                         | 11U         | D0 Repl. by ABAZOV 14     |
| −0.4 ±2.3 ±0.4                                                                                                                                                                                                                                                                        | <sup>6</sup> ABAZOV                                         | 10E         | D0 Repl. by ABAZOV 13     |
| −3.6 ±1.9                                                                                                                                                                                                                                                                             | <sup>7</sup> ABAZOV                                         | 10H         | D0 Repl. by ABAZOV 11U    |
| 6.1 ±4.8 ±0.9                                                                                                                                                                                                                                                                         | <sup>8</sup> ABAZOV                                         | 07A         | D0 Repl. by ABAZOV 10E    |
| <sup>1</sup> AAIJ 16G reports a measurement of time-integrated flavor-specific asymmetry in $B_s^0 \rightarrow \mu^+ D_s^- X$ decays, $A_{SL}^s = (0.39 \pm 0.26 \pm 0.20)\%$ , which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 +  \epsilon_{B_s^0} ^2)$ . |                                                             |             |                           |

- <sup>2</sup> ABAZOV 14 uses the dimuon charge asymmetry with different impact parameters from which it reports  $A_{SL}^s = (-0.86 \pm 0.74) \times 10^{-2}$ .
- <sup>3</sup> ABAZOV 13 reports a measurement of time-integrated flavor-specific asymmetry in mixed semileptonic  $B_s^0 \rightarrow \mu^+ D_s^- X$  decays  $A_{SL}^s = (-1.12 \pm 0.74 \pm 0.17)\%$  which is approximately equal to  $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$ .
- <sup>4</sup> AAIJ 14D reports a measurement of time-integrated flavor-specific asymmetry in  $B_s^0 \rightarrow \mu^+ D_s^- X$  decays,  $A_{SL}^s = (-0.06 \pm 0.50 \pm 0.36)\%$ , which is approximately equal to  $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$ .
- <sup>5</sup> ABAZOV 11U uses the dimuon charge asymmetry with different impact parameters from which it reports  $A_{SL}^s = (-18.1 \pm 10.6) \times 10^{-3}$ .
- <sup>6</sup> ABAZOV 10E reports a measurement of flavor-specific asymmetry in  $B_{(s)}^0 \rightarrow \mu^+ D_{(s)}^{*-} X$  decays with a decay-time analysis including initial-state flavor tagging,  $A_{SL}^s = (-1.7 \pm 9.1_{-1.5}^{+1.4}) \times 10^{-3}$  which is approximately equal to  $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$ .
- <sup>7</sup> ABAZOV 10H reports a measurement of like-sign dimuon charge asymmetry of  $A_{SL}^s = (-9.57 \pm 2.51 \pm 1.46) \times 10^{-3}$  in semileptonic  $b$ -hadron decays. Using the measured production ratio of  $B_d^0$  and  $B_s^0$ , and the asymmetry of  $B_d^0$   $A_{SL}^s = (-4.7 \pm 4.6) \times 10^{-3}$  measured from  $B$ -factories, they obtain the asymmetry for  $B_s^0$ .
- <sup>8</sup> The first direct measurement of the time integrated flavor untagged charge asymmetry in semileptonic  $B_s^0$  decays is reported as  $2 \times A_{SL}^s(\text{untagged}) = A_{SL}^s = (2.45 \pm 1.93 \pm 0.35) \times 10^{-2}$ .



**$C_{KK}(B_s^0 \rightarrow K^+ K^-)$** 

| VALUE                                      | DOCUMENT ID | TECN      | COMMENT       |
|--------------------------------------------|-------------|-----------|---------------|
| <b><math>0.14 \pm 0.11 \pm 0.03</math></b> | AAIJ        | 13B0 LHCb | $pp$ at 7 TeV |

 **$S_{KK}(B_s^0 \rightarrow K^+ K^-)$** 

| VALUE                                      | DOCUMENT ID | TECN      | COMMENT       |
|--------------------------------------------|-------------|-----------|---------------|
| <b><math>0.30 \pm 0.12 \pm 0.04</math></b> | AAIJ        | 13B0 LHCb | $pp$ at 7 TeV |

 **$r_B(B_s^0 \rightarrow D_s^\mp K^\pm)$** 

$r_B$  and  $\delta_B$  are the amplitude ratio and relative strong phase between the amplitudes of  $A(B_s^0 \rightarrow D_s^+ K^-)$  and  $A(B_s^0 \rightarrow D_s^- K^+)$ ,

| VALUE                                    | DOCUMENT ID       | TECN     | COMMENT          |
|------------------------------------------|-------------------|----------|------------------|
| <b><math>0.37^{+0.10}_{-0.09}</math></b> | <sup>1</sup> AAIJ | 18U LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                        |                   |           |                   |
|------------------------|-------------------|-----------|-------------------|
| $0.53^{+0.17}_{-0.16}$ | <sup>2</sup> AAIJ | 14BF LHCb | Repl. by AAIJ 18U |
|------------------------|-------------------|-----------|-------------------|

<sup>1</sup> Measured in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays, constraining  $-2\beta_s$  by the measurement of  $\phi_s = -0.030 \pm 0.033$  from HFLAV.

<sup>2</sup> Measured in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays, constraining  $-2\beta_s$  by the measurement of  $\phi_s = 0.01 \pm 0.07 \pm 0.0$  from AAIJ 13AR. At 68% CL.

 **$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp)$** 

| VALUE (°)                           | DOCUMENT ID       | TECN     | COMMENT          |
|-------------------------------------|-------------------|----------|------------------|
| <b><math>358^{+13}_{-14}</math></b> | <sup>1</sup> AAIJ | 18U LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                 |                   |           |                   |
|-----------------|-------------------|-----------|-------------------|
| $3^{+19}_{-20}$ | <sup>2</sup> AAIJ | 14BF LHCb | Repl. by AAIJ 18U |
|-----------------|-------------------|-----------|-------------------|

<sup>1</sup> Measured in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays, constraining  $-2\beta_s$  by the measurement of  $\phi_s = 0.030 \pm 0.033$  from HFLAV. The value is modulo  $180^\circ$ .

<sup>2</sup> Measured in  $B_s^0 \rightarrow D_s^\mp K^\pm$  decays, constraining  $-2\beta_s$  by the measurement of  $\phi_s = 0.01 \pm 0.07 \pm 0.0$  from AAIJ 13AR. The value is modulo  $180^\circ$  at 68% CL.

**CP Violation phase  $\beta_s$** 

$-2\beta_s$  is the weak phase difference between  $B_s^0$  mixing amplitude and the  $B_s^0 \rightarrow J/\psi \phi$  decay amplitude driven by the  $b \rightarrow c \bar{c} s$  transition (such as  $B_s \rightarrow J/\psi \phi$ ,  $J/\psi K^+ K^-$ ,  $J/\psi \pi^+ \pi^-$ , and  $D_s^+ D_s^-$ ). The Standard Model value of  $\beta_s$  is  $\arg(-\frac{V_{ts} V_{tb}^*}{V_{cs} V_{cb}^*})$  if penguin contributions are neglected.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV, <https://hflav.web.cern.ch/>) and are described at <https://hflav.web.cern.ch/>. The averaging/scaling procedure takes into account correlation between the measurements.

| VALUE ( $10^{-2}$ rad)                           | DOCUMENT ID         | TECN      | COMMENT              |
|--------------------------------------------------|---------------------|-----------|----------------------|
| <b><math>2.55 \pm 1.15</math> OUR EVALUATION</b> |                     |           |                      |
| <b><math>4.1 \pm 1.6</math> OUR AVERAGE</b>      |                     |           |                      |
| $3.7 \pm 5.8 \pm 1.4$                            | <sup>1,2</sup> AAIJ | 19AP LHCb | $pp$ at 7, 8, 13 TeV |

|                                 |                          |           |                        |
|---------------------------------|--------------------------|-----------|------------------------|
| $4.15 \pm 2.05 \pm 0.30$        | <sup>3</sup> AAIJ        | 19Q LHCb  | $pp$ at 13 TeV         |
| $11.9 \pm 10.7 \pm 3.4$         | <sup>4</sup> AAIJ        | 17V LHCb  | $pp$ at 7, 8 TeV       |
| $4.5 \pm 3.9 \pm 2.1$           | <sup>5</sup> AAD         | 16AP ATLS | $pp$ at 7, 8 TeV       |
| $-11.5 \pm^{+14}_{-14.5} \pm 1$ | <sup>6</sup> AAIJ        | 16AK LHCb | $pp$ at 7, 8 TeV       |
| $3.75 \pm 4.85 \pm 1.55$        | <sup>7</sup> KHACHATRYAN | 16S CMS   | $pp$ at 8 TeV          |
| $-1 \pm 9 \pm 1$                | <sup>8</sup> AAIJ        | 14AY LHCb | $pp$ at 7, 8 TeV       |
|                                 | <sup>9</sup> AALTONEN    | 12AJ CDF  | $p\bar{p}$ at 1.96 TeV |
| $28 \pm^{+18}_{-19}$            | <sup>10</sup> ABAZOV     | 12D D0    | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                                |                         |           |                        |
|------------------------------------------------|-------------------------|-----------|------------------------|
| $-0.1 \pm 2.2 \pm 0.6$                         | <sup>11</sup> AAIJ      | 19AF LHCb | $pp$ at 7, 8, 13 TeV   |
| $5.0 \pm 6.5 \pm 7.0$                          | <sup>12</sup> AAIJ      | 18S LHCb  | $pp$ at 7, 8 TeV       |
| $2.9 \pm 2.5 \pm 0.3$                          | <sup>13</sup> AAIJ      | 15I LHCb  | Repl. by AAIJ 19Q      |
| $6 \pm^{+8}_{-7}$                              | <sup>14,15</sup> AAIJ   | 15K LHCb  | $pp$ at 7, 8 TeV       |
| $-6 \pm 13 \pm 3$                              | <sup>16</sup> AAD       | 14U ATLS  | Repl. by AAD 16AP      |
| $8.5 \pm 7.5 \pm 1.5$                          | <sup>17</sup> AAIJ      | 14AE LHCb | Repl. by AAIJ 19AP     |
| $-3.5 \pm 3.4 \pm 0.4$                         | <sup>18</sup> AAIJ      | 14S LHCb  | Repl. by AAIJ 19AF     |
| $-0.5 \pm 3.5 \pm 0.5$                         | <sup>19</sup> AAIJ      | 13AR LHCb | Repl. by AAIJ 15I      |
|                                                | <sup>20</sup> AAIJ      | 13AY LHCb | $pp$ at 7 TeV          |
| $-11.0 \pm 20.5 \pm 5.0$                       | <sup>21</sup> AAD       | 12CV ATLS | Repl. by AAD 14U       |
| $22 \pm 22 \pm 1$                              | <sup>22</sup> AAIJ      | 12B LHCb  | Repl. by AAIJ 12Q      |
| $-8 \pm 9 \pm 3$                               | <sup>23</sup> AAIJ      | 12D LHCb  | Repl. by AAIJ 13AR     |
| $0.95 \pm^{+8.70}_{-8.65} \pm^{+0.15}_{-0.20}$ | <sup>24</sup> AAIJ      | 12Q LHCb  | Repl. by AAIJ 13AR     |
|                                                | <sup>25</sup> AALTONEN  | 12D CDF   | Repl. by AALTONEN 12AJ |
|                                                | <sup>26</sup> AALTONEN  | 08G CDF   | Repl. by AALTONEN 12D  |
| $28 \pm^{+12}_{-15} \pm^{+4}_{-1}$             | <sup>10,27</sup> ABAZOV | 08AMD0    | Repl. by ABAZOV 12D    |
| $39.5 \pm 28.0 \pm^{+0.5}_{-7.0}$              | <sup>28,29</sup> ABAZOV | 07 D0     | Repl. by ABAZOV 07N    |
| $35 \pm^{+20}_{-24}$                           | <sup>29,30</sup> ABAZOV | 07N D0    | Repl. by ABAZOV 08AM   |

<sup>1</sup> AAIJ 19AP reports  $\phi_s^{s\bar{s}s} = -0.073 \pm 0.115 \pm 0.027$  rad and  $|\lambda| = 0.99 \pm 0.05 \pm 0.01$ .

Measured using a time-dependent fit to  $B_s^0 \rightarrow \phi\phi$  decays, assuming independence of the helicity of the  $\phi\phi$  decay.

<sup>2</sup> AAIJ 19AP reports also polarisation-dependent results assuming that the longitudinal weak phase is CP-conserving and that there is no direct CP violation, giving  $\phi_{s,\parallel} = 0.014 \pm 0.055 \pm 0.011$  rad and  $\phi_{s,\perp} = 0.044 \pm 0.059 \pm 0.019$  rad.

<sup>3</sup> AAIJ 19Q reports  $\phi_s = -2\beta_s = -0.083 \pm 0.041 \pm 0.006$  rad. that was measured using a time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>4</sup> Measured using time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi K^+ K^-$  in the region  $m(KK) > 1.05$  GeV.

<sup>5</sup> AAD 16AP reports  $\phi_s = -2\beta_s = -0.090 \pm 0.078 \pm 0.041$  rad. that was measured using a time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays.

<sup>6</sup> AAIJ 16AK reports  $\phi_s = -2\beta_s = 0.23 \pm^{+0.29}_{-0.28} \pm 0.02$  rad. that was measured using a time-dependent angular analysis of  $B_s^0 \rightarrow \psi(2S)\phi$  decays.

<sup>7</sup> KHACHATRYAN 16S reports  $\phi_s = -2\beta_s = -0.075 \pm 0.097 \pm 0.031$  rad. that was measured using a time-dependent angular analysis of  $B_s^0 \rightarrow J/\psi\phi$  decays.



- <sup>8</sup> AAIJ 14AY reports  $\phi_S = -2\beta_S = 0.02 \pm 0.17 \pm 0.02$  rad. in a time-dependent fit to  $B_S^0 \rightarrow D_S^+ D_S^-$ , while allowing  $CP$  violation in decay.
- <sup>9</sup> AALTONEN 12AJ reports  $-\pi/2 < \beta_S < -1.51$  or  $-0.06 < \beta_S < 0.30$ , or  $1.26 < \beta_S < \pi/2$  rad. at 68% CL. Measured using the time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi \phi$  decays.
- <sup>10</sup> ABAZOV 12D reports  $\phi_S = -2\beta_S = -0.55^{+0.38}_{-0.36}$  rad. that was measured using a time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi \phi$  decays. A single error includes both statistical and systematic uncertainties.
- <sup>11</sup> AAIJ 19AF reports  $\phi_S = -2\beta_S = 0.002 \pm 0.044 \pm 0.012$  rad. and  $|\lambda| = 0.949 \pm 0.036 \pm 0.019$ , when direct  $CP$  violation is allowed. Measured using a time-dependent fit to  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  decays, which is sensitive to  $\phi_S(s\bar{s}s)$ , not  $\phi_S(c\bar{c}s)$ .
- <sup>12</sup> AAIJ 18S reports  $\phi_S = -2\beta_S = -0.10 \pm 0.13 \pm 0.14$  rad measured in  $B_S^0 \rightarrow (K^+ \pi^-)(K^- \pi^+)$  in the region  $0.75 < m(K^\pm \pi^\mp) < 1.6$  GeV. This is a  $b \rightarrow d\bar{d}s$  transition with a decay amplitude phase different from that of  $b \rightarrow c\bar{c}s$  transition.
- <sup>13</sup> AAIJ 15I reports  $\phi_S = -2\beta_S = -0.058 \pm 0.049 \pm 0.006$  rad. that was measured using a time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi K^+ K^-$  decays. It also combines this result with that of AAIJ 14S and quotes  $\phi_S = -2\beta_S = -0.010 \pm 0.039$  rad.
- <sup>14</sup> AAIJ 15K reports  $-2\beta_S = -0.12^{+0.14}_{-0.16}$  rad. The value was obtained by measuring time-dependent  $CP$  asymmetry in  $B_S^0 \rightarrow K^+ K^-$  and using a U-spin relation between  $B_S^0 \rightarrow K^+ K^-$  and  $B^0 \rightarrow \pi^+ \pi^-$ .
- <sup>15</sup> Results are also presented using additional inputs on  $B^0 \rightarrow \pi^0 \pi^0$  and  $B^+ \rightarrow \pi^+ \pi^0$  decays from other experiments and isospin symmetry assumptions. The dependence of the results on the maximum allowed amount of U-spin breaking up to 50% is also included.
- <sup>16</sup> AAD 14U reports  $\phi_S = -2\beta_S = 0.12 \pm 0.25 \pm 0.05$  rad. that was measured using a time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi \phi$  decays.
- <sup>17</sup> AAIJ 14AE value measured in  $B_S^0 \rightarrow \phi \phi$  decays. This is a  $b \rightarrow s\bar{s}s$  transition with a decay amplitude phase different from that of  $b \rightarrow c\bar{c}s$  transition. Also reports  $\phi_S = -0.17 \pm 0.15 \pm 0.03$  rad.
- <sup>18</sup> AAIJ 14S reports  $\phi_S = -2\beta_S = 0.070 \pm 0.068 \pm 0.008$  rad. and  $|\lambda| = 0.89 \pm 0.05 \pm 0.01$ , when direct  $CP$  violation is allowed. Measured using a time-dependent fit to  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  decays.
- <sup>19</sup> AAIJ 13AR reports  $\phi_S = -2\beta_S = 0.01 \pm 0.07 \pm 0.01$  rad. obtained from combined fit to  $B_S^0 \rightarrow J/\psi K^+ K^-$  and  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  data sets. Also reports separate results of  $\phi_S = 0.07 \pm 0.09 \pm 0.01$  rad. from  $B_S^0 \rightarrow J/\psi K^+ K^-$  decays and  $\phi_S = -0.14^{+0.17}_{-0.16} \pm 0.01$  rad. from  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  decays.
- <sup>20</sup> AAIJ 13AY uses  $B_S^0 \rightarrow \phi \phi$  mode, and reports the 68% CL interval of  $\phi_S = -2\beta_S$  as  $[-2.46, -0.76]$  rad.
- <sup>21</sup> AAD 12CV reports  $\phi_S = -2\beta_S = 0.22 \pm 0.41 \pm 0.10$  rad. that was measured using a time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi \phi$  decays.
- <sup>22</sup> Reports  $\phi_S = -2\beta_S = -0.44 \pm 0.44 \pm 0.02$  rad. that was measured using a time-dependent fit to  $B_S^0 \rightarrow J/\psi f_0(980)$  decays.
- <sup>23</sup> Reports  $\phi_S = -2\beta_S = 0.15 \pm 0.18 \pm 0.06$  rad. that was measured using a time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi \phi$  decays.
- <sup>24</sup> Reports  $\phi_S = -2\beta_S = -0.019^{+0.173+0.004}_{-0.174-0.003}$  rad. which was measured using a time-dependent fit to  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  decays, with the  $\pi^+ \pi^-$  mass within 775–1550

MeV. Searches for, but finds no evidence, for direct  $CP$  violation in  $B_S^0 \rightarrow J/\psi \pi \pi$  decays.

<sup>25</sup> Reports  $0.02 < \phi_S < 0.52$  or  $1.08 < \phi_S < 1.55$  rad. at 68% C.L. confidence regions in the two-dimensional space of  $\phi_S$  and  $\Delta\Gamma_{B_S^0}$  from  $B_S^0 \rightarrow J/\psi \phi$  decays.

<sup>26</sup> Reports  $0.32 < 2\beta_S < 2.82$  rad. at 68% C.L. and confidence regions in the two-dimensional space of  $2\beta_S$  and  $\Delta\Gamma$  from the first measurement of  $B_S^0 \rightarrow J/\psi \phi$  decays using flavor tagging. The probability of a deviation from SM prediction as large as the level of observed data is 15%.

<sup>27</sup> Reports  $\phi_S = -2\beta_S$  and obtains 90% CL interval  $-0.03 < \beta_S < 0.60$  rad.

<sup>28</sup> The first direct measurement of the  $CP$ -violating mixing phase is reported from the time-dependent analysis of flavor untagged  $B_S^0 \rightarrow J/\psi \phi$  decays.

<sup>29</sup> Reports  $\phi_S$  which equals to  $-2\beta_S$ .

<sup>30</sup> Combines D0 collaboration measurements of time-dependent angular distributions in  $B_S^0 \rightarrow J/\psi \phi$  and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

### $|\lambda| (B_S^0 \rightarrow J/\psi(1S)\phi)$

| VALUE                                                                         | DOCUMENT ID | TECN | COMMENT                |
|-------------------------------------------------------------------------------|-------------|------|------------------------|
| <b><math>1.012 \pm 0.016 \pm 0.006</math></b>                                 | AAIJ        | 19Q  | LHCB $pp$ at 13 TeV    |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |                        |
| $0.964 \pm 0.019 \pm 0.007$                                                   | AAIJ        | 15I  | LHCB Repl. by AAIJ 19Q |

### $|\lambda|$

| VALUE                                                                         | DOCUMENT ID       | TECN | COMMENT                   |
|-------------------------------------------------------------------------------|-------------------|------|---------------------------|
| <b><math>0.999 \pm 0.017</math> OUR AVERAGE</b>                               |                   |      |                           |
| $0.99 \pm 0.05 \pm 0.01$                                                      | <sup>1</sup> AAIJ | 19AP | LHCB $pp$ at 7, 8, 13 TeV |
| $1.035 \pm 0.034 \pm 0.089$                                                   | <sup>2</sup> AAIJ | 18S  | LHCB $pp$ at 7, 8 TeV     |
| $0.994 \pm 0.018 \pm 0.006$                                                   | <sup>3</sup> AAIJ | 17V  | LHCB $pp$ at 7, 8 TeV     |
| $1.045^{+0.069}_{-0.050} \pm 0.007$                                           | <sup>4</sup> AAIJ | 16AK | LHCB $pp$ at 7, 8 TeV     |
| $0.91^{+0.18}_{-0.15} \pm 0.02$                                               | <sup>5</sup> AAIJ | 14AY | LHCB $pp$ at 7, 8 TeV     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                   |      |                           |
| $0.949 \pm 0.036 \pm 0.019$                                                   | <sup>6</sup> AAIJ | 19AF | LHCB $pp$ at 7, 8, 13 TeV |
| $1.04 \pm 0.07 \pm 0.03$                                                      | <sup>7</sup> AAIJ | 14AE | LHCB Repl. by AAIJ 19AP   |

<sup>1</sup> Measured in  $B_S^0 \rightarrow \phi \phi$  decays.

<sup>2</sup> Measured in  $B_S^0 \rightarrow (K^+ \pi^-)(K^- \pi^+)$  in the region  $0.75 < m(K^\pm \pi^\mp) < 1.6$  GeV.

<sup>3</sup> Measured using time-dependent angular analysis of  $B_S^0 \rightarrow J/\psi K^+ K^-$  in the region  $m(K K) > 1.05$  GeV.

<sup>4</sup> Measured using time-dependent angular analysis of  $B_S^0 \rightarrow \psi(2S)\phi$  decays.

<sup>5</sup> Measured in  $B_S^0 \rightarrow D_S^+ D_S^-$  decays.

<sup>6</sup> Measured using time-dependent analysis of  $B_S^0 \rightarrow J/\psi \pi^+ \pi^-$  decays.

<sup>7</sup> Measured in  $B_S^0 \rightarrow \phi \phi$  decays.

**A, CP violation parameter**

$$A = -2 \operatorname{Re}(\lambda) / (1 + |\lambda|^2)$$

| VALUE                                          | DOCUMENT ID       | TECN      | COMMENT          |
|------------------------------------------------|-------------------|-----------|------------------|
| <b><math>-0.75 \pm 0.12</math> OUR AVERAGE</b> |                   |           |                  |
| $-0.79 \pm 0.07 \pm 0.10$                      | <sup>1</sup> AAIJ | 18O LHCb  | $pp$ at 7, 8 TeV |
| $0.49^{+0.77}_{-0.65} \pm 0.06$                | <sup>2</sup> AAIJ | 15AL LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow K^+ K^-$  decays.

<sup>2</sup> Measured in  $B_s^0 \rightarrow J/\psi K_S^0$  decays.

**C, CP violation parameter**

$$C = (1 - |\lambda|^2) / (1 + |\lambda|^2)$$

| VALUE                                         | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------------|-------------------|-----------|------------------|
| <b><math>0.19 \pm 0.06</math> OUR AVERAGE</b> |                   |           |                  |
| $0.20 \pm 0.06 \pm 0.02$                      | <sup>1</sup> AAIJ | 18O LHCb  | $pp$ at 7, 8 TeV |
| $-0.28 \pm 0.41 \pm 0.08$                     | <sup>2</sup> AAIJ | 15AL LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow K^+ K^-$  decays.

<sup>2</sup> Measured in  $B_s^0 \rightarrow J/\psi K_S^0$  decays.

**S, CP violation parameter**

$$S = -2 \operatorname{Im}(\lambda) / (1 + |\lambda|^2)$$

| VALUE                                         | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------------|-------------------|-----------|------------------|
| <b><math>0.17 \pm 0.06</math> OUR AVERAGE</b> |                   |           |                  |
| $0.18 \pm 0.06 \pm 0.02$                      | <sup>1</sup> AAIJ | 18O LHCb  | $pp$ at 7, 8 TeV |
| $-0.08 \pm 0.40 \pm 0.08$                     | <sup>2</sup> AAIJ | 15AL LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow K^+ K^-$  decays.

<sup>2</sup> Measured in  $B_s^0 \rightarrow J/\psi K_S^0$  decays.

 **$A_{CP}^L(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$** 

| VALUE                                          | DOCUMENT ID | TECN      | COMMENT          |
|------------------------------------------------|-------------|-----------|------------------|
| <b><math>-0.048 \pm 0.057 \pm 0.020</math></b> | AAIJ        | 15AV LHCb | $pp$ at 7, 8 TeV |

 **$A_{CP}^{\parallel}(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$** 

| VALUE                                         | DOCUMENT ID | TECN      | COMMENT          |
|-----------------------------------------------|-------------|-----------|------------------|
| <b><math>0.171 \pm 0.152 \pm 0.028</math></b> | AAIJ        | 15AV LHCb | $pp$ at 7, 8 TeV |

 **$A_{CP}^{\perp}(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$** 

| VALUE                                          | DOCUMENT ID | TECN      | COMMENT          |
|------------------------------------------------|-------------|-----------|------------------|
| <b><math>-0.049 \pm 0.096 \pm 0.025</math></b> | AAIJ        | 15AV LHCb | $pp$ at 7, 8 TeV |

 **$A_{CP}(B_s \rightarrow \pi^+ K^-)$** 

$A_{CP}$  is defined as

$$\frac{B(\bar{B}_s^0 \rightarrow f) - B(B_s^0 \rightarrow \bar{f})}{B(\bar{B}_s^0 \rightarrow f) + B(B_s^0 \rightarrow \bar{f})},$$

the  $CP$ -violation asymmetry of exclusive  $B_s^0$  and  $\bar{B}_s^0$  decay.

| VALUE                                                                         | DOCUMENT ID | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-------------|-----------|------------------------|
| <b><math>0.221 \pm 0.015</math> OUR AVERAGE</b>                               |             |           |                        |
| $0.213 \pm 0.015 \pm 0.007$                                                   | AAIJ        | 18O LHCb  | $pp$ at 7, 8 TeV       |
| $0.22 \pm 0.07 \pm 0.02$                                                      | AALTONEN    | 14P CDF   | $p\bar{p}$ at 1.96 TeV |
| $0.27 \pm 0.04 \pm 0.01$                                                      | AAIJ        | 13AX LHCb | $pp$ at 7 TeV          |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |           |                        |
| $0.27 \pm 0.08 \pm 0.02$                                                      | AAIJ        | 12V LHCb  | Repl. by AAIJ 13AX     |
| $0.39 \pm 0.15 \pm 0.08$                                                      | AALTONEN    | 11N CDF   | Repl. by AALTONEN 14P  |

### $A_{CP}(B_s^0 \rightarrow [K^+ K^-]_D \bar{K}^*(892)^0)$

| VALUE                                                                         | DOCUMENT ID | TECN      | COMMENT            |
|-------------------------------------------------------------------------------|-------------|-----------|--------------------|
| <b><math>-0.04 \pm 0.07 \pm 0.02</math></b>                                   | AAIJ        | 14BN LHCb | $pp$ at 7, 8 TeV   |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |           |                    |
| $0.04 \pm 0.16 \pm 0.01$                                                      | AAIJ        | 13L LHCb  | Repl. by AAIJ 14BN |

### $A_{CP}(B_s^0 \rightarrow [\pi^+ K^-]_D K^*(892)^0)$

| VALUE                                       | DOCUMENT ID | TECN      | COMMENT          |
|---------------------------------------------|-------------|-----------|------------------|
| <b><math>-0.01 \pm 0.03 \pm 0.02</math></b> | AAIJ        | 14BN LHCb | $pp$ at 7, 8 TeV |

### $A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0)$

| VALUE                                      | DOCUMENT ID | TECN      | COMMENT          |
|--------------------------------------------|-------------|-----------|------------------|
| <b><math>0.06 \pm 0.13 \pm 0.02</math></b> | AAIJ        | 14BN LHCb | $pp$ at 7, 8 TeV |

### $S(B_s^0 \rightarrow \phi \gamma)$

| VALUE                                      | DOCUMENT ID       | TECN      | COMMENT          |
|--------------------------------------------|-------------------|-----------|------------------|
| <b><math>0.43 \pm 0.30 \pm 0.11</math></b> | <sup>1</sup> AAIJ | 19AE LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in flavor tagged time dependent analysis.

### $C(B_s^0 \rightarrow \phi \gamma)$

| VALUE                                      | DOCUMENT ID       | TECN      | COMMENT          |
|--------------------------------------------|-------------------|-----------|------------------|
| <b><math>0.11 \pm 0.29 \pm 0.11</math></b> | <sup>1</sup> AAIJ | 19AE LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> Measured in flavor tagged time dependent analysis.

### $A^\Delta(B_s \rightarrow \phi \gamma)$

$A^\Delta(B_s \rightarrow \phi \gamma)$  is the multiplicative coefficient of the  $\sinh(\Delta\Gamma t/2)$  term in the  $B_s \rightarrow \phi \gamma$  decay rate time dependence.

| VALUE                                              | DOCUMENT ID       | TECN      | COMMENT          |
|----------------------------------------------------|-------------------|-----------|------------------|
| <b><math>-0.67^{+0.37}_{-0.41} \pm 0.17</math></b> | <sup>1</sup> AAIJ | 19AE LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                   |                   |          |                    |
|-----------------------------------|-------------------|----------|--------------------|
| $-0.98^{+0.46+0.23}_{-0.52-0.20}$ | <sup>2</sup> AAIJ | 17B LHCb | Repl. by AAIJ 19AE |
|-----------------------------------|-------------------|----------|--------------------|

<sup>1</sup> Measured in flavor tagged time dependent analysis, using tagged and un-tagged events. This result updates AAIJ 17B with better selection efficiency and other analysis improvements.

<sup>2</sup> Measured in time dependent analysis without initial flavor tagging.

## CPT VIOLATION PARAMETERS

In the  $B_s^0$  mixing, propagating mass eigenstates can be written as

$$\begin{aligned} |B_{sL}\rangle &\propto p \sqrt{1-\xi} |B_s^0\rangle + q \sqrt{1+\xi} |\bar{B}_s^0\rangle \\ |B_{sH}\rangle &\propto p \sqrt{1+\xi} |B_s^0\rangle - q \sqrt{1-\xi} |\bar{B}_s^0\rangle \end{aligned}$$

where parameter  $\xi$  controls  $CPT$  violation. If  $\xi$  is zero, then  $CPT$  is conserved. The parameter  $\xi$  can be written as

$$\xi = \frac{2(M_{11}-M_{22})-i(\Gamma_{11}-\Gamma_{22})}{-2\Delta m_s+i\Delta\Gamma_s} \approx \frac{-2\beta^\mu \Delta a_\mu}{2\Delta m_s-i\Delta\Gamma_s},$$

where  $M_{ii}$ ,  $\Gamma_{ii}$ ,  $\Delta m_s$ , and  $\Delta\Gamma_s$  are parameters of Hamiltonian governing  $B_s$  oscillations,  $\beta^\mu$  is the  $B_s^0$  meson velocity and  $\Delta a_\mu$  characterizes Lorentz-invariance violation.

### $\Delta a_\perp$

| VALUE ( $10^{-12}$ GeV)                     | CL% | DOCUMENT ID         | TECN | COMMENT                   |
|---------------------------------------------|-----|---------------------|------|---------------------------|
| <b><math>-0.47 \pm 0.39 \pm 0.08</math></b> |     | <sup>1</sup> AAIJ   | 16E  | LHCB $pp$ at 7, 8 TeV     |
| <b><math>&lt; 1.2</math></b>                | 95  | <sup>2</sup> ABAZOV | 15L  | D0 $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

<sup>2</sup> Measured in semileptonic  $B_s^0 \rightarrow D_s^- \mu^+ X$  decays. Also extracts limit on time and longitudinal components (  $-0.8 < \Delta a_T - 0.396 \Delta a_Z < 3.9$  )  $10^{-13}$  GeV.

### $\Delta a_\parallel$

| VALUE ( $10^{-14}$ GeV)                     | DOCUMENT ID       | TECN | COMMENT               |
|---------------------------------------------|-------------------|------|-----------------------|
| <b><math>-0.89 \pm 1.41 \pm 0.36</math></b> | <sup>1</sup> AAIJ | 16E  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

### $\Delta a_X$

| VALUE ( $10^{-14}$ GeV)                     | DOCUMENT ID       | TECN | COMMENT               |
|---------------------------------------------|-------------------|------|-----------------------|
| <b><math>+1.01 \pm 2.08 \pm 0.71</math></b> | <sup>1</sup> AAIJ | 16E  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

### $\Delta a_Y$

| VALUE ( $10^{-14}$ GeV)                     | DOCUMENT ID       | TECN | COMMENT               |
|---------------------------------------------|-------------------|------|-----------------------|
| <b><math>-3.83 \pm 2.09 \pm 0.71</math></b> | <sup>1</sup> AAIJ | 16E  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

### $\text{Re}(\xi)$

| VALUE                                          | DOCUMENT ID       | TECN | COMMENT               |
|------------------------------------------------|-------------------|------|-----------------------|
| <b><math>-0.022 \pm 0.033 \pm 0.003</math></b> | <sup>1</sup> AAIJ | 16E  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

### $\text{Im}(\xi)$

| VALUE                                         | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------------------------------|-------------------|------|-----------------------|
| <b><math>0.004 \pm 0.011 \pm 0.002</math></b> | <sup>1</sup> AAIJ | 16E  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> Uses  $B_s^0 \rightarrow J/\psi K^+ K^-$  decays.

**PARTIAL BRANCHING FRACTIONS IN  $B_s \rightarrow \phi \ell^+ \ell^-$**  **$B(B_s \rightarrow \phi \ell^+ \ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$** 

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID        | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|--------------------|-----------|------------------------|
| <b><math>1.14 \pm 0.16</math></b>                                             | <b>OUR AVERAGE</b> |           |                        |
| $1.11^{+0.14}_{-0.13} \pm 0.09$                                               | <sup>1</sup> AAIJ  | 15AQ LHCB | $pp$ at 7, 8 TeV       |
| $2.78 \pm 0.95 \pm 0.89$                                                      | AALTONEN           | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |           |                        |
| $0.897^{+0.207}_{-0.186} \pm 0.097$                                           | <sup>1</sup> AAIJ  | 13X LHCB  | Repl. by AAIJ 15AQ     |

<sup>1</sup> Measured in  $B_s^0 \rightarrow \phi \mu^+ \mu^-$  decays. **$B(B_s \rightarrow \phi \ell^+ \ell^-) (2.0 < q^2 < 5.0 \text{ GeV}^2/c^4)$** 

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>0.77 \pm 0.12 \pm 0.06</math></b>                                    | <sup>1</sup> AAIJ     | 15AQ LHCB | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| $0.529^{+0.182}_{-0.159} \pm 0.057$                                           | <sup>1,2</sup> AAIJ   | 13X LHCB  | Repl. by AAIJ 15AQ     |
| $0.58 \pm 0.55 \pm 0.19$                                                      | <sup>2</sup> AALTONEN | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow \phi \mu^+ \mu^-$  decays.<sup>2</sup> Measured in  $2 < q^2 < 4.3 \text{ GeV}^2/c^4$ . **$B(B_s \rightarrow \phi \ell^+ \ell^-) (5.0 < q^2 < 8.0 \text{ GeV}^2/c^4)$** 

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>0.96 \pm 0.13 \pm 0.08</math></b>                                    | <sup>1</sup> AAIJ     | 15AQ LHCB | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| $1.38^{+0.25}_{-0.23} \pm 0.14$                                               | <sup>1,2</sup> AAIJ   | 13X LHCB  | Repl. by AAIJ 15AQ     |
| $1.34 \pm 0.83 \pm 0.43$                                                      | <sup>2</sup> AALTONEN | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow \phi \mu^+ \mu^-$  decays.<sup>2</sup> Measured in  $4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$ . **$B(B_s \rightarrow \phi \ell^+ \ell^-) (11.0 < q^2 < 12.5 \text{ GeV}^2/c^4)$** 

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>0.71 \pm 0.10 \pm 0.06</math></b>                                    | <sup>1</sup> AAIJ     | 15AQ LHCB | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| $1.18^{+0.22}_{-0.21} \pm 0.14$                                               | <sup>1,2</sup> AAIJ   | 13X LHCB  | Repl. by AAIJ 15AQ     |
| $2.98 \pm 0.95 \pm 0.95$                                                      | <sup>2</sup> AALTONEN | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |

<sup>1</sup> Measured in  $B_s^0 \rightarrow \phi \mu^+ \mu^-$  decays.<sup>2</sup> Measured in  $10.9 < q^2 < 12.86 \text{ GeV}^2/c^4$ .

**$B(B_s \rightarrow \phi \ell^+ \ell^-)$  ( $15.0 < q^2 < 17.0 \text{ GeV}^2/c^4$ )**

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>0.90 \pm 0.11 \pm 0.07</math></b>                                    | <sup>1</sup> AAIJ     | 15AQ LHCb | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| $0.760^{+0.189}_{-0.169} \pm 0.087$                                           | <sup>1,2</sup> AAIJ   | 13X LHCb  | Repl. by AAIJ 15AQ     |
| $1.86 \pm 0.66 \pm 0.59$                                                      | <sup>2</sup> AALTONEN | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |
| <sup>1</sup> Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.         |                       |           |                        |
| <sup>2</sup> Measured in $14.18 < q^2 < 16 \text{ GeV}^2/c^4$ .               |                       |           |                        |

 **$B(B_s \rightarrow \phi \ell^+ \ell^-)$  ( $17.0 < q^2 < 19.0 \text{ GeV}^2/c^4$ )**

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID           | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-----------------------|-----------|------------------------|
| <b><math>0.79 \pm 0.11 \pm 0.07</math></b>                                    | <sup>1</sup> AAIJ     | 15AQ LHCb | $pp$ at 7, 8 TeV       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                       |           |                        |
| $1.06^{+0.23}_{-0.21} \pm 0.12$                                               | <sup>1,2</sup> AAIJ   | 13X LHCb  | Repl. by AAIJ 15AQ     |
| $2.32 \pm 0.76 \pm 0.74$                                                      | <sup>2</sup> AALTONEN | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |
| <sup>1</sup> Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.         |                       |           |                        |
| <sup>2</sup> Measured in $16 < q^2 < 19 \text{ GeV}^2/c^4$ .                  |                       |           |                        |

 **$B(B_s \rightarrow \phi \ell^+ \ell^-)$  ( $1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$ )**

| VALUE (units $10^{-7}$ )                                                      | DOCUMENT ID       | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-------------------|-----------|------------------------|
| <b><math>1.28 \pm 0.18</math> OUR AVERAGE</b>                                 |                   |           |                        |
| $1.29 \pm 0.16 \pm 0.10$                                                      | <sup>1</sup> AAIJ | 15AQ LHCb | $pp$ at 7, 8 TeV       |
| $1.14 \pm 0.79 \pm 0.36$                                                      | AALTONEN          | 11AI CDF  | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                   |           |                        |
| $1.14^{+0.25}_{-0.23} \pm 0.13$                                               | <sup>1</sup> AAIJ | 13X LHCb  | Repl. by AAIJ 15AQ     |
| <sup>1</sup> Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.         |                   |           |                        |

 **$B(B_s \rightarrow \phi \ell^+ \ell^-)$  ( $0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$ )**

| VALUE (units $10^{-7}$ )                   | DOCUMENT ID | TECN     | COMMENT                |
|--------------------------------------------|-------------|----------|------------------------|
| <b><math>3.30 \pm 1.09 \pm 1.05</math></b> | AALTONEN    | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

**PRODUCTION ASYMMETRIES** **$A_P(B_s^0)$** 

$$A_P(B_s^0) = [\sigma(\bar{B}_s^0) - \sigma(B_s^0)] / [\sigma(\bar{B}_s^0) + \sigma(B_s^0)]$$

| VALUE (units $10^{-2}$ )                                                                                                                                  | DOCUMENT ID       | TECN      | COMMENT                           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|-----------|-----------------------------------|
| <b><math>1.2 \pm 1.6</math> OUR AVERAGE</b>                                                                                                               |                   |           |                                   |
| $-0.65 \pm 2.88 \pm 0.59$                                                                                                                                 | <sup>1</sup> AAIJ | 17BF LHCb | $pp$ at 7 TeV                     |
| $1.98 \pm 1.90 \pm 0.59$                                                                                                                                  | <sup>1</sup> AAIJ | 17BF LHCb | $pp$ at 8 TeV                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                                             |                   |           |                                   |
| $1.09 \pm 2.61 \pm 0.66$                                                                                                                                  | <sup>2</sup> AAIJ | 14BP LHCb | Repl. by AAIJ 17BF, $pp$ at 7 TeV |
| <sup>1</sup> Based on time-dependent analysis of $B_s^0 \rightarrow D_s^- \pi^+$ in kinematic range $2 < p_T < 30 \text{ GeV}/c$ and $2.1 < \eta < 4.5$ . |                   |           |                                   |
| <sup>2</sup> Based on time-dependent analysis of $B_s^0 \rightarrow D_s^- \pi^+$ in kinematic range $4 < p_T < 30 \text{ GeV}/c$ and $2.5 < \eta < 4.5$ . |                   |           |                                   |

**$B_s^0$  REFERENCES**

|              |      |                |                              |                        |
|--------------|------|----------------|------------------------------|------------------------|
| AABOUD       | 19L  | JHEP 1904 098  | M. Aaboud <i>et al.</i>      | (ATLAS Collab.)        |
| AAIJ         | 19AE | PRL 123 081802 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19AF | PL B797 134789 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19AK | PRL 123 211801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19AP | JHEP 1912 155  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19K  | JHEP 1906 114  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19L  | JHEP 1907 032  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19Q  | EPJ C79 706    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 19U  | PRL 122 191804 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18AB | JHEP 1807 020  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18AC | JHEP 1808 191  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18AY | PR D98 071103  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18AZ | PR D98 072006  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18O  | PR D98 032004  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18S  | JHEP 1803 140  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18T  | JHEP 1803 078  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 18U  | JHEP 1803 059  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| SIRUNYAN     | 18BY | EPJ C78 457    | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)          |
| AAIJ         | 17A  | PR D95 012006  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17AI | PRL 118 191801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17AJ | PRL 118 251802 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17AL | PRL 119 041802 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17AN | PRL 119 101801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17B  | PRL 118 021801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BA | JHEP 1705 158  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BB | EPJ C77 609    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BD | PR D96 051103  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BF | PL B774 139    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BJ | PRL 119 232001 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17BP | JHEP 1711 027  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17G  | PRL 118 081801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17N  | JHEP 1703 001  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17U  | JHEP 1707 021  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 17V  | JHEP 1708 037  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AABOUD       | 16L  | EPJ C76 513    | M. Aaboud <i>et al.</i>      | (ATLAS Collab.)        |
| AAD          | 16AP | JHEP 1608 147  | G. Aad <i>et al.</i>         | (ATLAS Collab.)        |
| AAIJ         | 16   | JHEP 1601 012  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16AK | PL B762 253    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16AL | PL B762 484    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16C  | PRL 116 161802 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16E  | PRL 116 241601 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16G  | PRL 117 061803 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16P  | PR D93 092008  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 16U  | JHEP 1603 040  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| ABAZOV       | 16C  | PR D94 012001  | V.M. Abazov <i>et al.</i>    | (D0 Collab.)           |
| KHACHATRY... | 16Q  | PL B756 84     | V. Khachatryan <i>et al.</i> | (CMS Collab.)          |
| KHACHATRY... | 16S  | PL B757 97     | V. Khachatryan <i>et al.</i> | (CMS Collab.)          |
| PAL          | 16   | PRL 116 161801 | B. Pal <i>et al.</i>         | (BELLE Collab.)        |
| AAIJ         | 15AC | JHEP 1505 019  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AD | JHEP 1506 130  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AF | JHEP 1507 166  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AG | JHEP 1508 005  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AL | JHEP 1506 131  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AQ | JHEP 1509 179  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AS | JHEP 1510 053  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15AV | JHEP 1511 082  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15BB | PR D92 112002  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15D  | JHEP 1501 024  | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15I  | PRL 114 041801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15K  | PL B741 1      | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15O  | PRL 115 051801 | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15S  | PL B743 46     | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| AAIJ         | 15U  | PL B747 484    | R. Aaij <i>et al.</i>        | (LHCb Collab.)         |
| ABAZOV       | 15A  | PRL 114 062001 | V.M. Abazov <i>et al.</i>    | (D0 Collab.)           |
| ABAZOV       | 15L  | PRL 115 161601 | V.M. Abazov <i>et al.</i>    | (D0 Collab.)           |
| DUTTA        | 15   | PR D91 011101  | D. Dutta <i>et al.</i>       | (BELLE Collab.)        |
| KHACHATRY... | 15BE | NAT 522 68     | V. Khachatryan <i>et al.</i> | (CMS and LHCb Collab.) |
| OSWALD       | 15   | PR D92 072013  | C. Oswald <i>et al.</i>      | (BELLE Collab.)        |



|            |      |                        |                             |                 |
|------------|------|------------------------|-----------------------------|-----------------|
| AAD        | 14U  | PR D90 052007          | G. Aad <i>et al.</i>        | (ATLAS Collab.) |
| AAIJ       | 14AA | PRL 112 202001         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14AE | PR D90 052011          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14AX | PRL 113 172001         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14AY | PRL 113 211801         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BF | JHEP 1411 060          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BH | PR D90 072003          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BM | NJP 16 123001          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BN | PR D90 112002          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BP | PL B739 218            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14BR | PR D89 092006          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14D  | PL B728 607            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14E  | JHEP 1404 114          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14F  | PRL 112 111802         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14L  | JHEP 1407 140          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14R  | PL B736 446            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14S  | PL B736 186            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 14Y  | PRL 112 091802         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AALTONEN   | 14P  | PRL 113 242001         | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 14   | PR D89 012002          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| PDG        | 14   | CP C38 070001          | K. Olive <i>et al.</i>      | (PDG Collab.)   |
| AAIJ       | 13   | NP B867 1              | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13A  | NP B867 547            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AA | NP B871 403            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AB | NP B873 275            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AC | NP B874 663            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AL | PR D87 071101          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AN | PR D87 072004          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AP | PR D87 092007          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AQ | PR D87 112009          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AR | PR D87 112010          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AW | PRL 110 211801         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AX | PRL 110 221601         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13AY | PRL 110 241802         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13B  | PRL 110 021801         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BA | PRL 111 101805         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BI | NJP 15 053021          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BM | PRL 111 141801         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BO | JHEP 1310 183          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BP | JHEP 1310 143          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BQ | JHEP 1310 005          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BW | JHEP 1311 092          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13BX | PL B727 403            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13CF | EPJ C73 2655           | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13L  | JHEP 1303 067          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13X  | JHEP 1307 084          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 13Z  | JHEP 1309 006          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AALTONEN   | 13F  | PR D87 072003          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 13   | PRL 110 011801         | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 13C  | PR D87 072006          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| CHATRCHYAN | 13AW | PRL 111 101804         | S. Chatrchyan <i>et al.</i> | (CMS Collab.)   |
| ESEN       | 13   | PR D87 031101          | S. Esen <i>et al.</i>       | (BELLE Collab.) |
| OSWALD     | 13   | PR D87 072008          | C. Oswald <i>et al.</i>     | (BELLE Collab.) |
| Also       |      | PR D90 119901 (errat.) | C. Oswald <i>et al.</i>     | (BELLE Collab.) |
| SOLOVIEVA  | 13   | PL B726 206            | E. Solovieva <i>et al.</i>  | (BELLE Collab.) |
| THORNE     | 13   | PR D88 114006          | F. Thorne <i>et al.</i>     | (BELLE Collab.) |
| AAD        | 12AE | PL B713 387            | G. Aad <i>et al.</i>        | (ATLAS Collab.) |
| AAD        | 12CV | JHEP 1212 072          | G. Aad <i>et al.</i>        | (ATLAS Collab.) |
| AAIJ       | 12   | PL B707 349            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12A  | PL B708 55             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AE | PR D85 112013          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AG | JHEP 1206 115          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AM | PRL 109 131801         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AN | PRL 109 152002         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AO | PR D86 052006          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AP | PR D86 071102          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AR | JHEP 1210 037          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12AX | PR D86 112005          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12B  | PL B707 497            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12D  | PRL 108 101803         | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12E  | PL B708 241            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |

|            |      |                         |                             |                 |
|------------|------|-------------------------|-----------------------------|-----------------|
| AAIJ       | 12F  | PL B709 50              | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12I  | PL B709 177             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12L  | EPJ C72 2118            | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12O  | PL B713 172             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12P  | PL B713 369             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12Q  | PL B713 378             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12R  | PL B716 393             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12S  | PRL 108 151801          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12V  | PRL 108 201601          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 12W  | PRL 108 231801          | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AALTONEN   | 12AJ | PRL 109 171802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 12C  | PRL 108 201801          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 12D  | PR D85 072002           | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 12L  | PRL 108 211803          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 12AF | PR D86 092011           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 12C  | PR D85 011103           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 12D  | PR D85 032006           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| CHATRCHYAN | 12A  | JHEP 1204 033           | S. Chatrchyan <i>et al.</i> | (CMS Collab.)   |
| LEES       | 12A  | PR D85 011101           | J.P. Lees <i>et al.</i>     | (BABAR Collab.) |
| LI         | 12   | PRL 108 181808          | J. Li <i>et al.</i>         | (BELLE Collab.) |
| PDG        | 12   | PR D86 010001           | J. Beringer <i>et al.</i>   | (PDG Collab.)   |
| AAIJ       | 11   | PL B698 115             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 11A  | PL B698 14              | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 11B  | PL B699 330             | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AAIJ       | 11D  | PL B706 32              | R. Aaij                     | (LHCb Collab.)  |
| AAIJ       | 11E  | PR D84 092001           | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| Also       |      | PR D85 039904 (errat.)  | R. Aaij <i>et al.</i>       | (LHCb Collab.)  |
| AALTONEN   | 11A  | PR D83 052012           | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11AB | PR D84 052012           | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11AG | PRL 107 191801          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| Also       |      | PRL 107 239903 (errat.) | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11AI | PRL 107 201802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11AN | PRL 107 261802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11AP | PRL 107 272001          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11L  | PRL 106 161801          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 11N  | PRL 106 181802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 11U  | PR D84 052007           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| CHATRCHYAN | 11T  | PRL 107 191802          | S. Chatrchyan <i>et al.</i> | (CMS Collab.)   |
| LI         | 11   | PRL 106 121802          | J. Li <i>et al.</i>         | (BELLE Collab.) |
| ABAZOV     | 10E  | PR D82 012003           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 10H  | PRL 105 081801          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| Also       |      | PR D82 032001           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 10S  | PL B693 539             | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ESEN       | 10   | PRL 105 201802          | S. Esen <i>et al.</i>       | (BELLE Collab.) |
| LOUVOT     | 10   | PRL 104 231801          | R. LOUVOT <i>et al.</i>     | (BELLE Collab.) |
| PENG       | 10   | PR D82 072007           | C.-C. Peng <i>et al.</i>    | (BELLE Collab.) |
| AALTONEN   | 09AQ | PRL 103 191802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 09B  | PR D79 011104           | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 09C  | PRL 103 031801          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 09P  | PRL 102 201801          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 09E  | PRL 102 032001          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 09G  | PRL 102 051801          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 09I  | PRL 102 091801          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 09Y  | PR D79 111102           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| LOUVOT     | 09   | PRL 102 021801          | R. Louvot <i>et al.</i>     | (BELLE Collab.) |
| AALTONEN   | 08F  | PRL 100 021803          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 08G  | PRL 100 161802          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| AALTONEN   | 08J  | PRL 100 121803          | T. Aaltonen <i>et al.</i>   | (CDF Collab.)   |
| ABAZOV     | 08AM | PRL 101 241801          | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| WICHT      | 08A  | PRL 100 121801          | J. Wicht <i>et al.</i>      | (BELLE Collab.) |
| ABAZOV     | 07   | PRL 98 121801           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 07A  | PRL 98 151801           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 07N  | PR D76 057101           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 07Y  | PRL 99 241801           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABULENCIA  | 07C  | PRL 98 061802           | A. Abulencia <i>et al.</i>  | (CDF Collab.)   |
| DRUTSKOY   | 07   | PRL 98 052001           | A. Drutskoy <i>et al.</i>   | (BELLE Collab.) |
| DRUTSKOY   | 07A  | PR D76 012002           | A. Drutskoy <i>et al.</i>   | (BELLE Collab.) |
| ABAZOV     | 06B  | PRL 97 021802           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 06G  | PR D74 031107           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABAZOV     | 06V  | PRL 97 241801           | V.M. Abazov <i>et al.</i>   | (D0 Collab.)    |
| ABULENCIA  | 06J  | PRL 96 191801           | A. Abulencia <i>et al.</i>  | (CDF Collab.)   |

|             |     |                      |                             |                  |
|-------------|-----|----------------------|-----------------------------|------------------|
| ABULENCIA   | 06N | PRL 96 231801        | A. Abulencia <i>et al.</i>  | (CDF Collab.)    |
| ABULENCIA   | 06Q | PRL 97 062003        | A. Abulencia <i>et al.</i>  | (CDF Collab.)    |
| ABULENCIA,A | 06D | PRL 97 211802        | A. Abulencia <i>et al.</i>  | (CDF Collab.)    |
| ABULENCIA,A | 06G | PRL 97 242003        | A. Abulencia <i>et al.</i>  | (CDF Collab.)    |
| ACOSTA      | 06  | PRL 96 202001        | D. Acosta <i>et al.</i>     | (CDF Collab.)    |
| ABAZOV      | 05B | PRL 94 042001        | V.M. Abazov <i>et al.</i>   | (D0 Collab.)     |
| ABAZOV      | 05W | PRL 95 171801        | V.M. Abazov <i>et al.</i>   | (D0 Collab.)     |
| ACOSTA      | 05  | PRL 94 101803        | D. Acosta <i>et al.</i>     | (CDF Collab.)    |
| ACOSTA      | 05J | PRL 95 031801        | D. Acosta <i>et al.</i>     | (CDF Collab.)    |
| ABDALLAH    | 04A | PL B585 63           | J. Abdallah <i>et al.</i>   | (DELPHI Collab.) |
| ABDALLAH    | 04J | EPJ C35 35           | J. Abdallah <i>et al.</i>   | (DELPHI Collab.) |
| ABDALLAH    | 03B | EPJ C28 155          | J. Abdallah <i>et al.</i>   | (DELPHI Collab.) |
| ABE         | 03  | PR D67 012006        | K. Abe <i>et al.</i>        | (SLD Collab.)    |
| HEISTER     | 03E | EPJ C29 143          | A. Heister <i>et al.</i>    | (ALEPH Collab.)  |
| ABE         | 02V | PR D66 032009        | K. Abe <i>et al.</i>        | (SLD Collab.)    |
| ACOSTA      | 02D | PR D65 111101        | D. Acosta <i>et al.</i>     | (CDF Collab.)    |
| ACOSTA      | 02G | PR D66 112002        | D. Acosta <i>et al.</i>     | (CDF Collab.)    |
| ABBIENDI    | 01D | EPJ C19 241          | G. Abbiendi <i>et al.</i>   | (OPAL Collab.)   |
| ABE         | 00C | PR D62 071101        | K. Abe <i>et al.</i>        | (SLD Collab.)    |
| ABREU       | 00Y | EPJ C16 555          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| ABREU,P     | 00G | EPJ C18 229          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| AFFOLDER    | 00N | PRL 85 4668          | T. Affolder <i>et al.</i>   | (CDF Collab.)    |
| BARATE      | 00K | PL B486 286          | R. Barate <i>et al.</i>     | (ALEPH Collab.)  |
| ABBIENDI    | 99S | EPJ C11 587          | G. Abbiendi <i>et al.</i>   | (OPAL Collab.)   |
| ABE         | 99D | PR D59 032004        | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABE         | 99J | PRL 82 3576          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| BARATE      | 99J | EPJ C7 553           | R. Barate <i>et al.</i>     | (ALEPH Collab.)  |
| Also        |     | EPJ C12 181 (errat.) | R. Barate <i>et al.</i>     | (ALEPH Collab.)  |
| ABE         | 98B | PR D57 5382          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABE         | 98V | PRL 81 5742          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ACCIARRI    | 98S | PL B438 417          | M. Acciarri <i>et al.</i>   | (L3 Collab.)     |
| ACKERSTAFF  | 98F | EPJ C2 407           | K. Ackerstaff <i>et al.</i> | (OPAL Collab.)   |
| ACKERSTAFF  | 98G | PL B426 161          | K. Ackerstaff <i>et al.</i> | (OPAL Collab.)   |
| BARATE      | 98C | EPJ C4 367           | R. Barate <i>et al.</i>     | (ALEPH Collab.)  |
| BARATE      | 98Q | EPJ C4 387           | R. Barate <i>et al.</i>     | (ALEPH Collab.)  |
| PDG         | 98  | EPJ C3 1             | C. Caso <i>et al.</i>       | (PDG Collab.)    |
| ACCIARRI    | 97B | PL B391 474          | M. Acciarri <i>et al.</i>   | (L3 Collab.)     |
| ACCIARRI    | 97C | PL B391 481          | M. Acciarri <i>et al.</i>   | (L3 Collab.)     |
| ACKERSTAFF  | 97U | ZPHY C76 401         | K. Ackerstaff <i>et al.</i> | (OPAL Collab.)   |
| ACKERSTAFF  | 97V | ZPHY C76 417         | K. Ackerstaff <i>et al.</i> | (OPAL Collab.)   |
| ADAM        | 97  | PL B414 382          | W. Adam <i>et al.</i>       | (DELPHI Collab.) |
| ABE         | 96B | PR D53 3496          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABE         | 96N | PRL 77 1945          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABE         | 96Q | PR D54 6596          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABREU       | 96F | ZPHY C71 11          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| ADAM        | 96D | ZPHY C72 207         | W. Adam <i>et al.</i>       | (DELPHI Collab.) |
| BUSKULIC    | 96E | ZPHY C69 585         | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| BUSKULIC    | 96M | PL B377 205          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| BUSKULIC    | 96V | PL B384 471          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| PDG         | 96  | PR D54 1             | R. M. Barnett <i>et al.</i> | (PDG Collab.)    |
| ABE         | 95R | PRL 74 4988          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ABE         | 95Z | PRL 75 3068          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ACCIARRI    | 95H | PL B363 127          | M. Acciarri <i>et al.</i>   | (L3 Collab.)     |
| ACCIARRI    | 95I | PL B363 137          | M. Acciarri <i>et al.</i>   | (L3 Collab.)     |
| AKERS       | 95G | PL B350 273          | R. Akers <i>et al.</i>      | (OPAL Collab.)   |
| AKERS       | 95J | ZPHY C66 555         | R. Akers <i>et al.</i>      | (OPAL Collab.)   |
| BUSKULIC    | 95J | PL B356 409          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| BUSKULIC    | 95O | PL B361 221          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| ABREU       | 94D | PL B324 500          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| ABREU       | 94E | ZPHY C61 407         | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| Also        |     | PL B289 199          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| AKERS       | 94J | PL B337 196          | R. Akers <i>et al.</i>      | (OPAL Collab.)   |
| AKERS       | 94L | PL B337 393          | R. Akers <i>et al.</i>      | (OPAL Collab.)   |
| BUSKULIC    | 94B | PL B322 441          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| BUSKULIC    | 94C | PL B322 275          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| ABE         | 93F | PRL 71 1685          | F. Abe <i>et al.</i>        | (CDF Collab.)    |
| ACTON       | 93H | PL B312 501          | P.D. Acton <i>et al.</i>    | (OPAL Collab.)   |
| BUSKULIC    | 93G | PL B311 425          | D. Buskalic <i>et al.</i>   | (ALEPH Collab.)  |
| ABREU       | 92M | PL B289 199          | P. Abreu <i>et al.</i>      | (DELPHI Collab.) |
| ACTON       | 92N | PL B295 357          | P.D. Acton <i>et al.</i>    | (OPAL Collab.)   |

|              |     |             |                               |                   |
|--------------|-----|-------------|-------------------------------|-------------------|
| BUSKULIC     | 92E | PL B294 145 | D. Buskulic <i>et al.</i>     | (ALEPH Collab.)   |
| LEE-FRANZINI | 90  | PRL 65 2947 | J. Lee-Franzini <i>et al.</i> | (CUSB II Collab.) |

---