



$$I(J^P) = 0(0^-)$$

The angular distributions of the decays of the ϕ and $\bar{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\bar{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\bar{s}$ ground state.

D_s^\pm MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements. Measurements of the D_s^\pm mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.


VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

1968.35 ± 0.07 OUR FIT

1969.0 ± 1.4 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

1967.0 ± 1.0 ± 1.0	54	BARLAG	90C	ACCM	π^- Cu 230 GeV
1969.3 ± 1.4 ± 1.4		ALBRECHT	88	ARG	e^+e^- 9.4–10.6 GeV
1972.7 ± 1.5 ± 1.0	21	BECKER	87B	SILI	200 GeV π, K, p
1972.4 ± 3.7 ± 3.7	27	BLAYLOCK	87	MRK3	e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B	HRS	e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C	CLEO	e^+e^- 10.5 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1968.3 ± 0.7 ± 0.7	290	¹ ANJOS	88	E691	Photoproduction
1980 ± 15	6	USHIDA	86	EMUL	ν wideband
1973.6 ± 2.6 ± 3.0	163	ALBRECHT	85D	ARG	e^+e^- 10 GeV
1948 ± 28 ± 10	65	AIHARA	84D	TPC	e^+e^- 29 GeV
1975 ± 9 ± 10	49	ALTHOFF	84	TASS	e^+e^- 14–25 GeV
1975 ± 4	3	BAILEY	84	ACCM	hadron ⁺ Be → $\phi\pi^+X$

WEIGHTED AVERAGE
1969.0 ± 1.4 (Error scaled by 1.5)



¹ ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).

$m_{D_s^\pm} - m_{D^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
98.69±0.05 OUR FIT				
98.69±0.05 OUR AVERAGE				
98.68±0.03±0.04		AAIJ	13V LHCb	$D_s^+ \rightarrow K^+ K^- \pi^+$
99.41±0.38±0.21		ACOSTA	03D CDF2	$\bar{p}p, \sqrt{s}=1.96$ TeV
98.4 ±0.1 ±0.3	48k	AUBERT	02G BABR	$e^+e^- \approx \Upsilon(4S)$
99.5 ±0.6 ±0.3		BROWN	94 CLE2	$e^+e^- \approx \Upsilon(4S)$
98.5 ±1.5	555	CHEN	89 CLEO	e^+e^- 10.5 GeV
99.0 ±0.8	290	ANJOS	88 E691	Photoproduction

D_s^\pm MEAN LIFE

Measurements with an error greater than 100×10^{-15} s or with fewer than 100 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
504 ± 4 OUR AVERAGE Error includes scale factor of 1.2.				
506.4± 3.0± 1.7±1.7		¹ AAIJ	17AN LHCb	pp at 7, 8 TeV
507.4± 5.5± 5.1	13.6k	LINK	05J FOCS	$\phi\pi^+$ and $\bar{K}^{*0}K^+$
472.5±17.2± 6.6	760	IORI	01 SELX	600 GeV Σ^-, π^-, p
518 ±14 ± 7	1662	AITALA	99 E791	π^- nucleus, 500 GeV
486.3±15.0 ⁺ _{5.1}	2167	² BONVICINI	99 CLE2	$e^+e^- \approx \Upsilon(4S)$
475 ±20 ± 7	900	FRABETTI	93F E687	$\gamma\text{Be}, \phi\pi^+$
500 ±60 ±30	104	FRABETTI	90 E687	$\gamma\text{Be}, \phi\pi^+$
470 ±40 ±20	228	RAAB	88 E691	Photoproduction

¹ This AAIJ 17AN value is derived from the difference between the D_s^- and D^- widths.

The 3rd uncertainty, $\pm 1.7 \times 10^{-15}$ s, arises from the uncertainty of the D^- width.

² BONVICINI 99 obtains 1.19 ± 0.04 for the ratio of D_s^+ to D^0 lifetimes.

D_s^+ DECAY MODES

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Inclusive modes

Γ_1	e^+ semileptonic	[a] (6.33 \pm 0.15) %	
Γ_2	π^+ anything	(119.3 \pm 1.4) %	
Γ_3	π^- anything	(43.2 \pm 0.9) %	
Γ_4	π^0 anything	(123 \pm 7) %	
Γ_5	K^- anything	(18.7 \pm 0.5) %	
Γ_6	K^+ anything	(28.9 \pm 0.7) %	
Γ_7	K_S^0 anything	(19.0 \pm 1.1) %	
Γ_8	η anything	[b] (29.9 \pm 2.8) %	
Γ_9	ω anything	(6.1 \pm 1.4) %	
Γ_{10}	η' anything	[c] (10.3 \pm 1.4) %	S=1.1
Γ_{11}	$f_0(980)$ anything, $f_0 \rightarrow \pi^+\pi^-$	< 1.3 %	CL=90%
Γ_{12}	ϕ anything	(15.7 \pm 1.0) %	
Γ_{13}	K^+K^- anything	(15.8 \pm 0.7) %	
Γ_{14}	$K_S^0K^+$ anything	(5.8 \pm 0.5) %	
Γ_{15}	$K_S^0K^-$ anything	(1.9 \pm 0.4) %	
Γ_{16}	$2K_S^0$ anything	(1.70 \pm 0.32) %	
Γ_{17}	$2K^+$ anything	< 2.6 $\times 10^{-3}$	CL=90%
Γ_{18}	$2K^-$ anything	< 6 $\times 10^{-4}$	CL=90%

Leptonic and semileptonic modes

Γ_{19}	$e^+\nu_e$	< 8.3 $\times 10^{-5}$	CL=90%
Γ_{20}	$\mu^+\nu_\mu$	(5.43 \pm 0.15) $\times 10^{-3}$	
Γ_{21}	$\tau^+\nu_\tau$	(5.32 \pm 0.11) %	
Γ_{22}	$\gamma e^+\nu_e$	< 1.3 $\times 10^{-4}$	CL=90%
Γ_{23}	$K^+K^-e^+\nu_e$	—	
Γ_{24}	$K_S^0K_S^0e^+\nu_e$	< 3.8 $\times 10^{-4}$	CL=90%
Γ_{25}	$\phi e^+\nu_e$	[d] (2.39 \pm 0.16) %	S=1.3
Γ_{26}	$\phi\mu^+\nu_\mu$	(1.9 \pm 0.5) %	
Γ_{27}	$\eta e^+\nu_e + \eta'(958)e^+\nu_e$	[d] (3.03 \pm 0.24) %	
Γ_{28}	$\eta e^+\nu_e$	[d] (2.32 \pm 0.08) %	
Γ_{29}	$\eta'(958)e^+\nu_e$	[d] (8.0 \pm 0.7) $\times 10^{-3}$	
Γ_{30}	$\eta\mu^+\nu_\mu$	(2.4 \pm 0.5) %	
Γ_{31}	$\eta'(958)\mu^+\nu_\mu$	(1.1 \pm 0.5) %	
Γ_{32}	$\omega e^+\nu_e$	[e] < 2.0 $\times 10^{-3}$	CL=90%
Γ_{33}	$K^0 e^+\nu_e$	(3.4 \pm 0.4) $\times 10^{-3}$	
Γ_{34}	$K^*(892)^0 e^+\nu_e$	[d] (2.15 \pm 0.28) $\times 10^{-3}$	S=1.1
Γ_{35}	$f_0(500)e^+\nu_e, f_0 \rightarrow \pi^0\pi^0$	< 7.3 $\times 10^{-4}$	CL=90%
Γ_{36}	$f_0(980)e^+\nu_e, f_0 \rightarrow \pi^0\pi^0$	(7.9 \pm 1.5) $\times 10^{-4}$	
Γ_{37}	$f_0(980)e^+\nu_e, f_0 \rightarrow \pi^+\pi^-$		
Γ_{38}	$a_0(980)^0 e^+\nu_e, a_0(980)^0 \rightarrow \pi^0\eta$	< 1.2 $\times 10^{-4}$	CL=90%
Γ_{39}	$\pi^0 e^+\nu_e$	< 6.4 $\times 10^{-5}$	CL=90%

Hadronic modes with a $K\bar{K}$ pair

Γ_{40}	$K^+ K_S^0$		(1.450 ± 0.035) %	
Γ_{41}	$K^+ K_L^0$		(1.49 ± 0.06) %	
Γ_{42}	$K^+ \bar{K}^0$		(2.95 ± 0.14) %	
Γ_{43}	$K^+ K^- \pi^+$	[f]	(5.37 ± 0.10) %	S=1.1
Γ_{44}	$\phi \pi^+$	[d,g]	(4.5 ± 0.4) %	
Γ_{45}	$\phi \pi^+, \phi \rightarrow K^+ K^-$	[g]	(2.21 ± 0.06) %	
Γ_{46}	$K^+ \bar{K}^*(892)^0$		(12.7 $\begin{smallmatrix} +4.0 \\ -3.1 \end{smallmatrix}$) %	
Γ_{47}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$		(2.58 ± 0.06) %	
Γ_{48}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$		(4.8 ± 0.5) × 10 ⁻³	
Γ_{49}	$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$		(1.11 ± 0.19) %	
Γ_{50}	$f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$		(7.1 ± 2.9) × 10 ⁻⁴	
Γ_{51}	$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$		(6.7 ± 2.8) × 10 ⁻⁴	
Γ_{52}	$a_0(980)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$		(1.1 ± 0.4) × 10 ⁻³	
Γ_{53}	$a_0(1710)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$		(3.5 ± 0.6) × 10 ⁻³	
Γ_{54}	$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow$ $K^- \pi^+$		(1.76 ± 0.25) × 10 ⁻³	
Γ_{55}	$K^+ \bar{K}_0^*(1410)^0, \bar{K}_0^* \rightarrow$ $K_S^0 \pi^0$		(8.8 ± 2.8) × 10 ⁻⁴	
Γ_{56}	$K^+ K_S^0 \pi^0$		(1.47 ± 0.07) %	
Γ_{57}	$2K_S^0 \pi^+$		(7.1 ± 0.4) × 10 ⁻³	S=1.3
Γ_{58}	$f_0(980) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	< 1.8	× 10 ⁻⁴	CL=90%
Γ_{59}	$f_0(1710) \pi^+, f_0 \rightarrow K_S^0 K_S^0$		(3.3 ± 0.4) × 10 ⁻³	
Γ_{60}	$K^0 \bar{K}^0 \pi^+$		—	
Γ_{61}	$K^*(892)^+ \bar{K}^0$	[d]	(5.4 ± 1.2) %	
Γ_{62}	$K^*(892)^+ K_S^0$		(3.09 ± 0.33) × 10 ⁻³	
Γ_{63}	$K^*(892)^+ K_S^0, K^{*+} \rightarrow$ $K^+ \pi^0$		(2.04 ± 0.33) × 10 ⁻³	
Γ_{64}	$K^+ K^- \pi^+ \pi^0$		(5.50 ± 0.24) %	S=1.3
Γ_{65}	$\phi \rho^+$	[d]	(5.59 ± 0.34) %	
Γ_{66}	$\bar{K}_1(1270)^0 K^+, \bar{K}_1(1270)^0 \rightarrow K^- \rho^+$		(5.7 ± 0.6) × 10 ⁻³	
Γ_{67}	$\bar{K}_1(1270)^0 K^+, \bar{K}_1(1270)^0 \rightarrow K^*(892) \pi$		(1.31 ± 0.25) %	
Γ_{68}	$\bar{K}_1(1400)^0 K^+, \bar{K}_1(1400)^0 \rightarrow K^*(892) \pi$		(2.0 ± 0.4) %	
Γ_{69}	$a_0(980)^0 \rho^+, a_0(980)^0 \rightarrow$ $K^+ K^-$		(1.9 ± 0.4) × 10 ⁻³	
Γ_{70}	$f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow$ $K^*(892)^\mp K^\pm$		(3.9 ± 0.7) × 10 ⁻³	

Γ_{71}	$f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow$ $a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow$ $K^+ K^-$	(4.0 \pm 1.4) $\times 10^{-4}$	
Γ_{72}	$\eta(1475) \pi^+, \eta(1475) \rightarrow$ $a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow$ $K^+ K^-$	(7.0 \pm 2.8) $\times 10^{-4}$	
Γ_{73}	$K_S^0 K^- 2\pi^+$	(1.53 \pm 0.08) %	S=1.5
Γ_{74}	$K^*(892)^+ \bar{K}^*(892)^0$	[d] (5.64 \pm 0.35) %	
Γ_{75}	$\eta(1475) K_S^0, \eta \rightarrow$ $K^*(892)^0 \pi^+, K^{*0} \rightarrow$ $K^- \pi^+$	(3.4 \pm 1.0) $\times 10^{-4}$	
Γ_{76}	$\eta(1475) \pi^+, \eta \rightarrow$ $\bar{K}^*(892)^+ K^-, \bar{K}^{*+} \rightarrow$ $K_S^0 \pi^+$	(3.4 \pm 1.0) $\times 10^{-4}$	
Γ_{77}	$\eta(1475) \pi^+, \eta \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(1.7 \pm 0.9) $\times 10^{-3}$	
Γ_{78}	$f_1(1285) \pi^+, f_1 \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(3.4 \pm 0.8) $\times 10^{-4}$	
Γ_{79}	$K^+ K_S^0 \pi^+ \pi^-$	(9.5 \pm 0.8) $\times 10^{-3}$	S=1.1
Γ_{80}	$K^+ K^- 2\pi^+ \pi^-$	(6.6 \pm 0.6) $\times 10^{-3}$	
Γ_{81}	$\phi 2\pi^+ \pi^-$	[d] (1.21 \pm 0.16) %	
Γ_{82}	$\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-$	(4.9 \pm 0.7) $\times 10^{-3}$	
Γ_{83}	$\phi a_1(1260)^+, \phi \rightarrow$ $K^+ K^-, a_1^+ \rightarrow$ $\rho^0 \pi^+$	(7.4 \pm 1.2) $\times 10^{-3}$	
Γ_{84}	$\phi 2\pi^+ \pi^-$ non- $\rho, \phi \rightarrow$ $K^+ K^-$	(1.4 \pm 0.5) $\times 10^{-3}$	
Γ_{85}	$K^+ K^- \rho^0 \pi^+$ non- ϕ	< 2.0 $\times 10^{-4}$	CL=90%
Γ_{86}	$K^+ K^- 2\pi^+ \pi^-$ nonresonant	(1.0 \pm 0.4) $\times 10^{-3}$	
Γ_{87}	$2K_S^0 2\pi^+ \pi^-$	(7.8 \pm 3.3) $\times 10^{-4}$	

Hadronic modes without K 's

Γ_{88}	$\pi^+ \pi^0$	< 1.2 $\times 10^{-4}$	CL=90%
Γ_{89}	$2\pi^+ \pi^-$	(1.08 \pm 0.04) %	
Γ_{90}	$\rho^0 \pi^+$	(1.2 \pm 0.6) $\times 10^{-4}$	
Γ_{91}	$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	[h] (9.0 \pm 0.4) $\times 10^{-3}$	
Γ_{92}	$f_0(980) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{93}	$f_0(1370) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{94}	$f_0(1500) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{95}	$f_2(1270) \pi^+, f_2 \rightarrow \pi^+ \pi^-$	(1.11 \pm 0.12) $\times 10^{-3}$	
Γ_{96}	$\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	(1.6 \pm 0.7) $\times 10^{-4}$	
Γ_{97}	$\pi^+ 2\pi^0$	(5.2 \pm 0.5) $\times 10^{-3}$	S=1.1

Γ_{98}	$f_0(980)\pi^+$, $f_0 \rightarrow \pi^0\pi^0$	(2.9 \pm 0.6) $\times 10^{-3}$	
Γ_{99}	$f_0(1370)\pi^+$, $f_0 \rightarrow \pi^0\pi^0$	(1.3 \pm 0.6) $\times 10^{-3}$	
Γ_{100}	$f_2(1270)\pi^+$, $f_2 \rightarrow \pi^0\pi^0$	(5.0 \pm 3.5) $\times 10^{-4}$	
Γ_{101}	$2\pi^+\pi^-\pi^0$	—	
Γ_{102}	$\eta\pi^+$	[d] (1.67 \pm 0.09) %	S=1.1
Γ_{103}	$\omega\pi^+$	[d] (1.92 \pm 0.30) $\times 10^{-3}$	
Γ_{104}	$3\pi^+2\pi^-$	(7.8 \pm 0.8) $\times 10^{-3}$	
Γ_{105}	$2\pi^+\pi^-2\pi^0$	—	
Γ_{106}	$\eta\rho^+$	[d] (8.9 \pm 0.8) %	
Γ_{107}	$\eta\pi^+\pi^0$	(9.5 \pm 0.5) %	
Γ_{108}	$\eta(\pi^+\pi^0)_{P\text{-wave}}$	(5.1 \pm 3.1) $\times 10^{-3}$	
Γ_{109}	$a_0(980)^{+0}\pi^{0+}$, $a_0(980)^{+0} \rightarrow \eta\pi^{+0}$	(2.2 \pm 0.4) %	
Γ_{110}	$\omega\pi^+\pi^0$	[d] (2.8 \pm 0.7) %	
Γ_{111}	$2\pi^+\pi^-\eta$	(3.12 \pm 0.16) %	
Γ_{112}	$a_1(1260)^+\eta$, $a_1^+ \rightarrow$ $\rho(770)^0\pi^+$, $\rho^0 \rightarrow \pi^+\pi^-$	(1.73 \pm 0.16) %	
Γ_{113}	$a_1(1260)^+\eta$, $a_1^+ \rightarrow$ $f_0(500)\pi^+$, $f_0 \rightarrow \pi^+\pi^-$	(2.5 \pm 0.9) $\times 10^{-3}$	
Γ_{114}	$a_0(980)^+\rho(770)^0$, $a_0^+ \rightarrow$ $\eta\pi^+$	(2.1 \pm 0.9) $\times 10^{-3}$	
Γ_{115}	$\eta(1405)\pi^+$, $\eta(1405) \rightarrow$ $a_0(980)^-\pi^+$, $a_0^- \rightarrow \eta\pi^-$	(2.2 \pm 0.7) $\times 10^{-4}$	
Γ_{116}	$\eta(1405)\pi^+$, $\eta(1405) \rightarrow$ $a_0(980)^+\pi^-$, $a_0^+ \rightarrow \eta\pi^+$	(2.2 \pm 0.7) $\times 10^{-4}$	
Γ_{117}	$f_1(1420)\pi^+$, $f_1 \rightarrow$ $a_0(980)^-\pi^+$, $a_0^- \rightarrow \eta\pi^-$	(5.9 \pm 1.8) $\times 10^{-4}$	
Γ_{118}	$f_1(1420)\pi^+$, $f_1 \rightarrow$ $a_0(980)^+\pi^-$, $a_0^+ \rightarrow \eta\pi^+$	(5.3 \pm 1.8) $\times 10^{-4}$	
Γ_{119}	$3\pi^+2\pi^-\pi^0$	(4.9 \pm 3.2) %	
Γ_{120}	$\omega 2\pi^+\pi^-$	[d] (1.6 \pm 0.5) %	
Γ_{121}	$\eta'(958)\pi^+$	[c,d] (3.94 \pm 0.25) %	
Γ_{122}	$3\pi^+2\pi^-2\pi^0$	—	
Γ_{123}	$\omega\eta\pi^+$	[d] < 2.13 %	CL=90%
Γ_{124}	$\eta'(958)\rho^+$	[c,d] (5.8 \pm 1.5) %	
Γ_{125}	$\eta'(958)\pi^+\pi^0$	(6.08 \pm 0.29) %	
Γ_{126}	$\eta'(958)\pi^+\pi^0$ nonresonant	< 5.1 %	CL=90%

Modes with one or three K's

Γ_{127}	$K^+\pi^0$	(7.4 \pm 0.5) $\times 10^{-4}$	
Γ_{128}	$K_S^0\pi^+$	(1.09 \pm 0.05) $\times 10^{-3}$	
Γ_{129}	$K^+\eta$	[d] (1.73 \pm 0.08) $\times 10^{-3}$	
Γ_{130}	$K^+\omega$	[d] (9.9 \pm 1.5) $\times 10^{-4}$	

Γ_{131}	$K^+ \eta'(958)$	[d]	$(2.64 \pm 0.24) \times 10^{-3}$	
Γ_{132}	$K^+ \pi^+ \pi^-$		$(6.20 \pm 0.19) \times 10^{-3}$	
Γ_{133}	$K^+ \rho^0$		$(2.17 \pm 0.25) \times 10^{-3}$	
Γ_{134}	$K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-$		$(7.2 \pm 1.7) \times 10^{-4}$	
Γ_{135}	$K^+ f_0(500), f_0 \rightarrow \pi^+ \pi^-$		$(4.5 \pm 3.0) \times 10^{-4}$	
Γ_{136}	$K^+ f_0(980), f_0 \rightarrow \pi^+ \pi^-$		$(2.8 \pm 1.1) \times 10^{-4}$	
Γ_{137}	$K^+ f_0(1370), f_0 \rightarrow \pi^+ \pi^-$		$(1.2 \pm 0.6) \times 10^{-3}$	
Γ_{138}	$K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$		$(1.67 \pm 0.26) \times 10^{-3}$	
Γ_{139}	$K^*(1410)^0 \pi^+, K^{*0} \rightarrow$ $K^+ \pi^-$		$(6 \pm 4) \times 10^{-4}$	
Γ_{140}	$K^*(1430)^0 \pi^+, K^{*0} \rightarrow$ $K^+ \pi^-$		$(9.3 \pm 3.1) \times 10^{-4}$	
Γ_{141}	$K^+ \pi^+ \pi^-$ nonresonant		$(9.9 \pm 3.2) \times 10^{-4}$	
Γ_{142}	$K^0 \pi^+ \pi^0$		$(1.08 \pm 0.06) \%$	
Γ_{143}	$K_S^0 2\pi^+ \pi^-$		$(2.8 \pm 1.0) \times 10^{-3}$	
Γ_{144}	$K^+ \pi^+ \pi^- \pi^0$		$(9.7 \pm 0.6) \times 10^{-3}$	
Γ_{145}	$K^*(892)^0 \rho^+, K^{*0} \rightarrow K^+ \pi^-$		$(3.9 \pm 0.4) \times 10^{-3}$	
Γ_{146}	$K^*(892)^+ \rho^0, K^{*+} \rightarrow K^+ \pi^0$		$(4.2 \pm 1.2) \times 10^{-4}$	
Γ_{147}	$K_1(1270)^0 \pi^+, K_1^0 \rightarrow K^+ \rho^-$		$(3.9 \pm 1.3) \times 10^{-4}$	
Γ_{148}	$K_1(1400)^0 \pi^+, K_1^0 \rightarrow$ $K^*(890)^+ \pi^-, K^{*+} \rightarrow$ $K^+ \pi^0$		$(5.4 \pm 0.9) \times 10^{-4}$	
Γ_{149}	$K_1(1400)^0 \pi^+, K_1^0 \rightarrow$ $K^*(890)^0 \pi^0, K^{*0} \rightarrow$ $K^+ \pi^-$		$(5.9 \pm 1.0) \times 10^{-4}$	
Γ_{150}	$K^+ a_1(1260)^0, a_1 \rightarrow \rho^+ \pi^-$		$(1.8 \pm 1.1) \times 10^{-4}$	
Γ_{151}	$K^+ a_1(1260)^0, a_1 \rightarrow \rho^- \pi^+$		$(1.8 \pm 1.1) \times 10^{-4}$	
Γ_{152}	$K^+ \pi^+ \pi^- \pi^0$ nonresonant		$(9.2 \pm 2.4) \times 10^{-4}$	
Γ_{153}	$(K^+ \pi^0) P\text{-wave } \rho^0$		$(1.01 \pm 0.21) \times 10^{-3}$	
Γ_{154}	$K^+ \omega \pi^0$	[d]	$< 8.2 \times 10^{-3}$	CL=90%
Γ_{155}	$K^+ \omega \pi^+ \pi^-$	[d]	$< 5.4 \times 10^{-3}$	CL=90%
Γ_{156}	$K^+ \omega \eta$	[d]	$< 7.9 \times 10^{-3}$	CL=90%
Γ_{157}	$2K^+ K^-$		$(2.15 \pm 0.20) \times 10^{-4}$	
Γ_{158}	$\phi K^+, \phi \rightarrow K^+ K^-$		$(8.8 \pm 2.0) \times 10^{-5}$	

Doubly Cabibbo-suppressed modes

Γ_{159}	$2K^+ \pi^-$		$(1.274 \pm 0.031) \times 10^{-4}$	
Γ_{160}	$K^+ K^*(892)^0, K^{*0} \rightarrow$ $K^+ \pi^-$		$(6.0 \pm 3.4) \times 10^{-5}$	

Baryon-antibaryon mode

Γ_{161}	$p \bar{n}$		$(1.22 \pm 0.11) \times 10^{-3}$	
Γ_{162}	$p \bar{p} e^+ \nu_e$	<	2.0×10^{-4}	CL=90%

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

Γ_{163}	$\pi^+ e^+ e^-$		$[i] < 5.5$	$\times 10^{-6}$	CL=90%
Γ_{164}	$\pi^+ \phi, \phi \rightarrow e^+ e^-$		$[j] (6 \quad +8) \times 10^{-6}$		
				-4	
Γ_{165}	$\pi^+ \mu^+ \mu^-$		$[i] < 1.8$	$\times 10^{-7}$	CL=90%
Γ_{166}	$K^+ e^+ e^-$	C1	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{167}	$K^+ \mu^+ \mu^-$	C1	< 1.4	$\times 10^{-7}$	CL=90%
Γ_{168}	$K^*(892)^+ \mu^+ \mu^-$	C1	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{169}	$\pi^+ e^+ \mu^-$	LF	< 1.1	$\times 10^{-6}$	CL=90%
Γ_{170}	$\pi^+ e^- \mu^+$	LF	< 9.4	$\times 10^{-7}$	CL=90%
Γ_{171}	$K^+ e^+ \mu^-$	LF	< 7.9	$\times 10^{-7}$	CL=90%
Γ_{172}	$K^+ e^- \mu^+$	LF	< 5.6	$\times 10^{-7}$	CL=90%
Γ_{173}	$\pi^- 2e^+$	L	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{174}	$\pi^- 2\mu^+$	L	< 8.6	$\times 10^{-8}$	CL=90%
Γ_{175}	$\pi^- e^+ \mu^+$	L	< 6.3	$\times 10^{-7}$	CL=90%
Γ_{176}	$K^- 2e^+$	L	< 7.7	$\times 10^{-7}$	CL=90%
Γ_{177}	$K^- 2\mu^+$	L	< 2.6	$\times 10^{-8}$	CL=90%
Γ_{178}	$K^- e^+ \mu^+$	L	< 2.6	$\times 10^{-7}$	CL=90%
Γ_{179}	$K^*(892)^- 2\mu^+$	L	< 1.4	$\times 10^{-3}$	CL=90%

- [a] This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+ exclusive fractions — an $e^+ \nu_e$ with an η, η', ϕ, K^0 , or K^{*0} — is 5.99 ± 0.31 %.
- [b] This fraction includes η from η' decays.
- [c] The sum of our exclusive η' fractions — $\eta' e^+ \nu_e, \eta' \mu^+ \nu_\mu, \eta' \pi^+, \eta' \rho^+$, and $\eta' K^+$ — is 11.8 ± 1.6 %.
- [d] This branching fraction includes all the decay modes of the final-state resonance.
- [e] A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and $\omega-\phi$ mixing is an unlikely explanation for any fraction above about 2×10^{-4} .
- [f] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [g] We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+, \phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

- [h] This is the average of a model-independent and a K -matrix parametrization of the $\pi^+\pi^-$ S -wave and is a sum over several f_0 mesons.
- [i] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [j] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+\ell^+\ell^-$ final state.

CONSTRAINED FIT INFORMATION

An overall fit to 16 branching ratios uses 25 measurements and one constraint to determine 12 parameters. The overall fit has a $\chi^2 = 12.7$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{43}	27									
x_{64}	8	0								
x_{73}	24	4	14							
x_{79}	18	3	12	45						
x_{89}	17	33	1	6	4					
x_{102}	1	15	-8	-15	-12	6				
x_{103}	0	1	0	-1	0	0	4			
x_{130}	0	0	0	0	0	0	0	0		
x_{132}	7	3	4	8	6	2	-4	0	0	
x_{144}	0	0	0	0	0	0	0	0	26	0
	x_{40}	x_{43}	x_{64}	x_{73}	x_{79}	x_{89}	x_{102}	x_{103}	x_{130}	x_{132}

See the related review(s):

D_s⁺ Branching Fractions

D_s⁺ BRANCHING RATIOS

A number of older, now obsolete results have been omitted. They may be found in earlier editions.

Inclusive modes

$\Gamma(e^+ \text{ semileptonic}) / \Gamma_{\text{total}}$

Γ_1 / Γ

This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.33 ± 0.15 OUR AVERAGE				
6.30 ± 0.13 ± 0.10	17k	^{1,2} ABLIKIM	21AC BES3	e^+e^- at 4.178–4.230 GeV
6.52 ± 0.39 ± 0.15	0.5k	³ ASNER	10 CLEO	e^+e^- at 3774 MeV

¹ ABLIKIM 21AC finds that the ratio of the D_s^+ and D^0 semielectronic widths is $0.790 \pm 0.016 \pm 0.020$.

² ABLIKIM 21AC reports a value of $(6.30 \pm 0.13 \pm 0.09 \pm 0.04) \times 10^{-2}$, where the last uncertainty is an external systematic from $B(D_S^+ \rightarrow \tau \nu)$. We have added the systematic uncertainties in quadrature.

³ Using the D_S^+ and D^0 lifetimes, ASNER 10 finds that the ratio of the D_S^+ and D^0 semileptonic widths is $0.828 \pm 0.051 \pm 0.025$.

$\Gamma(\pi^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_2/Γ

Events with two π^+ 's count twice, etc. But π^+ 's from $K_S^0 \rightarrow \pi^+ \pi^-$ are not included.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
119.3±1.2±0.7	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\pi^- \text{ anything})/\Gamma_{\text{total}}$ Γ_3/Γ

Events with two π^- 's count twice, etc. But π^- 's from $K_S^0 \rightarrow \pi^+ \pi^-$ are not included.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
43.2±0.9±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_4/Γ

Events with two π^0 's count twice, etc. But π^0 's from $K_S^0 \rightarrow 2\pi^0$ are not included.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
123.4±3.8±5.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
18.7±0.5±0.2	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
28.9±0.6±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19.0±1.0±0.4	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

This ratio includes η particles from η' decays.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29.9±2.2±1.7		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

23.5±3.1±2.0	674 ± 91	HUANG	06B	CLEO See DOBBS 09
--------------	----------	-------	-----	-------------------

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±1.4±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.3±1.4 OUR AVERAGE	Error includes scale factor of 1.1.			
8.8±1.8±0.5	68	ABLIKIM	15Z BES3	482 pb ⁻¹ , 4009 MeV
11.7±1.7±0.7		DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.7±1.9±0.8	68	HUANG	06B CLEO	See DOBBS 09

$\Gamma(f_0(980) \text{ anything}, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
15.7±0.8±0.6		DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
16.1±1.2±1.1	398 ± 27	HUANG	06B CLEO	See DOBBS 09

$\Gamma(K^+ K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
15.8±0.6±0.3	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(K_S^0 K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.8±0.5±0.1	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(K_S^0 K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.9±0.4±0.1	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(2K_S^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.7±0.3±0.1	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(2K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.26	90	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(2K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.06	90	DOBBS	09 CLEO	e ⁺ e ⁻ at 4170 MeV

———— **Leptonic and semileptonic modes** ————

See the related review(s):

[Leptonic Decays of Charged Pseudoscalar Mesons](#)

$\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.83 × 10⁻⁴	90	¹ ZUPANC	13 BELL	e ⁺ e ⁻ at $\Upsilon(4S), \Upsilon(5S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.3 \times 10^{-4}$	90	DEL-AMO-SA..10J	BABR	e^+e^- , 10.58 GeV
$<1.2 \times 10^{-4}$	90	ALEXANDER 09	CLEO	e^+e^- at 4170 MeV
$<1.3 \times 10^{-4}$	90	PEDLAR 07A	CLEO	See ALEXANDER 09

¹ZUPANC 13 also gives the limit as $< 1.0 \times 10^{-4}$ at 95% CL.

$\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$

Γ_{20}/Γ

See the note on "Decay Constants of Charged Pseudoscalar Mesons."

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.43±0.15 OUR AVERAGE				
5.35±0.13±0.16	2.2k	ABLIKIM	21BE BES3	e^+e^- , 4.178, 4.226 GeV
5.17±0.75±0.21	69	¹ ABLIKIM	160 BES3	e^+e^- at 4.009 GeV
5.31±0.28±0.20	492 ± 26	² ZUPANC	13 BELL	e^+e^- at $\Upsilon(4S), \Upsilon(5S)$
6.02±0.38±0.34	275 ± 17	³ DEL-AMO-SA..10J	BABR	e^+e^- , 10.58 GeV
5.65±0.45±0.17	235 ± 14	ALEXANDER 09	CLEO	e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.49±0.16±0.15	1.1k	ABLIKIM	19E BES3	e^+e^- at 4178 MeV
6.44±0.76±0.57	169 ± 18	⁴ WIDHALM	08 BELL	See ZUPANC 13
5.94±0.66±0.31	88	⁵ PEDLAR	07A CLEO	See ALEXANDER 09
6.8 ± 1.1 ± 1.8	553	⁶ HEISTER	02I ALEP	Z decays

¹ABLIKIM 160 also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+\nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+\nu_\mu) = 9.76$, the branching fraction is found to be $(0.495 \pm 0.067 \pm 0.026)\%$. The constrained value is used to obtain the decay constant, $f_{D_s^+} = (241.0 \pm 16.3 \pm 6.6)$ MeV.

²ZUPANC 13 uses both $\mu^+\nu$ and $\tau^+\nu$ events to get $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$ MeV.

³DEL-AMO-SANCHEZ 10J uses $\mu^+\nu_\mu$ and $\tau^+\nu_\tau$ events together to get $f_{D_s} = (258.6 \pm 6.4 \pm 7.5)$ MeV.

⁴WIDHALM 08 gets $f_{D_s} = (275 \pm 16 \pm 12)$ MeV from the branching fraction.

⁵PEDLAR 07A also fits μ^+ and τ^+ events together and gets an effective $\mu^+\nu_\mu$ branching fraction of $(6.38 \pm 0.59 \pm 0.33) \times 10^{-3}$

⁶This HEISTER 02I result is not actually an independent measurement of the absolute $\mu^+\nu_\mu$ branching fraction, but is in fact based on our $\phi\pi^+$ branching fraction of 3.6 ± 0.9%, so it cannot be included in our overall fit. HEISTER 02I combines its $D_s^+ \rightarrow \tau^+\nu_\tau$ and $\mu^+\nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

$\Gamma(\mu^+\nu_\mu)/\Gamma(\phi\pi^+)$

Γ_{20}/Γ_{44}

See the note on "Decay Constants of Charged Pseudoscalar Mesons" above.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.143±0.018±0.006	489 ± 55	¹ AUBERT	07V BABR	$e^+e^- \approx \Upsilon(4S)$
0.23 ± 0.06 ± 0.04	18	² ALEXANDROV	00 BEAT	π^- nucleus, 350 GeV
0.173±0.023±0.035	182	³ CHADHA	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
0.245±0.052±0.074	39	⁴ ACOSTA	94 CLE2	See CHADHA 98

¹AUBERT 07V gets $f_{D_s^+} = (283 \pm 17 \pm 16)$ MeV, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = (4.71 \pm 0.46)\%$.

² ALEXANDROV 00 uses $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$ from a lattice-gauge-theory calculation to get the relative numbers of $D^+ \rightarrow \mu^+ \nu_\mu$ and $D_s^+ \rightarrow \mu^+ \nu_\mu$ events. The present result leads to $f_{D_s} = (323 \pm 44 \pm 36)$ MeV.

³ CHADHA 98 obtains $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$.

⁴ ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$

Γ_{21}/Γ

See the note on “Decay Constants of Charged Pseudoscalar Mesons” above.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.32±0.11 OUR AVERAGE				
5.29±0.25±0.20	1.7k	¹ ABLIKIM	21AF BES3	$e^+ e^-$ at 4.178, 4.226 GeV
5.27±0.10±0.12	4.9k	² ABLIKIM	21AZ BES3	$e^+ e^-$ at 4.178, 4.226 GeV
5.21±0.25±0.17	950	³ ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV
3.28±1.83±0.37	33	⁴ ABLIKIM	160 BES3	$e^+ e^-$ at 4.009 GeV
5.70±0.21 ^{+0.31} _{-0.30}	2.2k	⁵ ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S)$, $\Upsilon(5S)$
4.96±0.37±0.57	748	⁶ DEL-AMO-SA..10J	BABR	$e^- \bar{\nu}_e \nu_\tau, \mu^- \bar{\nu}_\mu \nu_\tau$
6.42±0.81±0.18	126	⁷ ALEXANDER 09	CLEO	$\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$
5.52±0.57±0.21	155	⁷ NAIK 09A	CLEO	$\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$
5.30±0.47±0.22	181	⁷ ONYISI 09	CLEO	$\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
6.17±0.71±0.34	102	⁸ ECKLUND 08	CLEO	See ONYISI 09
8.0 ±1.3 ±0.4	47	⁸ PEDLAR 07A	CLEO	See ALEXANDER 09
5.79±0.77±1.84	881	⁹ HEISTER 02I	ALEP	Z decays
7.0 ±2.1 ±2.0	22	¹⁰ ABBIENDI 01L	OPAL	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's
7.4 ±2.8 ±2.4	16	¹¹ ACCIARRI 97F	L3	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's

¹ ABLIKIM 21F uses $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}$ decays.

² ABLIKIM 21AZ uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays.

³ ABLIKIM 21BE uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays. When constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the branching fraction is found to be $(5.22 \pm 0.10 \pm 0.14)\%$.

⁴ ABLIKIM 160 also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.76$; the branching fraction is found to be $(4.83 \pm 0.65 \pm 0.26)\%$.

⁵ ZUPANC 13 uses both $\mu^+ \nu$ and $\tau^+ \nu$ events to get $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$ MeV.

⁶ DEL-AMO-SANCHEZ 10J (with a small correction; see LEES 15D) uses $\mu^+ \nu_\mu$ and $\tau^+ \nu_\tau$ events together to get $f_{D_s} = (259.9 \pm 6.6 \pm 7.6)$ MeV.

⁷ ALEXANDER 09, NAIK 09A, and ONYISI 09 use different τ decay modes and are independent. The three papers combined give $f_{D_s} = (259.7 \pm 7.8 \pm 3.4)$ MeV.

⁸ ECKLUND 08 and PEDLAR 07A are independent: ECKLUND 08 uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ events, PEDLAR 07A uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ events.

⁹ HEISTER 02I combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

¹⁰ This ABBIENDI 01L value gives a decay constant f_{D_s} of $(286 \pm 44 \pm 41)$ MeV.

¹¹ The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives $f_{D_s} = (309 \pm 58 \pm 33 \pm 38)$ MeV.

$\Gamma(\tau^+ \nu_\tau) / \Gamma(\mu^+ \nu_\mu)$ $\Gamma_{21} / \Gamma_{20}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.73 \pm 0.69^{+0.56}_{-0.53}$	2.2k/492	¹ ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$
----------------------------------	----------	---------------------	---------	---

$11.0 \pm 1.4 \pm 0.6$	102	² ECKLUND	08 CLEO	See ONYISI 09
------------------------	-----	----------------------	---------	---------------

¹ This ZUPANC 13 ratio is not independent of the separate $\tau\nu$ and $\mu\nu$ fractions listed above.

² This ECKLUND 08 value also uses results from PEDLAR 07A, and it is not independent of other results in these Listings. Combined with earlier CLEO results, the decay constant f_{D_s} is $274 \pm 10 \pm 5$ MeV.

$\Gamma(\gamma e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{22} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 3.8 \times 10^{-4}$	90	ABLIKIM	19AD BES3	for $E_\gamma > 10$ MeV
------------------------	----	---------	-----------	-------------------------

$\Gamma(K^+ K^- e^+ \nu_e) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{23} / \Gamma_{43}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.558 \pm 0.007 \pm 0.016$	¹ AUBERT	08AN BABR	$e^+ e^-$ at $\Upsilon(4S)$
-----------------------------	---------------------	-----------	-----------------------------

¹ This AUBERT 08AN ratio is only for the $K^+ K^-$ mass in the range 1.01–to–1.03 GeV in the numerator and 1.0095–to–1.0295 GeV in the denominator.

$\Gamma(K_S^0 K_S^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{24} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 3.8 \times 10^{-4}$	90	ABLIKIM	22J BES3	$e^+ e^-$ at 4.178–4.226 GeV
------------------------	----	---------	----------	------------------------------

$\Gamma(\phi e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{25} / Γ

See the end of the D_s^+ Listings for measurements of $D_s^+ \rightarrow \phi e^+ \nu_e$ form factors. Unseen decay modes of the ϕ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.39 ± 0.16 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

$2.26 \pm 0.45 \pm 0.09$	26	ABLIKIM	18A BES3	$e^+ e^-$ at 4.009 GeV
--------------------------	----	---------	----------	------------------------


$2.14 \pm 0.17 \pm 0.08$	207	HIETALA	15	Uses CLEO data
--------------------------	-----	---------	----	----------------

$2.61 \pm 0.03 \pm 0.17$	25k	AUBERT	08AN BABR	$e^+ e^-$ at $\Upsilon(4S)$
--------------------------	-----	--------	-----------	-----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.36 \pm 0.23 \pm 0.13$	106	ECKLUND	09 CLEO	See HIETALA 15
--------------------------	-----	---------	---------	----------------

$2.29 \pm 0.37 \pm 0.11$	45	YELTON	09 CLEO	See ECKLUND 09
--------------------------	----	--------	---------	----------------



$\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$

$\Gamma_{25} / \Gamma_{44}$

As noted in the comment column, most of these measurements use $\phi \mu^+ \nu_\mu$ events in addition to or instead of $\phi e^+ \nu_e$ events.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.540 \pm 0.033 \pm 0.048$	793	LINK	02J FOCs	Uses $\phi \mu^+ \nu_\mu$
$0.54 \pm 0.05 \pm 0.04$	367	BUTLER	94 CLE2	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$
$0.58 \pm 0.17 \pm 0.07$	97	FRABETTI	93G E687	Uses $\phi \mu^+ \nu_\mu$
$0.57 \pm 0.15 \pm 0.15$	104	ALBRECHT	91 ARG	Uses $\phi e^+ \nu_e$
$0.49 \pm 0.10 \begin{smallmatrix} +0.10 \\ -0.14 \end{smallmatrix}$	54	ALEXANDER	90B CLEO	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$

$\Gamma(\phi \mu^+ \nu_\mu) / \Gamma_{\text{total}}$

Γ_{26} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.94 \pm 0.53 \pm 0.09$	22	ABLIKIM	18A BES3	$e^+ e^-$ at 4.009 GeV

$\Gamma(\eta e^+ \nu_e) / \Gamma_{\text{total}}$

Γ_{28} / Γ

Unseen decay modes of the η are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.32 ± 0.08 OUR AVERAGE				
$2.323 \pm 0.063 \pm 0.063$	1.8k	ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV
$2.30 \pm 0.31 \pm 0.08$	63	ABLIKIM	16T BES3	$e^+ e^-$ at 4.009 GeV
$2.28 \pm 0.14 \pm 0.19$	358	¹ HIETALA	15	Uses CLEO data
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.48 \pm 0.29 \pm 0.13$	82	YELTON	09 CLEO	See HIETALA 15

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\eta e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$ Γ_{28}/Γ_{25}

Unseen decay modes of the η and the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.24 ± 0.12 ± 0.15 440 ¹ BRANDENB... 95 CLE2 See HIETALA 15

¹ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma_{total}$ Γ_{29}/Γ

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.80 ± 0.07 OUR AVERAGE

0.824 ± 0.073 ± 0.027 261 ABLIKIM 19S BES3 $e^+ e^-$ at 4178 MeV

0.93 ± 0.30 ± 0.05 14 ABLIKIM 16T BES3 $e^+ e^-$ at 4009 MeV

0.68 ± 0.15 ± 0.06 20 ¹ HIETALA 15 Uses CLEO data

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.91 ± 0.33 ± 0.05 7.5 YELTON 09 CLEO See HIETALA 15

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$ Γ_{29}/Γ_{25}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.43 ± 0.11 ± 0.07 29 ¹ BRANDENB... 95 CLE2 See HIETALA 15

¹ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$[\Gamma(\eta e^+ \nu_e) + \Gamma(\eta'(958) e^+ \nu_e)]/\Gamma(\phi e^+ \nu_e)$ $\Gamma_{27}/\Gamma_{25} = (\Gamma_{28} + \Gamma_{29})/\Gamma_{25}$

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.67 ± 0.17 ± 0.17 ¹ BRANDENB... 95 CLE2 See HIETALA 15

¹ This BRANDENBURG 95 data is redundant with data in previous blocks.

$\Gamma(\eta \mu^+ \nu_\mu)/\Gamma_{total}$ Γ_{30}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.42 ± 0.46 ± 0.11 44 ABLIKIM 18A BES3 $e^+ e^-$ at 4.009 GeV

$\Gamma(\eta'(958) \mu^+ \nu_\mu)/\Gamma_{total}$ Γ_{31}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.06 ± 0.54 ± 0.07 10 ABLIKIM 18A BES3 $e^+ e^-$ at 4.009 GeV

$\Gamma(\omega e^+ \nu_e)/\Gamma_{total}$ Γ_{32}/Γ

A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and $\omega - \phi$ mixing is an unlikely explanation for any fraction above about 2×10^{-4} .

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

<0.20 90 MARTIN 11 CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.04 OUR AVERAGE				
0.325 ± 0.038 ± 0.016	117	¹ ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV
0.39 ± 0.08 ± 0.03	42	HIETALA	15	Uses CLEO data
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.37 ± 0.10 ± 0.02	14	YELTON	09 CLEO	See HIETALA 15
¹ K^0 reconstructed via $K^0 \rightarrow K_S^0 \rightarrow \pi^+ \pi^-$ decays.				

$\Gamma(K^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{34}/Γ

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.215 ± 0.028 OUR AVERAGE				Error includes scale factor of 1.1.
0.237 ± 0.026 ± 0.020	155	ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV
0.18 ± 0.04 ± 0.01	32	¹ HIETALA	15	$e^+ e^-$ at 4.170 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.18 ± 0.07 ± 0.01	7.5	YELTON	09 CLEO	See HIETALA 15
¹ Uses CLEO data, but not authored by the CLEO collaboration				

$\Gamma(f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7.3 × 10⁻⁴	90	ABLIKIM	22J BES3	$e^+ e^-$ at 4.178-4.226 GeV

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
7.9 ± 1.4 ± 0.4	55	¹ ABLIKIM	22J BES3	$e^+ e^-$ at 4.178-4.226 GeV
¹ Assuming $B(f_0 \rightarrow \pi^0 \pi^0) = 1/3$ via the isospin limit, this result implies $B(D_s^+ \rightarrow f_0(980) e^+ \nu_e) = (2.4 \pm 0.4) \times 10^{-3}$.				

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.13 ± 0.03 ± 0.01	42	¹ HIETALA	15	Uses CLEO data
0.20 ± 0.03 ± 0.01	44	ECKLUND	09 CLEO	See HIETALA 15
0.13 ± 0.04 ± 0.01	13	YELTON	09 CLEO	See ECKLUND 09
¹ HIETALA 15 uses a tighter cut on the reconstructed $\pi^+ \pi^-$ mass (± 60 MeV around the f^0) than ECKLUND 09. It finds that applying the same tight cut to both analyses gives consistent results.				

$\Gamma(a_0(980)^0 e^+ \nu_e, a_0(980)^0 \rightarrow \pi^0 \eta)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2 × 10⁻⁴	90	ABLIKIM	21Y BES3	$e^+ e^-$ at 4.178-4.226 GeV

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.4 × 10⁻⁵	90	ABLIKIM	22BH BES3	6.32 fb ⁻¹ of $e^+ e^-$ at 4.178-4.226 GeV

————— **Hadronic modes with a $K\bar{K}$ pair** —————

$\Gamma(K^+ K_S^0)/\Gamma_{\text{total}}$ **Γ_{40}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.450 ± 0.035				OUR FIT
1.46 ± 0.05				OUR AVERAGE Error includes scale factor of 1.2.
$1.425 \pm 0.038 \pm 0.031$	1.8k	ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV
$1.52 \pm 0.05 \pm 0.03$		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.49 \pm 0.07 \pm 0.05$		¹ ALEXANDER 08	CLEO	See ONYISI 13

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(K^+ K_S^0)/\Gamma(K^+ K^- \pi^+)$ **Γ_{40}/Γ_{43}**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$27.55 \pm 0.18 \pm 0.50$	40k	ABLIKIM	20R BES3	$e^+ e^-$, 4178 ~ 4226 MeV

$\Gamma(K^+ K_L^0)/\Gamma_{\text{total}}$ **Γ_{41}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.485 \pm 0.039 \pm 0.046$	2.3k	ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV

$\Gamma(K^+ \bar{K}^0)/\Gamma_{\text{total}}$ **Γ_{42}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.95 \pm 0.11 \pm 0.09$	2.0k	¹ ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$

¹ ZUPANC 13 finds the \bar{K}^0 from its missing-mass squared, not from $K_S^0 \rightarrow \pi^+ \pi^-$.

The DCS ($D_S^+ \rightarrow K^+ K^0$) contribution to this fraction is estimated to be an order of magnitude below the statistical uncertainty.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$ **Γ_{43}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.37 ± 0.10				OUR FIT Error includes scale factor of 1.1.
5.45 ± 0.11				OUR AVERAGE Error includes scale factor of 1.1.
$5.47 \pm 0.08 \pm 0.13$	5.1k	ABLIKIM	21AE BES3	$e^+ e^-$ at 4.178 GeV
$5.55 \pm 0.14 \pm 0.13$		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
$5.06 \pm 0.15 \pm 0.21$	4.1k	ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$
$5.78 \pm 0.20 \pm 0.30$		DEL-AMO-SA..10J	BABR	$e^+ e^-$, 10.58 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$5.50 \pm 0.23 \pm 0.16$		¹ ALEXANDER 08	CLEO	See ONYISI 13

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\phi \pi^+)/\Gamma_{\text{total}}$ **Γ_{44}/Γ**

The results here are model-independent. For earlier, model-dependent results, see our PDG 06 edition. We decouple the $D_S^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_S^+ \rightarrow \phi \pi^+, \phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis

of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.4 OUR AVERAGE				
4.62 ± 0.36 ± 0.51		¹ AUBERT	06N BABR	$e^+ e^-$ at $\Upsilon(4S)$
4.81 ± 0.52 ± 0.38	212 ± 19	² AUBERT	05V BABR	$e^+ e^- \approx \Upsilon(4S)$
3.59 ± 0.77 ± 0.48		³ ARTUSO	96 CLE2	$e^+ e^-$ at $\Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.9 ^{+5.1} _{-1.9} ^{+1.8} _{-1.1}		⁴ BAI	95C BES	$e^+ e^-$ 4.03 GeV

¹ This AUBERT 06N measurement uses $\bar{B}^0 \rightarrow D_s^{(*)-} D^{(*)+}$ and $B^- \rightarrow D_s^{(*)-} D^{(*)0}$ decays, including some from other papers. However, the result is independent of AUBERT 05V.

² AUBERT 05V uses the ratio of $B^0 \rightarrow D^{*-} D_s^{*+}$ events seen in two different ways, in both of which the $D^{*-} \rightarrow \bar{D}^0 \pi^-$ decay is fully reconstructed: (1) The $D_s^{*+} \rightarrow D_s^+ \gamma$, $D_s^+ \rightarrow \phi \pi^+$ decay is fully reconstructed. (2) The number of events in the D_s^+ peak in the missing mass spectrum against the $D^{*-} \gamma$ is measured.

³ ARTUSO 96 uses partially reconstructed $\bar{B}^0 \rightarrow D^{*+} D_s^{*-}$ decays to get a model-independent value for $\Gamma(D_s^- \rightarrow \phi \pi^-) / \Gamma(D^0 \rightarrow K^- \pi^+)$ of $0.92 \pm 0.20 \pm 0.11$.

⁴ BAI 95C uses $e^+ e^- \rightarrow D_s^+ D_s^-$ events in which one or both of the D_s^\pm are observed to obtain the first model-independent measurement of the $D_s^+ \rightarrow \phi \pi^+$ branching fraction, without assumptions about $\sigma(D_s^\pm)$. However, with only two “doubly-tagged” events, the statistical error is very large.

$\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{45} / \Gamma_{43}$

This is the “fit fraction” from the Dalitz-plot analysis. We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
41.2 ± 0.7 OUR AVERAGE				
40.5 ± 0.7 ± 0.9	18.6k	ABLIKIM	21AE BES3	$e^+ e^-$ at 4.178 GeV
41.4 ± 0.8 ± 0.5		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
42.2 ± 1.6 ± 0.3		MITCHELL	09A CLEO	Dalitz fit, 12k evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
39.6 ± 3.3 ± 4.7		FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(K^+ \bar{K}^*(892)^0) / \Gamma(K^*(892)^+ \bar{K}^0)$ $\Gamma_{46} / \Gamma_{61}$

VALUE	DOCUMENT ID	TECN	COMMENT
2.35 ^{+0.42} _{-0.23} ± 0.10	ABLIKIM	22AH BES3	Dalitz plot fit to 990 $D_s^\pm \rightarrow K^\pm K_S \pi^0$ evts

$\Gamma(K^+\bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{47}/Γ_{43}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
47.9±0.6 OUR AVERAGE				
48.3±0.9±0.6	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
47.9±0.5±0.5		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
47.4±1.5±0.4		MITCHELL	09A CLEO	Dalitz fit, 12k evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
47.8±4.6±4.0		FRABETTI	95B E687	Dalitz fit, 701 evts


$\Gamma(K^+\bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K_S^0\pi^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{48}/Γ_{56}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
32.7±2.2±1.9	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{49}/Γ_{43}

This is the "fit fraction" from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(980)\pi$ and $D_s^+ \rightarrow a_0(980)\pi$ which are indistinguishable in such an analysis.


VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
20.6±3.5 OUR AVERAGE		Error includes scale factor of 2.5. See the ideogram below.		
19.3±1.7±2.0	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
16.4±0.7±2.0		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
28.2±1.9±1.8		MITCHELL	09A CLEO	Dalitz fit, 12k evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
11.0±3.5±2.6		FRABETTI	95B E687	Dalitz fit, 701 evts



$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{50}/Γ_{43}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3±0.5 OUR AVERAGE		Error includes scale factor of 2.8. See the ideogram below.		
1.2±0.4±0.2	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
1.1±0.1±0.2		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
4.3±0.6±0.5		MITCHELL	09A CLEO	Dalitz fit, 12k evts



$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{51}/Γ_{43}

This is the "fit fraction" from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(1710)\pi$ and $D_s^+ \rightarrow a_0(1710)\pi$ which are indistinguishable in such an analysis.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3±0.5 OUR AVERAGE		Error includes scale factor of 3.8.		
1.9±0.4±0.6	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
1.1±0.1±0.1		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
3.4±0.5±0.3		MITCHELL	09A CLEO	Dalitz fit, 12k evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.4±2.3±3.5		FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(a_0(980)^+\pi^0, a_0^+ \rightarrow K^+K_S^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{52}/Γ_{56}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.7±1.7±1.8	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(a_0(1710)^+\pi^0, a_0^+ \rightarrow K^+ K_S^0)/\Gamma(K^+ K_S^0 \pi^0)$ Γ_{53}/Γ_{56}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
23.6±3.4±2.0	¹ ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

¹ ABLIKIM 22AH observe an a_0 -like state with mass $m_{a_0} = 1.817 \pm 0.008 \pm 0.020$ GeV, and name the intermediate resonance $a_0(1817)$. We interpret this as the $a_0(1710)$ observed by LEES 21A.

$\Gamma(K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+)$ Γ_{54}/Γ_{43}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±0.5 OUR AVERAGE				

3.0±0.6±0.5 18.6k ABLIKIM 21AE BES3 $e^+ e^-$ at 4.178 GeV

2.4±0.3±1.0 DEL-AMO-SA..11G BABR Dalitz fit, 96k evts

3.9±0.5±0.5 MITCHELL 09A CLEO Dalitz fit, 12k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.3±3.2±3.2 FRABETTI 95B E687 Dalitz fit, 701 evts

$\Gamma(K^+ \bar{K}^*(1410)^0, \bar{K}_0^* \rightarrow K_S^0 \pi^0)/\Gamma(K^+ K_S^0 \pi^0)$ Γ_{55}/Γ_{56}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
6.0±1.4±1.3	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(K^+ K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.07 OUR AVERAGE				

1.46±0.06±0.05 990 ABLIKIM 22AH BES3 $e^+ e^-$ at 4.178-4.226 GeV

1.52±0.09±0.20 ONYISI 13 CLEO $e^+ e^-$ at 4.17 GeV

$\Gamma(2K_S^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.71±0.04 OUR AVERAGE				Error includes scale factor of 1.3.

0.68±0.04±0.01 370 ABLIKIM 22F BES3 $e^+ e^-$ at 4.178-4.226 GeV

0.77±0.05±0.03 ONYISI 13 CLEO $e^+ e^-$ at 4.17 GeV

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{58}/Γ

This is the "fit fraction" from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(980)\pi$ and $D_s^+ \rightarrow a_0(980)\pi$ which are indistinguishable in such an analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.8 × 10⁻⁴	90	¹ ABLIKIM	22F BES3	Dalitz plot fit

¹ Based on isospin considerations, the authors interpret the suppression in the observed rate of this mode compared to $D_s^+ \rightarrow f_0(980)\pi^+, f_0 \rightarrow K^+ K^-$ as likely due to the destructive interference between $a_0(980)$ and $f_0(980)$ in decays to $K_S^0 K_S^0$.

$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K_S^0 K_S^0)/\Gamma(2K_S^0 \pi^+)$ Γ_{59}/Γ_{57}

This is the "fit fraction" from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(1710)\pi$ and $D_s^+ \rightarrow a_0(1700)\pi$ which are indistinguishable in such an analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
46.3±4.0±1.2	ABLIKIM	22F BES3	Dalitz plot fit, 400 evts

$\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{61}/Γ_{44}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
1.20±0.21±0.13	CHEN	89	CLEO e^+e^- 10 GeV

$\Gamma(K^*(892)^+K_S^0)/\Gamma(2K_S^0\pi^+)$ Γ_{62}/Γ_{57}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
43.5±3.9±0.5	ABLIKIM	22F	BES3 Dalitz plot fit, 400 evts

$\Gamma(K^*(892)^+K_S^0, K^{*+} \rightarrow K^+\pi^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{63}/Γ_{56}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
13.9±1.7±1.3	ABLIKIM	22AH	BES3 Dalitz plot fit, 990 evts

$\Gamma(K^+K^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.50±0.24 OUR FIT				Error includes scale factor of 1.3.
5.51±0.28 OUR AVERAGE				Error includes scale factor of 1.5.

5.42±0.10±0.17	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV
6.37±0.21±0.56		ONYISI	13	CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.65±0.29±0.40		² ALEXANDER	08	CLEO	See ONYISI 13
----------------	--	------------------------	----	------	---------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_S^+ \rightarrow K^+K^-\pi^+\pi^0$ with 9 components.

² ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\phi\rho^+)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.59±0.15±0.30	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_S^+ \rightarrow K^+K^-\pi^+\pi^0$ with 9 components.

$\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$ Γ_{65}/Γ_{44}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.26^{+0.29}_{-0.40}	253	AVERY	92	CLE2 $e^+e^- \simeq 10.5$ GeV

$\Gamma(\bar{K}_1(1270)^0K^+, \bar{K}_1(1270)^0 \rightarrow K^-\rho^+)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.57±0.05±0.04	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_S^+ \rightarrow K^+K^-\pi^+\pi^0$ with 9 components.

$\Gamma(\bar{K}_1(1270)^0K^+, \bar{K}_1(1270)^0 \rightarrow K^*(892)\pi)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.31±0.18±0.18	3k	^{1,2} ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_S^+ \rightarrow K^+K^-\pi^+\pi^0$ with 9 components.

² $\bar{K}_1(1270)^0 \rightarrow K^*(892)\pi$ denotes a sum over $\bar{K}(892)^0\pi^0$ and $K(892)^-\pi^+$ final states, which are assumed to have relative branching ratio 1/2, as per isospin.

$\Gamma(\bar{K}_1(1400)^0 K^+, \bar{K}_1(1400)^0 \rightarrow K^*(892)\pi)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.98±0.27±0.32	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ $\bar{K}_1(1400)^0 \rightarrow K^*(892)\pi$ denotes a sum over $\bar{K}(892)^0\pi^0$ and $K(892)^-\pi^+$ final states, which are assumed to have relative branching ratio 1/2, as per isospin.

$\Gamma(a_0(980)^0 \rho^+, a_0(980)^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.19±0.03±0.03	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow K^*(892)^\mp K^\pm)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.39±0.06±0.03	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.04±0.01±0.01	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(\eta(1475)\pi^+, \eta(1475) \rightarrow a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.07±0.02±0.02	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(K_S^0 K^- 2\pi^+)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.53±0.08 OUR FIT	Error includes scale factor of 1.5.			
--------------------------	-------------------------------------	--	--	--

1.53±0.11 OUR AVERAGE	Error includes scale factor of 1.8.			
------------------------------	-------------------------------------	--	--	--

1.46±0.05±0.05	1.3k	ABLIKIM	21K BES3	e^+e^- at 4.178–4.226 GeV
----------------	------	---------	----------	-----------------------------

1.69±0.07±0.08		ONYISI	13 CLEO	e^+e^- at 4.17 GeV
----------------	--	--------	---------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.64±0.10±0.07		¹ ALEXANDER 08	CLEO	See ONYISI 13
----------------	--	---------------------------	------	---------------

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.64±0.23±0.27	3k	¹ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV
-----------------------	----	----------------------	----------	-----------------------------

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{74}/Γ_{44}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.6±0.4±0.4	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV
--------------------	----------	---------	--------------------------

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{74} / \Gamma_{73}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
40.6 ± 2.9 ± 4.9	1.3k	^{1,2} ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ Predominantly S -wave, with a significant D -wave component.

² $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.

$\Gamma(\eta(1475) K_S^0, \eta \rightarrow K^*(892)^0 \pi^+, K^{*0} \rightarrow K^- \pi^+) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{75} / \Gamma_{73}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.6 ± 0.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.

$\Gamma(\eta(1475) \pi^+, \eta \rightarrow \bar{K}^*(892)^+ K^-, \bar{K}^{*+} \rightarrow K_S^0 \pi^+) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{76} / \Gamma_{73}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.6 ± 0.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.

$\Gamma(\eta(1475) \pi^+, \eta \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{77} / \Gamma_{73}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.8 ± 2.6 ± 5.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.

$\Gamma(f_1(1285) \pi^+, f_1 \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{78} / \Gamma_{73}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.5 ± 0.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.

$\Gamma(K^+ K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{79} / Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.95 ± 0.08 OUR FIT	Error includes scale factor of 1.1.		
1.03 ± 0.06 ± 0.08	ONYISI	13	CLEO $e^+ e^-$ at 4.17 GeV

$\Gamma(K^+ K_S^0 \pi^+ \pi^-) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{79} / \Gamma_{73}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.62 ± 0.05 OUR FIT				
0.586 ± 0.052 ± 0.043	476	LINK	01c FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- 2\pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{80} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.60 ± 0.47 ± 0.38	309	ABLIKIM	22AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

$\Gamma(K^+ K^- 2\pi^+ \pi^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{80} / \Gamma_{43}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.160 ± 0.027 OUR AVERAGE				
0.150 ± 0.019 ± 0.025	240	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.188 ± 0.036 ± 0.040	75	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\phi 2\pi^+ \pi^-)/\Gamma(\phi \pi^+)$ Γ_{81}/Γ_{44}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.269±0.027 OUR AVERAGE				
0.249±0.024±0.021	136	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV
0.28 ±0.06 ±0.01	40	FRABETTI	97C E687	$\gamma Be, \bar{E}_\gamma \approx 200$ GeV
0.58 ±0.21 ±0.10	21	FRABETTI	92 E687	γBe
0.42 ±0.13 ±0.07	19	ANJOS	88 E691	Photoproduction
1.11 ±0.37 ±0.28	62	ALBRECHT	85D ARG	$e^+ e^- 10$ GeV

$\Gamma(\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{82}/Γ_{80}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.75±0.06±0.04	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+ K^-, a_1^+ \rightarrow \rho^0 \pi^+)/\Gamma(K^+ K^- \pi^+)$ Γ_{83}/Γ_{43}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.137±0.019±0.011	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+ K^-, a_1^+ \rightarrow \rho^0 \pi^+)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{83}/Γ_{80}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.781±0.029±0.016	235	ABLIKIM	22AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

$\Gamma(\phi 2\pi^+ \pi^- \text{ non-}\rho, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{84}/Γ_{80}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21±0.05±0.06	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- \rho^0 \pi^+ \text{ non-}\phi)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{85}/Γ_{80}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.03	90	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- 2\pi^+ \pi^- \text{ nonresonant})/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{86}/Γ_{80}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.15 ±0.06 OUR AVERAGE				
0.218±0.029±0.08	235	ABLIKIM	22AB BES3	$e^+ e^-$ at 4.178–4.226 GeV
0.10 ±0.06 ±0.05		LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(2K_S^0 2\pi^+ \pi^-)/\Gamma(K_S^0 K^- 2\pi^+)$ Γ_{87}/Γ_{73}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.051±0.015±0.015	37 ± 10	LINK	04D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV

———— Pionic modes ————

$\Gamma(\pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{88}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2 × 10⁻⁴	90	¹ GUAN	21 BELL	$e^+ e^- \approx \Upsilon(4, 5S)$

¹ Uses $B(D_s^+ \rightarrow \pi^+ \phi, \phi \rightarrow K^+ K^-) = (2.24 \pm 0.08)\%$.

$\Gamma(\pi^+\pi^0)/\Gamma(K^+K_S^0)$ Γ_{88}/Γ_{40}

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	MENDEZ	10	CLEO e^+e^- at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4.1	90	ADAMS	07A	CLEO See MENDEZ 10

$\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.08±0.04 OUR FIT			
1.11±0.04±0.04	ONYISI	13	CLEO e^+e^- at 4.17 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.11±0.07±0.04	¹ ALEXANDER	08	CLEO See ONYISI 13
¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.			

$\Gamma(2\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{89}/Γ_{43}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.201±0.007 OUR FIT				
0.199±0.004±0.009	≈ 10.5k	AUBERT	090	BABR e^+e^- ≈ 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.265±0.041±0.031	98	FRABETTI	97D	E687 γ Be ≈ 200 GeV

$\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$ Γ_{90}/Γ_{89}

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
1.1±0.6 OUR AVERAGE				
0.9±0.4±0.5		ABLIKIM	22BI	BES3 Dalitz fit, 11.1k events
1.8±0.5±1.0		AUBERT	090	BABR Dalitz fit, ≈ 10.5k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		LINK	04	FOCS Dalitz fit, 1475 ± 50 evts
5.8±2.3±3.7		AITALA	01A	E791 Dalitz fit, 848 evts
<7.3	90	FRABETTI	97D	E687 γ Be ≈ 200 GeV

$\Gamma(\pi^+(\pi^+\pi^-)_{S\text{-wave}})/\Gamma(2\pi^+\pi^-)$ Γ_{91}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis. See also KLEMPT 08, which uses 568 $D_S^+ \rightarrow 3\pi$ decays (over 280 background events) from FNAL E791 to study various parametrizations of the decay amplitudes. The emphasis there is more on S -wave $\pi\pi$ decay products — 20 different solutions are given — than on D_S^+ fit fractions.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
83.9 ±1.2 OUR AVERAGE			
84.2 ±0.8 ±1.2	¹ ABLIKIM	22BI	BES3 Dalitz fit, 11.1k events
83.0 ±0.9 ±1.9	¹ AUBERT	090	BABR Dalitz fit, ≈ 10.5k evts
87.04±5.60±4.38	² LINK	04	FOCS Dalitz fit, 1475 ± 50 evts

¹AUBERT 090 and ABLIKIM 22BI give the amplitude and phase of the $\pi^+\pi^-$ S -wave in 29 $\pi^+\pi^-$ invariant-mass bins.

²LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full $\pi-\pi$ S -wave isoscalar scattering amplitude to describe the $\pi^+\pi^-$ S -wave component of the $\pi^+\pi^+\pi^-$ state. The fit fraction given above is a sum over five f_0 mesons, the $f_0(980)$, $f_0(1300)$, $f_0(1200-1600)$, $f_0(1500)$, and $f_0(1750)$. See LINK 04 for details and discussion.

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{92}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.565 \pm 0.043 \pm 0.047$	AITALA	01A E791	Dalitz fit, 848 evts
$1.074 \pm 0.140 \pm 0.043$	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{93}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.324 \pm 0.077 \pm 0.017$	AITALA	01A E791	Dalitz fit, 848 evts

$\Gamma(f_0(1500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{94}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.274 \pm 0.114 \pm 0.019$	¹ FRABETTI	97D E687	γ Be \approx 200 GeV
¹ FRABETTI 97D calls this mode $S(1475)\pi^+$, but finds the mass and width of this $S(1475)$ to be in excellent agreement with those of the $f_0(1500)$.			

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{95}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.3 \pm 1.1 OUR AVERAGE			
$10.5 \pm 0.8 \pm 1.1$	ABLIKIM	22BI BES3	Dalitz fit, 11.1k events
$10.1 \pm 1.5 \pm 1.1$	AUBERT	09O BABR	Dalitz fit, \approx 10.5k evts
$9.74 \pm 4.49 \pm 2.94$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$19.7 \pm 3.3 \pm 0.6$	AITALA	01A E791	Dalitz fit, 848 evts
$12.3 \pm 5.6 \pm 1.8$	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(\rho(1450)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{96}/Γ_{89}

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.5 \pm 0.6 OUR AVERAGE			
$1.3 \pm 0.4 \pm 0.5$	ABLIKIM	22BI BES3	Dalitz fit, 11.1k events
$2.3 \pm 0.8 \pm 1.7$	AUBERT	09O BABR	Dalitz fit, \approx 10.5k evts
$6.56 \pm 3.43 \pm 4.40$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$4.4 \pm 2.1 \pm 0.2$	AITALA	01A E791	Dalitz fit, 848 evts

$\Gamma(\pi^+2\pi^0)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.52 \pm 0.05 OUR AVERAGE Error includes scale factor of 1.1.				
$0.50 \pm 0.04 \pm 0.02$	590	ABLIKIM	22Z BES3	e^+e^- at 4.178–4.226 GeV
$0.65 \pm 0.13 \pm 0.03$	72 ± 16	NAIK	09A CLEO	e^+e^- at 4170 MeV

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ Γ_{98}/Γ_{97}

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
55.4±6.8±7.3	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ Γ_{99}/Γ_{97}

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
25.5±5.1±9.3	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ Γ_{100}/Γ_{97}

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
9.7±2.9±6.0	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{101}/Γ_{44}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
•••				We do not use the following data for averages, fits, limits, etc. •••
<3.3	90	ANJOS	89E E691	Photoproduction

$\Gamma(\eta\pi^+)/\Gamma_{total}$ Γ_{102}/Γ

Unseen decay modes of the η are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.09 OUR FIT				Error includes scale factor of 1.1.
1.71±0.08 OUR AVERAGE				
1.67±0.08±0.06		ONYISI	13	CLEO e^+e^- at 4.17 GeV
1.82±0.14±0.07	0.8k	ZUPANC	13	BELL e^+e^- at $\Upsilon(4S), \Upsilon(5S)$
•••				We do not use the following data for averages, fits, limits, etc. •••
1.58±0.11±0.18		¹ ALEXANDER	08	CLEO See ONYISI 13
		¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.		

$\Gamma(\eta\pi^+)/\Gamma(K^+K_S^0)$ Γ_{102}/Γ_{40}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.15 ±0.07 OUR FIT				Error includes scale factor of 1.1.
•••				We do not use the following data for averages, fits, limits, etc. •••
1.236±0.043±0.063	2587 ± 89	MENDEZ	10	CLEO See ONYISI 13

$\Gamma(\eta\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{102}/Γ_{43}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
31.94±0.33±0.49	19.5k	ABLIKIM	20R	BES3 e^+e^- , 4178 ~ 4226 MeV

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{102}/Γ_{44}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
•••				We do not use the following data for averages, fits, limits, etc. •••
0.48±0.03±0.04	920	JESSOP	98	CLE2 $e^+e^- \approx \Upsilon(4S)$
0.54±0.09±0.06	165	ALEXANDER	92	CLE2 See JESSOP 98

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+, \phi \rightarrow K^+ K^-)$					Γ_{102}/Γ_{45}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
84.80±0.47±2.64	22k	GUAN	21 BELL	$e^+ e^- \approx \Upsilon(4, 5S)$	

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$					Γ_{103}/Γ
Unseen decay modes of the ω are included.					
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.192±0.030 OUR FIT					
0.181±0.032 OUR AVERAGE					
0.177±0.032±0.013	65 ± 12	ABLIKIM	19AH BES3	$e^+ e^-$ at 4.178 GeV	
0.21 ±0.09 ±0.01	6 ± 2.4	GE	09A CLEO	$e^+ e^-$ at 4170 MeV	

$\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$					$\Gamma_{103}/\Gamma_{102}$
Unseen decay modes of the resonances are included.					
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.115±0.018 OUR FIT					
0.16 ±0.04 ±0.03		BALEST	97 CLE2	$e^+ e^- \approx \Upsilon(4S)$	

$\Gamma(3\pi^+ 2\pi^-)/\Gamma(K^+ K^- \pi^+)$					Γ_{104}/Γ_{43}
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.146±0.014 OUR AVERAGE					
0.145±0.011±0.010	671	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV	
0.158±0.042±0.031	37	FRABETTI	97C E687	$\gamma Be, \bar{E}_\gamma \approx 200$ GeV	

$\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$					Γ_{106}/Γ
Unseen decay modes of the η are included.					
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.9±0.6±0.5	328 ± 22	NAIK	09A CLEO	$\eta \rightarrow 2\gamma$	

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$					Γ_{106}/Γ_{44}
Unseen decay modes of the resonances are included.					
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.98±0.20±0.39	447	JESSOP	98 CLE2	$e^+ e^- \approx \Upsilon(4S)$	
2.86±0.38 ^{+0.36} _{-0.38}	217	AVERY	92 CLE2	See JESSOP 98	

$\Gamma(\eta\rho^+)/\Gamma(\eta\pi^+\pi^0)$					$\Gamma_{106}/\Gamma_{107}$
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
78.3±5.0±2.1	1.2k	ABLIKIM	19BE BES3	$\eta\pi^+\pi^0$ amplitude analysis	

$\Gamma(\eta\pi^+\pi^0)/\Gamma_{\text{total}}$					Γ_{107}/Γ
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
9.5 ±0.5 OUR AVERAGE					
9.50±0.28±0.41	2.6k	ABLIKIM	19BE BES3	$e^+ e^-$ at 4.178 GeV	
9.2 ±0.4 ±1.1		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV	

$\Gamma(\eta(\pi^+\pi^0)_{P\text{-wave}})/\Gamma(\eta\pi^+\pi^0)$					$\Gamma_{108}/\Gamma_{107}$
<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.4±2.1±2.5	1.2k	ABLIKIM	19BE BES3	$\eta\pi^+\pi^0$ amplitude analysis	

$\Gamma(a_0(980)^+ \pi^0, a_0(980)^+ \rightarrow \eta \pi^+ \pi^0) / \Gamma(\eta \pi^+ \pi^0)$ $\Gamma_{109} / \Gamma_{107}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
23.2 ± 2.3 ± 3.3	1.2k	¹ ABLIKIM	19BE BES3	$\eta \pi^+ \pi^0$ amplitude analysis

¹ Coherent sum of $D_s^+ \rightarrow a_0^+ \pi^0 \rightarrow \eta \pi^+ \pi^0$ and $D_s^+ \rightarrow a_0^0 \pi^+ \rightarrow \eta \pi^+ \pi^0$. ABLIKIM 19BE find $a_0(980)^0 - f(980)$ mixing effects negligibly small in this $D_s^+ \rightarrow \eta \pi^+ \pi^0$ Dalitz plot analysis.

$\Gamma(\omega \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{110} / Γ

Unseen decay modes of the ω are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.78 ± 0.65 ± 0.25	34 ± 7.9	GE	09A CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(2\pi^+ \pi^- \eta) / \Gamma_{\text{total}}$ Γ_{111} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.12 ± 0.13 ± 0.09	2.1k	ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

$\Gamma(a_1(1260)^+ \eta, a_1^+ \rightarrow \rho(770)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^- \eta)$ $\Gamma_{112} / \Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
55.4 ± 3.9 ± 2.0	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ amplitude analysis with 11 components.

$\Gamma(a_1(1260)^+ \eta, a_1^+ \rightarrow f_0(500) \pi^+, f_0 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^- \eta)$ $\Gamma_{113} / \Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
8.1 ± 1.9 ± 2.1	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ amplitude analysis with 11 components.

$\Gamma(a_0(980)^+ \rho(770)^0, a_0^+ \rightarrow \eta \pi^+) / \Gamma(2\pi^+ \pi^- \eta)$ $\Gamma_{114} / \Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
6.7 ± 2.5 ± 1.5	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ amplitude analysis with 11 components.

$\Gamma(\eta(1405) \pi^+, \eta(1405) \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow \eta \pi^-) / \Gamma(2\pi^+ \pi^- \eta)$ $\Gamma_{115} / \Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.2 ± 0.1	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ amplitude analysis with 11 components.

$\Gamma(\eta(1405) \pi^+, \eta(1405) \rightarrow a_0(980)^+ \pi^-, a_0^+ \rightarrow \eta \pi^+) / \Gamma(2\pi^+ \pi^- \eta)$ $\Gamma_{116} / \Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.2 ± 0.1	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ $D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ amplitude analysis with 11 components.

$\Gamma(f_1(1420)\pi^+, f_1 \rightarrow a_0(980)^-\pi^+, a_0^- \rightarrow \eta\pi^-)/\Gamma(2\pi^+\pi^-\eta)$ $\Gamma_{117}/\Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.9±0.5±0.3	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$\Gamma(f_1(1420)\pi^+, f_1 \rightarrow a_0(980)^+\pi^-, a_0^+ \rightarrow \eta\pi^+)/\Gamma(2\pi^+\pi^-\eta)$ $\Gamma_{118}/\Gamma_{111}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.7±0.5±0.3	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$\Gamma(3\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.049^{+0.033}_{-0.030}	BARLAG	92C ACCM	π^- 230 GeV
--	--------	----------	-----------------

$\Gamma(\omega 2\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{120}/Γ

Unseen decay modes of the ω are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.58±0.45±0.09	29 ± 8.2	GE	09A CLEO	e^+e^- at 4170 MeV
-----------------------	----------	----	----------	----------------------

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$ Γ_{121}/Γ

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

3.94±0.15±0.20	ONYISI	13 CLEO	e^+e^- at 4.17 GeV
-----------------------	--------	---------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.77±0.25±0.30	¹ ALEXANDER	08 CLEO	See ONYISI 13
----------------	------------------------	---------	---------------

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+K_S^0)$ Γ_{121}/Γ_{40}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.654±0.088±0.139	1436 ± 47	MENDEZ	10 CLEO	See ONYISI 13
-------------------	-----------	--------	---------	---------------

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{121}/Γ_{43}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

69.4±0.8±3.8	9.9k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV
---------------------	------	---------	----------	----------------------------

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ Γ_{121}/Γ_{44}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.03±0.06±0.07	537	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
----------------	-----	--------	---------	-------------------------------

1.20±0.15±0.11	281	ALEXANDER	92 CLE2	See JESSOP 98
----------------	-----	-----------	---------	---------------

2.5 ± 1.0 ^{+1.5} _{-0.4}	22	ALVAREZ	91 NA14	Photoproduction
---	----	---------	---------	-----------------

2.5 ± 0.5 ± 0.3	215	ALBRECHT	90D ARG	$e^+e^- \approx 10.4$ GeV
-----------------	-----	----------	---------	---------------------------

$\Gamma(\omega\eta\pi^+)/\Gamma_{\text{total}}$ Γ_{123}/Γ

Unseen decay modes of the ω and η are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.13 \times 10^{-2}$	90	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(\eta'(958)\rho^+)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$5.8 \pm 1.4 \pm 0.4$	ABLIKIM 15Z	BES3	482 pb^{-1} , 4009 MeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ Γ_{124}/Γ_{44}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.78 \pm 0.28 \pm 0.30$	137	¹ JESSOP	98	CLE2 $e^+e^- \approx \gamma(4S)$
$3.44 \pm 0.62^{+0.44}_{-0.46}$	68	AVERY	92	CLE2 See JESSOP 98

¹ This JESSOP 98 fraction, when combined with other η' fractions, greatly overshoots the inclusive η' fraction. See the measurement just above, which fits nicely.

$\Gamma(\eta'(958)\rho^+)/\Gamma(\eta'(958)\pi^+\pi^0)$ $\Gamma_{124}/\Gamma_{125}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≈ 1	395	¹ ABLIKIM	22AA	BES3 e^+e^- at 4.178–4.226 GeV
-------------	-----	----------------------	------	----------------------------------

¹ Result of an amplitude analysis of $D_s^+ \rightarrow \pi^+\pi^0\eta'$ which found that $D_s^+ \rightarrow \rho^+\eta'$ is the dominant decay mode, with other contributions negligible. No uncertainty is assigned to this 100% fit fraction; however, the fit fractions of non-resonant contributions are shown to be below 1%.

$\Gamma(\eta'(958)\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

6.08 ± 0.29 OUR AVERAGE

$6.15 \pm 0.25 \pm 0.18$	837	¹ ABLIKIM	22AA	BES3 e^+e^- at 4.178–4.226 GeV
$5.6 \pm 0.5 \pm 0.6$		ONYISI	13	CLEO e^+e^- at 4.17 GeV

¹ An amplitude analysis in the same publication finds that $D_s^+ \rightarrow \rho^+\eta'$ is the only statistically significant contribution to this decay.

$\Gamma(\eta'(958)\pi^+\pi^0 \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<5.1 \times 10^{-2}$	90	ABLIKIM	15Z	BES3 482 pb^{-1} , 4009 MeV
-----------------------	----	---------	-----	---------------------------------------

———— Modes with one or three K's ————

$\Gamma(K^+\pi^0)/\Gamma(K^+K_S^0)$ Γ_{127}/Γ_{40}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4.2 \pm 1.4 \pm 0.2$	202 ± 70	MENDEZ	10	CLEO e^+e^- at 4170 MeV
-----------------------	--------------	--------	----	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.5 \pm 1.3 \pm 0.7$	141 ± 34	ADAMS	07A	CLEO See MENDEZ 10
-----------------------	--------------	-------	-----	--------------------

$\Gamma(K^+\pi^0)/\Gamma(K^+K^-\pi^+)$ Γ_{127}/Γ_{43}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
13.73±0.90±0.33	2.3k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV

$\Gamma(K^+\pi^0)/\Gamma(\phi\pi^+, \phi \rightarrow K^+K^-)$ Γ_{127}/Γ_{45}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
3.28±0.23±0.13	12k	GUAN	21 BELL	$e^+e^- \approx \Upsilon(4,5S)$

$\Gamma(K_S^0\pi^+)/\Gamma(K^+K_S^0)$ Γ_{128}/Γ_{40}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.12±0.28 OUR AVERAGE				
8.5 ±0.7 ±0.2	393 ± 33	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
8.03±0.24±0.19	17.6k±481	WON	09 BELL	e^+e^- at $\Upsilon(4S)$
10.4 ±2.4 ±1.4	113 ± 26	LINK	08 FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2 ±0.9 ±0.2	206 ± 22	ADAMS	07A CLEO	See MENDEZ 10
---------------	----------	-------	----------	---------------

$\Gamma(K_S^0\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{128}/Γ_{43}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
20.35±0.62±0.42	2.7k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV

$\Gamma(K^+\eta)/\Gamma(K^+K_S^0)$ Γ_{129}/Γ_{40}

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
11.8±2.2±0.6	222 ± 41	MENDEZ	10 CLEO	e^+e^- at 4170 MeV

$\Gamma(K^+\eta)/\Gamma(K^+K^-\pi^+)$ Γ_{129}/Γ_{43}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.97±0.18±0.06	1.8k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV

$\Gamma(K^+\eta)/\Gamma(\phi\pi^+, \phi \rightarrow K^+K^-)$ Γ_{129}/Γ_{45}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
7.81±0.22±0.24	14k	GUAN	21 BELL	$e^+e^- \approx \Upsilon(4,5S)$

$\Gamma(K^+\eta)/\Gamma(\eta\pi^+)$ $\Gamma_{129}/\Gamma_{102}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.9±1.5±0.4	113 ± 18	ADAMS	07A CLEO	See MENDEZ 10
-------------	----------	-------	----------	---------------

$\Gamma(K^+\omega)/\Gamma_{total}$ Γ_{130}/Γ

Unseen decay modes of the ω are included.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.9±1.5 OUR FIT					
8.7±2.4±0.8	29	¹ ABLIKIM	19AH BES3	e^+e^- at 4.178 GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24	90	GE	09A CLEO	e^+e^- at 4170 MeV
-----	----	----	----------	----------------------

¹Evidence for mode at 4.4σ .

$\Gamma(K^+\omega)/\Gamma(K^+\pi^+\pi^-\pi^0)$ $\Gamma_{130}/\Gamma_{144}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

10.3±1.5 OUR FIT

10.9±1.8±0.1 ¹ ABLIKIM 22BL BES3 PWA, 550 $D_s^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

¹ ABLIKIM 22BL reports $[\Gamma(D_s^+ \rightarrow K^+\omega)/\Gamma(D_s^+ \rightarrow K^+\pi^+\pi^-\pi^0)] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (9.7 \pm 1.5 \pm 0.6) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+\eta'(958))/\Gamma(K^+K_S^0)$ Γ_{131}/Γ_{40}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

11.8±3.6±0.7 56 ± 17 MENDEZ 10 CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\eta'(958))/\Gamma(K^+K^-\pi^+)$ Γ_{131}/Γ_{43}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.91±0.31±0.31 675 ABLIKIM 20R BES3 e^+e^- , 4178 ~ 4226 MeV

$\Gamma(K^+\eta'(958))/\Gamma(\eta'(958)\pi^+)$ $\Gamma_{131}/\Gamma_{121}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2±1.3±0.3 28 ± 9 ADAMS 07A CLEO See MENDEZ 10

$\Gamma(K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.20±0.19 OUR FIT

6.20±0.19 OUR AVERAGE

6.11±0.18±0.11 1.3k ABLIKIM 22AC BES3 e^+e^- at 4.178–4.226 GeV

6.54±0.33±0.25 ONYISI 13 CLEO e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.9 ± 0.5 ± 0.3 ¹ ALEXANDER 08 CLEO See ONYISI 13

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(K^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{132}/Γ_{43}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.115±0.004 OUR FIT Error includes scale factor of 1.1.

0.127±0.007±0.014 567 ± 31 LINK 04F FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-)$ $\Gamma_{133}/\Gamma_{132}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.35 ± 0.04 OUR AVERAGE

0.321±0.037±0.037 ABLIKIM 22AC BES3 Dalitz plot fit, 1.3k evts

0.388±0.053±0.026 LINK 04F FOCS Dalitz plot fit, 567 evts

$$\Gamma(K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{134} / \Gamma_{132}$$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.117 ± 0.028 OUR AVERAGE			
0.131 ± 0.031 ± 0.029	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.106 ± 0.035 ± 0.010	LINK	04F FOCS	Dalitz plot fit, 567 evts

$$\Gamma(K^+ f_0(500), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{135} / \Gamma_{132}$$

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
7.2 ± 2.1 ± 4.4	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts

$$\Gamma(K^+ f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{136} / \Gamma_{132}$$

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
4.5 ± 1.3 ± 1.2	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts

$$\Gamma(K^+ f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{137} / \Gamma_{132}$$

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
19.9 ± 2.9 ± 9.3	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts

$$\Gamma(K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{138} / \Gamma_{132}$$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.27 ± 0.04 OUR AVERAGE			Error includes scale factor of 2.0.
0.302 ± 0.018 ± 0.020	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.2164 ± 0.0321 ± 0.0114	LINK	04F FOCS	Dalitz plot fit, 567 evts

$$\Gamma(K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{139} / \Gamma_{132}$$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.07 OUR AVERAGE			Error includes scale factor of 2.7.
0.045 ± 0.021 ± 0.025	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.1882 ± 0.0403 ± 0.0122	LINK	04F FOCS	Dalitz plot fit, 567 evts

$$\Gamma(K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{140} / \Gamma_{132}$$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ± 0.05 OUR AVERAGE			Error includes scale factor of 1.7.
0.185 ± 0.025 ± 0.026	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.0765 ± 0.0500 ± 0.0170	LINK	04F FOCS	Dalitz plot fit, 567 evts

$$\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^-) \quad \Gamma_{141} / \Gamma_{132}$$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1588 ± 0.0492 ± 0.0153	LINK	04F FOCS	Dalitz fit, 567 evts

$$\Gamma(K^0 \pi^+ \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{142} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.06 OUR AVERAGE				
1.086 ± 0.060 ± 0.030	666	¹ ABLIKIM	21AB BES3	e ⁺ e ⁻ at 4.178–4.226 GeV
1.00 ± 0.18 ± 0.04	44	NAIK	09A CLEO	e ⁺ e ⁻ at 4170 MeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component, and measures $B(D_s^+ \rightarrow K_S^0 \pi^+ \pi^0) = (5.43 \pm 0.30 \pm 0.15) \times 10^{-3}$.

$\Gamma(K_S^0 2\pi^+ \pi^-) / \Gamma(K_S^0 K^- 2\pi^+)$					$\Gamma_{143} / \Gamma_{73}$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.18 ± 0.04 ± 0.05	179 ± 36	LINK	08	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV	

$\Gamma(K^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$					Γ_{144} / Γ
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
9.7 ± 0.6 OUR FIT					
9.75 ± 0.54 ± 0.17	776	ABLIKIM	22BL BES3	$e^+ e^-$ at 4.178–4.226 GeV	

$\Gamma(K^*(892)^0 \rho^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{145} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
40.5 ± 2.8 ± 1.5	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K^*(892)^+ \rho^0, K^{*+} \rightarrow K^+ \pi^0) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{146} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
4.3 ± 1.1 ± 0.6	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K_1(1270)^0 \pi^+, K_1^0 \rightarrow K^+ \rho^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{147} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
4.0 ± 1.2 ± 0.6	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^*(890)^+ \pi^-, K^{*+} \rightarrow K^+ \pi^0) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{148} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
5.6 ± 0.9 ± 0.2	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^*(890)^0 \pi^0, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{149} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
6.1 ± 0.9 ± 0.2	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K^+ a_1(1260)^0, a_1 \rightarrow \rho^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{150} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.9 ± 0.7 ± 0.9	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K^+ a_1(1260)^0, a_1 \rightarrow \rho^- \pi^+) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{151} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.9 ± 0.7 ± 0.9	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma(K^+ \pi^+ \pi^- \pi^0 \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^- \pi^0)$					$\Gamma_{152} / \Gamma_{144}$
<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
9.5 ± 2.2 ± 0.9	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$		

$\Gamma((K^+\pi^0)_{P\text{-wave}}\rho^0)/\Gamma(K^+\pi^+\pi^-\pi^0)$ $\Gamma_{153}/\Gamma_{144}$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
10.4 ± 2.0 ± 0.6		ABLIKIM	22BL BES3	PWA, 550 $D_s^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$\Gamma(K^+\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{154}/Γ

Unseen decay modes of the ω are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.82	90	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{155}/Γ

Unseen decay modes of the ω are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.54	90	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\omega\eta)/\Gamma_{\text{total}}$ Γ_{156}/Γ

Unseen decay modes of the ω and η are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.79	90	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(2K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{157}/Γ_{43}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.3 ± 0.2	748 ± 60	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.95 ± 2.12 ^{+2.24} _{-2.31}	31	LINK	02I	FOCS $\gamma A, \approx 180$ GeV

$\Gamma(\phi K^+, \phi \rightarrow K^+K^-)/\Gamma(2K^+K^-)$ $\Gamma_{158}/\Gamma_{157}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.08 ± 0.03	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(2K^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{159}/Γ_{43}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.371 ± 0.034 OUR AVERAGE				
2.372 ± 0.024 ± 0.025	67k	AAIJ	19G	LHCB pp at 8 TeV
2.3 ± 0.3 ± 0.2	356 ± 52	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$
2.29 ± 0.28 ± 0.12	281 ± 34	KO	09	BELL e^+e^- at $\Upsilon(4S)$
5.2 ± 1.7 ± 1.1	27 ± 9	LINK	05k	FOCS <0.78%, CL = 90%

$\Gamma(K^+K^*(892)^0, K^{*0} \rightarrow K^+\pi^-)/\Gamma(2K^+\pi^-)$ $\Gamma_{160}/\Gamma_{159}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.22 ± 0.15	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$

———— Baryon-antibaryon mode ————

$\Gamma(p\bar{n})/\Gamma_{\text{total}}$ Γ_{161}/Γ

This is the only baryonic mode allowed kinematically.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.22 ± 0.11 OUR AVERAGE				
1.21 ± 0.10 ± 0.05	193 ± 17	ABLIKIM	190BES3	e^+e^- , $E_{\text{cm}} = 4178$ MeV
1.30 ± 0.36 ^{+0.12} _{-0.16}	13.0 ± 3.6	ATHAR	08	CLEO e^+e^- , $E_{\text{cm}} \approx 4170$ MeV

$\Gamma(\rho\bar{\rho}e^+\nu_e)/\Gamma_{\text{total}}$					Γ_{162}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-4}$	90	ABLIKIM	19BD BES3	e^+e^- at 4178 MeV	

————— Rare or forbidden modes —————

$\Gamma(\pi^+e^+e^-)/\Gamma_{\text{total}}$					Γ_{163}/Γ
---	--	--	--	--	-----------------------

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.5 \times 10^{-6}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<13 \times 10^{-6}$	90	LEES	11G BABR	$e^+e^- \approx \Upsilon(4S)$
$< 2.2 \times 10^{-5}$	90	¹ RUBIN	10 CLEO	e^+e^- at 4170 MeV
$<27 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

¹This RUBIN 10 limit is for the e^+e^- mass in the continuum away from the $\phi(1020)$. See the next data block.

$\Gamma(\pi^+\phi, \phi \rightarrow e^+e^-)/\Gamma_{\text{total}}$					Γ_{164}/Γ
--	--	--	--	--	-----------------------

This is *not* a test for the $\Delta C = 1$ weak neutral current, but leads to the $\pi^+e^+e^-$ final state.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$(6_{-4}^{+8} \pm 1) \times 10^{-6}$	3	RUBIN	10 CLEO	e^+e^- at 4170 MeV

$\Gamma(\pi^+\mu^+\mu^-)/\Gamma_{\text{total}}$					Γ_{165}/Γ
---	--	--	--	--	-----------------------

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<4.1 \times 10^{-7}$	90	AAIJ	13AF LHCb	pp at 7 TeV
$<4.3 \times 10^{-5}$	90	LEES	11G BABR	$e^+e^- \approx \Upsilon(4S)$
$<2.6 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$<1.4 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<4.3 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^+e^+e^-)/\Gamma_{\text{total}}$					Γ_{166}/Γ
---	--	--	--	--	-----------------------

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-6}$	90	LEES	11G BABR	$e^+e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<4.9 \times 10^{-6}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
$<5.2 \times 10^{-5}$	90	RUBIN	10 CLEO	e^+e^- at 4170 MeV
$<1.6 \times 10^{-3}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^+\mu^+\mu^-)/\Gamma_{\text{total}}$					Γ_{167}/Γ
---	--	--	--	--	-----------------------

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 21 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 3.6 \times 10^{-5}$	90	LINK	03F	FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$< 1.4 \times 10^{-4}$	90	AITALA	99G	E791	$\pi^- N 500 \text{ GeV}$
$< 5.9 \times 10^{-4}$	90	KODAMA	95	E653	$\pi^- \text{ emulsion } 600 \text{ GeV}$

$\Gamma(K^*(892)^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{168} / Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-3}$	90	KODAMA	95	E653 $\pi^- \text{ emulsion } 600 \text{ GeV}$

$\Gamma(\pi^+ e^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{169} / Γ**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.1 \times 10^{-6}$	90	AAIJ	21T	LHCB $1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 12 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
-----------------------	----	------	-----	------	--------------------------------

$\Gamma(\pi^+ e^- \mu^+) / \Gamma_{\text{total}}$ **Γ_{170} / Γ**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.4 \times 10^{-7}$	90	AAIJ	21T	LHCB $1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 20 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
-----------------------	----	------	-----	------	--------------------------------

$\Gamma(K^+ e^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_{171} / Γ**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.9 \times 10^{-7}$	90	AAIJ	21T	LHCB $1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 14 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
-----------------------	----	------	-----	------	--------------------------------

$\Gamma(K^+ e^- \mu^+) / \Gamma_{\text{total}}$ **Γ_{172} / Γ**

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-7}$	90	AAIJ	21T	LHCB $1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 9.7 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
------------------------	----	------	-----	------	--------------------------------

$\Gamma(\pi^- 2e^+) / \Gamma_{\text{total}}$ **Γ_{173} / Γ**

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-6}$	90	AAIJ	21T	LHCB $1.6 \text{ fb}^{-1} pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 4.1 \times 10^{-6}$	90	LEES	11G	BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 1.8 \times 10^{-5}$	90	RUBIN	10	CLEO	$e^+ e^- \text{ at } 4170 \text{ MeV}$
$< 69 \times 10^{-5}$	90	AITALA	99G	E791	$\pi^- N 500 \text{ GeV}$

$\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{174}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.6 \times 10^{-8}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.2 \times 10^{-7}$	90	AAIJ	13AF LHCB	pp at 7 TeV
$<1.4 \times 10^{-5}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<2.9 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$<8.2 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<4.3 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{175}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.3 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<8.4 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<7.3 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^- 2e^+)/\Gamma_{\text{total}}$ **Γ_{176}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.7 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5.2 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<1.7 \times 10^{-5}$	90	RUBIN	10 CLEO	$e^+ e^-$ at 4170 MeV
$<63 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{177}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.6 \times 10^{-8}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.3 \times 10^{-5}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<1.3 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$<1.8 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<5.9 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{178}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.6 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<6.1 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<6.8 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$ Γ_{179}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-3}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

D_s^\pm Amplitude analyses

$D_s^+ \rightarrow K^+ K^- \pi^+$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $K^+ K^- \pi^+$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	18.6k	¹ ABLIKIM	21AE BES3	$e^+ e^-$ at 4.178 GeV
seen	96k	¹ DEL-AMO-SA..11G	BABR	$e^+ e^-$ at $\Upsilon(4S)$
seen	12k	¹ MITCHELL	09A CLEO	$e^+ e^-$ at 4.17 GeV
seen	701	² FRABETTI	95B E687	

¹ Amplitude analysis with 6 components.

² Amplitude analysis with 5 components.

$D_s^+ \rightarrow K^+ K_S \pi^0$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	990	¹ ABLIKIM	22AH BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ Amplitude analysis with 5 components.

$D_s^+ \rightarrow 2\pi^+ \pi^-$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	11.1k	¹ ABLIKIM	22BI BES3	Dalitz fit
	10.5k	¹ AUBERT	09O BABR	Dalitz fit
	1.5k	² LINK	04 FOCS	Dalitz fit
	848	³ AITALA	01A E791	Dalitz fit

¹ Amplitude analysis with 4 components, one of which is a model-independent $\pi^+ \pi^-$ S-wave parametrisation as complex numbers in 29 $\pi^+ \pi^-$ mass bins.

² Amplitude analysis with 5 components.

³ Amplitude analysis with 6 components.

$D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $\pi^+ \pi^+ \pi^- \eta$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	2.1k	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ Amplitude analysis with 11 components.

$D_s^+ \rightarrow \pi^+ \pi^0 \eta'$ partial wave analyses.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	395	¹ ABLIKIM	22AA BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ The only significant contribution found in this analysis is $D_s^+ \rightarrow \rho^+ \eta'$.

$D_s^+ \rightarrow \pi^+ 2\pi^0$ partial wave analyses.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	440	ABLIKIM	22Z	BES3 $e^+ e^-$ at 4.178–4.226 GeV

$D_s^+ \rightarrow K^+ \pi^+ \pi^-$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	1.3k	¹ ABLIKIM	22AC	BES3 $e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 22AC uses an amplitude analysis with 8 components .

$D_s^+ \rightarrow K^+ \pi^+ \pi^- \pi^0$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	550	¹ ABLIKIM	22BL	BES3 $e^+ e^-$ at 4.178–4.226 GeV

¹ Amplitude analysis with 11 components.

$D_s^+ \rightarrow 2K_S^0 \pi^+$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	400	ABLIKIM	22F	BES3 $e^+ e^-$ at 4.178–4.226 GeV

$D_s^+ \rightarrow (KS)^0 K^- 2\pi^+$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $K_S^0 K^- 2\pi^+$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1.3k	¹ ABLIKIM	21K	BES3 $e^+ e^-$ at 4.178–4.226 GeV

¹ Amplitude analysis with 13 components.

$D_s^+ \rightarrow K^- K^+ \pi^+ \pi^0$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $K^- K^+ \pi^+ \pi^0$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	3k	¹ ABLIKIM	21U	BES3 $e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis with 9 components.

$D_s^+ \rightarrow K^- K^+ 2\pi^+ \pi^-$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	309	ABLIKIM	22AB	BES3 $e^+ e^-$ at 4.178–4.226 GeV

$D_s^+ - D_s^-$ CP-VIOLATING DECAY-RATE ASYMMETRIES

This is the difference between D_s^+ and D_s^- partial widths for the decay to state f , divided by the sum of the widths:

$$A_{CP}(f) = [\Gamma(D_s^+ \rightarrow f) - \Gamma(D_s^- \rightarrow \bar{f})] / [\Gamma(D_s^+ \rightarrow f) + \Gamma(D_s^- \rightarrow \bar{f})].$$

$A_{CP}(\mu^\pm \nu)$ in $D_s^+ \rightarrow \mu^+ \nu$, $D_s^- \rightarrow \mu^- \bar{\nu}_\mu$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.2 ± 2.5 OUR AVERAGE				
$-1.2 \pm 2.5 \pm 1.0$	2.2k	ABLIKIM	21BE	BES3 $e^+ e^-$ at 4.178, 4.226 GeV
4.8 ± 6.1		ALEXANDER	09	CLEO $e^+ e^-$ at 4170 MeV

$A_{CP}(\tau^\pm \nu)$ in $D_s^+ \rightarrow \tau^+ \nu_\tau$, $D_s^- \rightarrow \tau^- \bar{\nu}_\tau$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.9±4.8±1.0	950	¹ ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV

¹ ABLIKIM 21BE also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the result is $(-0.1 \pm 1.9 \pm 1.0)\%$.

$A_{CP}(K^\pm K_S^0)$ in $D_s^\pm \rightarrow K^\pm K_S^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.09±0.26 OUR AVERAGE				
0.6 ±2.8 ±0.6	1.8k	ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV
-0.05±0.23±0.24	288k	¹ LEES	13E BABR	$e^+ e^-$ at $\Upsilon(4S)$
2.6 ±1.5 ±0.6		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
0.12±0.36±0.22		KO	10 BELL	$e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.7 ±1.8 ±0.9	4.0k	MENDEZ	10 CLEO	See ONYISI 13
4.9 ±2.1 ±0.9		ALEXANDER	08 CLEO	See MENDEZ 10

¹ LEES 13E finds that after subtracting the contribution due to $K^0 - \bar{K}^0$ mixing, the CP asymmetry is $(+0.28 \pm 0.23 \pm 0.24)\%$.

$A_{CP}(K^\pm K_L^0)$ in $D_s^\pm \rightarrow K^\pm K_L^0$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-1.1±2.6±0.6	2.3k	ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV

$A_{CP}(K^+ K^- \pi^\pm)$ in $D_s^\pm \rightarrow K^+ K^- \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-0.5±0.8±0.4	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3±1.1±0.8	ALEXANDER 08	CLEO	See ONYISI 13
-------------	--------------	------	---------------

$A_{CP}(\phi \pi^\pm)$ in $D_s^\pm \rightarrow \phi \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-0.38±0.26±0.08	ABAZOV 14B	D0	$p\bar{p}$ at 1.96 TeV

$A_{CP}(K^\pm K_S^0 \pi^0)$ in $D_s^\pm \rightarrow K^\pm K_S^0 \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-1.6±6.0±1.1	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(2K_S^0 \pi^\pm)$ in $D_s^\pm \rightarrow 2K_S^0 \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
3.1±5.2±0.6	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(K^+ K^- \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow K^+ K^- \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.0±2.7±1.2	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

-5.9±4.2±1.2	ALEXANDER 08	CLEO	See ONYISI 13
--------------	--------------	------	---------------

$A_{CP}(K^\pm K_S^0 \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K^\pm K_S^0 \pi^+ \pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-5.7 \pm 5.3 \pm 0.9$	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(K_S^0 K^\mp 2\pi^\pm)$ in $D_s^+ \rightarrow K_S^0 K^\mp 2\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$4.1 \pm 2.7 \pm 0.9$	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.7 \pm 3.6 \pm 1.1$	ALEXANDER 08	CLEO	See ONYISI 13
------------------------	--------------	------	---------------

$A_{CP}(\pi^+ \pi^- \pi^\pm)$ in $D_s^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-0.7 \pm 3.0 \pm 0.6$	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.0 \pm 4.6 \pm 0.7$	ALEXANDER 08	CLEO	See ONYISI 13
-----------------------	--------------	------	---------------

$A_{CP}(\pi^\pm \eta)$ in $D_s^\pm \rightarrow \pi^\pm \eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.3 ± 0.4 OUR AVERAGE				

$0.8 \pm 0.7 \pm 0.5$	38k	AAIJ 21U	LHCB	pp at 13 TeV
-----------------------	-----	----------	------	----------------

$0.2 \pm 0.3 \pm 0.3$	22k	GUAN 21	BELL	$e^+ e^- \approx \Upsilon(4, 5S)$
-----------------------	-----	---------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.1 \pm 3.0 \pm 0.8$		ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV
-----------------------	--	-----------	------	-----------------------

$-4.6 \pm 2.9 \pm 0.3$	2.5k	MENDEZ 10	CLEO	See ONYISI 13
------------------------	------	-----------	------	---------------

$-8.2 \pm 5.2 \pm 0.8$		ALEXANDER 08	CLEO	See MENDEZ 10
------------------------	--	--------------	------	---------------

$A_{CP}(\pi^\pm \eta')$ in $D_s^\pm \rightarrow \pi^\pm \eta'$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.9 ± 0.5 OUR AVERAGE				

$-0.82 \pm 0.36 \pm 0.35$	152k	AAIJ 17AF	LHCB	pp at 7, 8 TeV
---------------------------	------	-----------	------	------------------

$-2.2 \pm 2.2 \pm 0.6$		ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV
------------------------	--	-----------	------	-----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-6.1 \pm 3.0 \pm 0.3$	1.4k	MENDEZ 10	CLEO	See ONYISI 13
------------------------	------	-----------	------	---------------

$-5.5 \pm 3.7 \pm 1.2$		ALEXANDER 08	CLEO	See MENDEZ 10
------------------------	--	--------------	------	---------------

$A_{CP}(\eta \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow \eta \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-0.5 \pm 3.9 \pm 2.0$	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(\eta' \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow \eta' \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$-0.4 \pm 7.4 \pm 1.9$	ONYISI 13	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(K^\pm \pi^0)$ in $D_s^\pm \rightarrow K^\pm \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2 ± 4 OUR AVERAGE		Error includes scale factor of 1.2.		

$-0.8 \pm 3.9 \pm 1.2$	2.8k	AAIJ 21U	LHCB	pp at 7, 8, 13 TeV
------------------------	------	----------	------	----------------------

$6.4 \pm 4.4 \pm 1.1$	12k	GUAN 21	BELL	$e^+ e^- \approx \Upsilon(4, 5S)$
-----------------------	-----	---------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-26.6 \pm 23.8 \pm 0.9$	202	MENDEZ	10	CLEO	e^+e^- at 4170 MeV
2 ± 29		ADAMS	07A	CLEO	See MENDEZ 10

$A_{CP}(\bar{K}^0/K^0\pi^\pm)$ in $D_s^+ \rightarrow \bar{K}^0\pi^+$, $D_s^- \rightarrow K^0\pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.4 ± 0.5 OUR AVERAGE

$0.38 \pm 0.46 \pm 0.17$	121k	¹ AAIJ	14BD	LHCB	pp at 7, 8 TeV
$0.3 \pm 2.0 \pm 0.3$	14k	LEES	13E	BABR	e^+e^- at $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.61 \pm 0.83 \pm 0.14$	26k	AAIJ	13W	LHCB	See AAIJ 14BD
--------------------------	-----	------	-----	------	---------------

¹AAIJ 14BD reports its result as $A_{CP}(D_s^\pm \rightarrow K_S^0 K^\pm)$ with CP -violation effects in the $K^0 - \bar{K}^0$ system subtracted. It also measures $A_{CP}(D^\pm \rightarrow \bar{K}^0/K^0 K^\pm) + A_{CP}(D_s^\pm \rightarrow \bar{K}^0/K^0 \pi^\pm) = (0.41 \pm 0.49 \pm 0.26)\%$.

$A_{CP}(K_S^0\pi^\pm)$ in $D_s^\pm \rightarrow K_S^0\pi^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.20 ± 0.18 OUR AVERAGE

$0.16 \pm 0.17 \pm 0.05$	721k	AAIJ	19T	LHCB	pp at 7, 8, 13 TeV
$0.6 \pm 2.0 \pm 0.3$	14k	LEES	13E	BABR	e^+e^- at $\Upsilon(4S)$
$5.45 \pm 2.50 \pm 0.33$		KO	10	BELL	$e^+e^- \approx \Upsilon(4S)$
$16.3 \pm 7.3 \pm 0.3$	0.4k	MENDEZ	10	CLEO	e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

27 ± 11		ADAMS	07A	CLEO	See MENDEZ 10
-------------	--	-------	-----	------	---------------

$A_{CP}(K^\pm\pi^+\pi^-)$ in $D_s^\pm \rightarrow K^\pm\pi^+\pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

3.7 ± 2.7 OUR AVERAGE

$3.3 \pm 3.0 \pm 1.3$	1.3k	ABLIKIM	22AC	BES3	e^+e^- at 4.178–4.226 GeV
$4.5 \pm 4.8 \pm 0.6$		ONYISI	13	CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$11.2 \pm 7.0 \pm 0.9$		ALEXANDER	08	CLEO	See ONYISI 13
------------------------	--	-----------	----	------	---------------

$A_{CP}(K_S^0\pi^+\pi^0)$ in $D_s^\pm \rightarrow K_S^0\pi^+\pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.7 ± 5.5 ± 0.9 ¹ABLIKIM 21AB BES3 e^+e^- at 4.178–4.226 GeV

¹ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$A_{CP}(K^\pm\pi^+\pi^-\pi^0)$ in $D_s^\pm \rightarrow K^\pm\pi^+\pi^-\pi^0$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.6 ± 5.4 ± 0.7 776 ABLIKIM 22BL BES3 e^+e^- at 4.178–4.226 GeV

$A_{CP}(K^\pm\eta)$ in $D_s^\pm \rightarrow K^\pm\eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.8 ± 1.9 OUR AVERAGE

$0.9 \pm 3.7 \pm 1.1$	2.5k	AAIJ	21U	LHCB	pp at 13 TeV
$2.1 \pm 2.1 \pm 0.4$	14k	GUAN	21	BELL	$e^+e^- \approx \Upsilon(4,5S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.3 \pm 15.2 \pm 0.9$	222	MENDEZ	10	CLEO	e^+e^- at 4170 MeV
-20 ± 18		ADAMS	07A	CLEO	See MENDEZ 10

$A_{CP}(K^\pm \eta'(958))$ in $D_s^\pm \rightarrow K^\pm \eta'(958)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$6.0 \pm 18.9 \pm 0.9$	56 ± 17	MENDEZ	10	CLEO e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

-17 ± 37		ADAMS	07A	CLEO	See MENDEZ 10
--------------	--	-------	-----	------	---------------

CP VIOLATING ASYMMETRIES OF P-ODD (T-ODD) MOMENTS

$A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$ is a parity-odd correlation of the K^+ , π^+ , and π^- momenta for the D_s^+ . $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$ is the corresponding quantity for the D_s^- . Then

$$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)], \text{ and}$$

$$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)], \text{ and}$$

$A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$. C_T and \bar{C}_T are commonly referred to as T -odd moments, because they are odd under T reversal. However, the T -conjugate process $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D_s^\pm$ is not accessible, while the P -conjugate process is.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$-13.6 \pm 7.7 \pm 3.4$	$29.8 \pm 0.3k$	LEES	11E BABR	$e^+e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-36 \pm 67 \pm 23$	508 ± 34	LINK	05E FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
---------------------	--------------	------	----------	--

D_s^+ Semileptonic Form Factors and Decay Constants

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ± 0.11	OUR AVERAGE	Error includes scale factor of 2.4.		
$0.816 \pm 0.036 \pm 0.030$	$25 \pm 0.5k$	¹ AUBERT	08AN BABR	$\phi e^+ \nu_e$
$0.713 \pm 0.202 \pm 0.284$	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
$1.57 \pm 0.25 \pm 0.19$	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
$1.4 \pm 0.5 \pm 0.3$	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
$1.1 \pm 0.8 \pm 0.1$	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
$2.1 \begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix} \pm 0.2$	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

¹ To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at $m_A = 2.5 \text{ GeV}/c^2$ and $m_V = 2.1 \text{ GeV}/c^2$. A simultaneous fit to r_2 , r_V , r_0 (a significant s -wave contribution) and m_A , gives $r_2 = 0.763 \pm 0.071 \pm 0.065$.

$r_V \equiv V(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.80 ± 0.08 OUR AVERAGE				
1.807 ± 0.046 ± 0.065	25 ± 0.5k	¹ AUBERT	08AN BABR	$\phi e^+ \nu_e$
1.549 ± 0.250 ± 0.148	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
2.27 ± 0.35 ± 0.22	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
0.9 ± 0.6 ± 0.3	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.8 ± 0.9 ± 0.2	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
2.3 ^{+1.1} / _{-0.9} ± 0.4	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

¹ To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at $m_A = 2.5 \text{ GeV}/c^2$ and $m_V = 2.1 \text{ GeV}/c^2$. A simultaneous fit to r_2 , r_V , r_0 (a significant s -wave contribution) and m_A , gives $r_V = 1.849 \pm 0.060 \pm 0.095$.

Γ_L/Γ_T in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.18 OUR AVERAGE				
1.0 ± 0.3 ± 0.2	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.0 ± 0.5 ± 0.1	90	¹ FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
0.54 ± 0.21 ± 0.10	19	¹ KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

¹ FRABETTI 94F and KODAMA 93 evaluate Γ_L/Γ_T for a lepton mass of zero.

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.4455 ± 0.0053 ± 0.0044	1.8k	ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta' e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.477 ± 0.049 ± 0.011	261	ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV

$f_+(0) |V_{cd}|$ in $D_s^+ \rightarrow K^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.162 ± 0.019 ± 0.003	117	¹ ABLIKIM	19D BES3	$K_S^0 e^+ \nu_e$

¹ Using a two parameter fit in the z expansion.

$r_V \equiv V(0)/A_1(0)$ in $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.34 ± 0.16	155	ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.28 ± 0.07	155	ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV

$f_{D_s^+} |V_{cs}|$ in $D_s^+ \rightarrow \mu^+ \nu_\mu$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
243.1 ± 3.0 ± 3.6 ± 1.0	2.2K	¹ ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$246.2 \pm 3.6 \pm 3.5$ 1.1k ABLIKIM 19E BES3 e^+e^- at 4178 MeV

¹ The third uncertainty is dominated by the uncertainty of the D_s^+ lifetime.

$f_{D_s^+} |V_{cs}|$ in $D_s^+ \rightarrow \tau^+ \nu_\tau$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
245.3 ± 3.0 OUR AVERAGE				
251.6 ± 5.9 ± 4.9	1.7k	¹ ABLIKIM	21AF BES3	e^+e^- at 4.178, 4.226 GeV
244.4 ± 2.3 ± 2.9	4.9k	² ABLIKIM	21AZ BES3	e^+e^- at 4.178, 4.226 GeV
243.0 ± 5.8 ± 4.0 ± 1.0	950	^{3,4} ABLIKIM	21BE BES3	e^+e^- at 4.178, 4.226 GeV

¹ ABLIKIM 21F uses $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}$ decays.

² ABLIKIM 21AZ uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays.

³ ABLIKIM 21BE uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays. When constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau) / \Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the result is $243.2 \pm 2.3 \pm 3.3 \pm 1.0$.

⁴ The third uncertainty is dominated by the uncertainty of the D_s^+ lifetime.

D_s^\pm REFERENCES

ABLIKIM	22AA	JHEP 2204 058	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AB	JHEP 2207 051	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AC	JHEP 2208 196	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AH	PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BH	PR D106 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BI	PR D106 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BL	JHEP 2209 242	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22F	PR D105 L051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22J	PR D105 L031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Z	JHEP 2201 052	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	21T	JHEP 2106 044	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21U	JHEP 2106 019	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	21AB	JHEP 2106 181	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AC	PR D104 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AE	PR D104 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AF	PR D104 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AR	PR D104 L071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AZ	PRL 127 171801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21BE	PR D104 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21F	PR D103 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21K	PR D103 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Y	PR D103 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GUAN	21	PR D103 112005	Y. Guan <i>et al.</i>	(BELLE Collab.)
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	20R	JHEP 2008 146	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	19G	JHEP 1903 176	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19T	PRL 122 191803	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	19AD	PR D99 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AH	PR D99 091101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AM	PR D99 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BD	PR D100 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BE	PRL 123 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19D	PRL 122 061801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19E	PRL 122 071802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19O	PR D99 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19S	PRL 122 121801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18A	PR D97 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	17AF	PL B771 21	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AN	PRL 119 101801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	16O	PR D94 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16T	PR D94 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	15Z	PL B750 466	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HIETALA	15	PR D92 012009	J. Hietala <i>et al.</i>	(MINN, LUTH, OXF)
LEES	15D	PR D91 019901 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	14BD	JHEP 1410 025	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	14B	PRL 112 111804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AAIJ	13AF	PL B724 203	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13V	JHEP 1306 065	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13W	JHEP 1306 112	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	13E	PR D87 052012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ONYISI	13	PR D88 032009	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
ZUPANC	13	JHEP 1309 139	A. Zupanc <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	11G	PR D83 052001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
LEES	11E	PR D84 031103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MARTIN	11	PR D84 012005	L. Martin <i>et al.</i>	(CLEO Collab.)
ASNER	10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10J	PR D82 091103	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
Also		PR D91 019901 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)
KO	10	PRL 104 181602	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MENDEZ	10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
RUBIN	10	PR D82 092007	P. Rubin <i>et al.</i>	(CLEO Collab.)
ALEXANDER	09	PR D79 052001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AUBERT	09O	PR D79 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	09	PR D79 112008	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ECKLUND	09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
GE	09A	PR D80 051102	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KO	09	PRL 102 221802	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
NAIK	09A	PR D80 112004	P. Naik <i>et al.</i>	(CLEO Collab.)
ONYISI	09	PR D79 052002	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
WON	09	PR D80 111101	E. Won <i>et al.</i>	(BELLE Collab.)
YELTON	09	PR D80 052007	J. Yelton <i>et al.</i>	(CLEO Collab.)
ALEXANDER	08	PRL 100 161804	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ATHAR	08	PRL 100 181802	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	08AN	PR D78 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ECKLUND	08	PRL 100 161801	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
KLEMPPT	08	EPJ C55 39	E. Klempt, M. Matveev, A.V. Sarantsev	(BONN+)
LINK	08	PL B660 147	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
WIDHALM	08	PRL 100 241801	L. Widhalm <i>et al.</i>	(BELLE Collab.)
ADAMS	07A	PRL 99 191805	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	07V	PRL 98 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
PEDLAR	07A	PR D76 072002	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
Also		PRL 99 071802	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
AUBERT	05V	PR D71 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05J	PRL 95 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05K	PL B624 166	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04C	PL B586 183	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04D	PL B586 191	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACOSTA	03D	PR D68 072004	D. Acosta <i>et al.</i>	(FNAL CDF-II Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT	02G	PR D65 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
HEISTER	02I	PL B528 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABBIENDI	01L	PL B516 236	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
IORI	01	PL B523 22	M. Iori <i>et al.</i>	(FNAL SELEX Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ALEXANDROV	00	PL B478 31	Y. Alexandrov <i>et al.</i>	(CERN BEATRICE Collab.)
AITALA	99	PL B445 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99D	PL B450 294	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)

CHADHA	98	PR D58 032002	M. Chada <i>et al.</i>	(CLEO Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
BALEST	97	PRL 79 1436	R. Balest <i>et al.</i>	(CLEO Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ARTUSO	96	PL B378 364	M. Artuso <i>et al.</i>	(CLEO Collab.)
BAI	95C	PR D52 3781	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	95	PRL 75 3804	G.W. Brandenburg <i>et al.</i>	(CLEO Collab.)
FRABETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ACOSTA	94	PR D49 5690	D. Acosta <i>et al.</i>	(CLEO Collab.)
AVERY	94B	PL B337 405	P. Avery <i>et al.</i>	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
BUTLER	94	PL B324 255	F. Butler <i>et al.</i>	(CLEO Collab.)
FRABETTI	94F	PL B328 187	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93F	PRL 71 827	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	93	PL B309 483	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	J. Alexander <i>et al.</i>	(CLEO Collab.)
AVERY	92	PRL 68 1279	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	92	PL B281 167	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ALBRECHT	90D	PL B245 315	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander <i>et al.</i>	(CLEO Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
CHEN	89	PL B226 192	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88	PRL 60 897	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
BECKER	87B	PL B184 277	H. Becker <i>et al.</i>	(NA11 and NA32 Collabs.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
USHIDA	86	PRL 56 1767	N. Ushida <i>et al.</i>	(FNAL E531 Collab.)
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DERRICK	85B	PRL 54 2568	M. Derrick <i>et al.</i>	(HRS Collab.)
AIHARA	84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALTHOFF	84	PL 136B 130	M. Althoff <i>et al.</i>	(TASSO Collab.)
BAILEY	84	PL 139B 320	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
CHEN	83C	PRL 51 634	A. Chen <i>et al.</i>	(CLEO Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
---------	----	------------	----------------------------	--------------