

$B^\pm/B^0$ ADMIXTURE

B DECAY MODES

The branching fraction measurements are for an admixture of B mesons at the $\Upsilon(4S)$. The values quoted assume that $B(\Upsilon(4S) \rightarrow B\bar{B}) = 100\%$.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the treatment of multiple D 's in the final state must be defined. One possibility would be to count the number of events with one-or-more D 's and divide by the total number of B 's. Another possibility would be to count the total number of D 's and divide by the total number of B 's, which is the definition of average multiplicity. The two definitions are identical if only one D is allowed in the final state. Even though the "one-or-more" definition seems sensible, for practical reasons inclusive branching fractions are almost always measured using the multiplicity definition. For heavy final state particles, authors call their results inclusive branching fractions while for light particles some authors call their results multiplicities. In the B sections, we list all results as inclusive branching fractions, adopting a multiplicity definition. This means that inclusive branching fractions can exceed 100% and that inclusive partial widths can exceed total widths, just as inclusive cross sections can exceed total cross section.

$\bar{B}$ modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
Γ_1 $e^+ \nu_e$ anything	[a]	
Γ_2 $\mu^+ \nu_\mu$ anything	[a]	
Γ_3 $\ell^+ \nu_\ell$ anything	[a,b] (10.82 $\pm$ 0.15) %	
Γ_4 $D^- \ell^+ \nu_\ell$ anything	[b] (2.6 $\pm$ 0.5) %	
Γ_5 $\bar{D}^0 \ell^+ \nu_\ell$ anything	[b] (7.2 $\pm$ 1.5) %	
Γ_6 $\bar{D} \ell^+ \nu_\ell$	(2.41 $\pm$ 0.12) %	
Γ_7 $D^{*-} \ell^+ \nu_\ell$ anything	[c] (6.7 $\pm$ 1.3) $\times 10^{-3}$	
Γ_8 $D^{*0} \ell^+ \nu_\ell$ anything		
Γ_9 $\bar{D}^* \ell^+ \nu_\ell$	[d] (4.95 $\pm$ 0.11) %	
Γ_{10} $\bar{D}^* e^+ \nu_e$		
Γ_{11} $\bar{D}^* \mu^+ \nu_\mu$		
Γ_{12} $\bar{D}^{**} \ell^+ \nu_\ell$	[b,e] (2.7 $\pm$ 0.7) %	
Γ_{13} $\bar{D}_1(2420) \ell^+ \nu_\ell$ anything	(3.8 $\pm$ 1.3) $\times 10^{-3}$	S=2.4
Γ_{14} $\bar{D} \pi \ell^+ \nu_\ell$ anything + $\bar{D}^* \pi \ell^+ \nu_\ell$ anything	(2.6 $\pm$ 0.5) %	S=1.5
Γ_{15} $\bar{D} \pi \ell^+ \nu_\ell$ anything	(1.5 $\pm$ 0.6) %	
Γ_{16} $\bar{D}^* \pi \ell^+ \nu_\ell$ anything	(1.9 $\pm$ 0.4) %	
Γ_{17} $\bar{D}_2(2460) \ell^+ \nu_\ell$ anything	(4.4 $\pm$ 1.6) $\times 10^{-3}$	

Γ_{18}	$D^{*-}\pi^+\ell^+\nu_\ell$ anything	(	1.00 ± 0.34	) %	
Γ_{19}	$\bar{D}\pi^+\pi^-\ell^+\nu_\ell$	(	1.62 ± 0.32	) $\times 10^{-3}$	
Γ_{20}	$\bar{D}^*\pi^+\pi^-\ell^+\nu_\ell$	(	9.4 ± 3.2	) $\times 10^{-4}$	
Γ_{21}	$D_s^-\ell^+\nu_\ell$ anything	[b] <	7	$\times 10^{-3}$	CL=90%
Γ_{22}	$D_s^-\ell^+\nu_\ell K^+$ anything	[b] <	5	$\times 10^{-3}$	CL=90%
Γ_{23}	$D_s^-\ell^+\nu_\ell K^0$ anything	[b] <	7	$\times 10^{-3}$	CL=90%
Γ_{24}	$X_c\ell^+\nu_\ell$	(	10.63 ± 0.15	) %	
Γ_{25}	$X_u\ell^+\nu_\ell$	(	1.92 ± 0.21	) $\times 10^{-3}$	
Γ_{26}	$X_ue^+\nu_e$	(	1.57 ± 0.19	) $\times 10^{-3}$	
Γ_{27}	$X_u\mu^+\nu_\mu$	(	1.62 ± 0.21	) $\times 10^{-3}$	
Γ_{28}	$K^+\ell^+\nu_\ell$ anything	[b] (	6.3 ± 0.5	) %	
Γ_{29}	$K^-\ell^+\nu_\ell$ anything	[b] (	10 ± 4	) $\times 10^{-3}$	
Γ_{30}	$K^0/\bar{K}^0\ell^+\nu_\ell$ anything	[b] (	4.6 ± 0.5	) %	
Γ_{31}	$\bar{D}\tau^+\nu_\tau$	(	8.3 ± 0.8	) $\times 10^{-3}$	
Γ_{32}	$\bar{D}^*\tau^+\nu_\tau$	(	1.42 ± 0.07	) %	

D, D*, or D_s modes

Γ_{33}	$D^\pm$ anything	(	23.1 ± 1.2	) %	
Γ_{34}	$D^0/\bar{D}^0$ anything	(	64.6 ± 2.1	) %	S=1.5
Γ_{35}	$D^*(2010)^\pm$ anything	(	22.5 ± 1.5	) %	
Γ_{36}	$\bar{D}^*(2007)^0$ anything	(	26.0 ± 2.7	) %	
Γ_{37}	$D_s^\pm$ anything	[f] (	10.6 ± 0.6	) %	S=1.7
Γ_{38}	$D_s^{*\pm}$ anything	(	6.3 ± 1.0	) %	
Γ_{39}	$D_s^{*\pm}\bar{D}^*$	(	3.4 ± 0.6	) %	
Γ_{40}	$\bar{D}D_{s0}(2317)$	seen			
Γ_{41}	$\bar{D}D_{sJ}(2457)$	seen			
Γ_{42}	$D^*(*)\bar{D}^*(*)K^0 + D^*(*)\bar{D}^*(*)K^\pm$ [f,g]	(	7.1 ± 2.7 -1.7	) %	
Γ_{43}	$b \rightarrow c\bar{c}s$	(	22 ± 4	) %	
Γ_{44}	$D_s^*(*)\bar{D}^*(*)$	[f,g] (	5.0 ± 0.4	) %	
Γ_{45}	$D^*D^*(2010)^\pm$	[f] <	5.9	$\times 10^{-3}$	CL=90%
Γ_{46}	$DD^*(2010)^\pm + D^*D^\pm$	[f] <	5.5	$\times 10^{-3}$	CL=90%
Γ_{47}	$DD^\pm$	[f] <	3.1	$\times 10^{-3}$	CL=90%
Γ_{48}	$D_s^*(*)^\pm\bar{D}^*(*)X(n\pi^\pm)$	[f,g] (	9 ± 5 -4	) %	
Γ_{49}	$\bar{D}^*(2010)\gamma$	<	1.1	$\times 10^{-3}$	CL=90%
Γ_{50}	$D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-,$ $D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0,$ $D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0,$ $D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega$	[f] <	4	$\times 10^{-4}$	CL=90%
Γ_{51}	$D_{s1}(2536)^+$ anything	<	9.5	$\times 10^{-3}$	CL=90%

Charmonium modes

Γ_{52}	$J/\psi(1S)$ anything	(	1.094 ± 0.032	) %	S=1.1
Γ_{53}	$J/\psi(1S)$ (direct) anything	(	7.8 ± 0.4	) $\times 10^{-3}$	S=1.1

Γ_{54}	$\psi(2S)$ anything	(	3.07 ± 0.21	) $\times 10^{-3}$	
Γ_{55}	$\chi_{c1}(1P)$ anything	(	3.55 ± 0.27	) $\times 10^{-3}$	S=1.3
Γ_{56}	$\chi_{c1}(1P)$ (direct) anything	(	3.08 ± 0.19	) $\times 10^{-3}$	
Γ_{57}	$\chi_{c2}(1P)$ anything	(	9.9 ± 1.7	) $\times 10^{-4}$	S=1.6
Γ_{58}	$\chi_{c2}(1P)$ (direct) anything	(	7.5 ± 1.1	) $\times 10^{-4}$	
Γ_{59}	$\eta_c(1S)$ anything	<	9	$\times 10^{-3}$	CL=90%
Γ_{60}	$K\chi_{c1}(3872)$	(	1.9 ± 0.7	) $\times 10^{-4}$	
Γ_{61}	$KX(3940)$, $X \rightarrow D^{*0}D^0$	<	6.7	$\times 10^{-5}$	CL=90%
Γ_{62}	$K\chi_{c0}(3915)$, $\chi_{c0} \rightarrow \omega J/\psi$ [h]	(	7.1 ± 3.4	) $\times 10^{-5}$	

K or K* modes

Γ_{63}	$K^\pm$ anything	[f]	(	78.9 ± 2.5	) %	
Γ_{64}	K^+ anything	(	66	± 5	) %	
Γ_{65}	K^- anything	(	13	± 4	) %	
Γ_{66}	$K^0/\bar{K}^0$ anything	[f]	(	64	± 4	) %
Γ_{67}	$K^*(892)^\pm$ anything	(	18	± 6	) %	
Γ_{68}	$K^*(892)^0/\bar{K}^*(892)^0$ anything	[f]	(	14.6 ± 2.6	) %	
Γ_{69}	$K^*(892)\gamma$	(	4.2 ± 0.6	) $\times 10^{-5}$		
Γ_{70}	$\eta K\gamma$	(	8.5 ± 1.8	-1.6	) $\times 10^{-6}$	
Γ_{71}	$K_1(1400)\gamma$	<	1.27	$\times 10^{-4}$	CL=90%	
Γ_{72}	$K_2^*(1430)\gamma$	(	1.7 ± 0.6	-0.5	) $\times 10^{-5}$	
Γ_{73}	$K_2(1770)\gamma$	<	1.2	$\times 10^{-3}$	CL=90%	
Γ_{74}	$K_3^*(1780)\gamma$	<	3.7	$\times 10^{-5}$	CL=90%	
Γ_{75}	$K_4^*(2045)\gamma$	<	1.0	$\times 10^{-3}$	CL=90%	
Γ_{76}	$K\eta'(958)$	(	8.3 ± 1.1	) $\times 10^{-5}$		
Γ_{77}	$K^*(892)\eta'(958)$	(	4.1 ± 1.1	) $\times 10^{-6}$		
Γ_{78}	$K\eta$	<	5.2	$\times 10^{-6}$	CL=90%	
Γ_{79}	$K^*(892)\eta$	(	1.8 ± 0.5	) $\times 10^{-5}$		
Γ_{80}	$K\phi\phi$	(	2.3 ± 0.9	) $\times 10^{-6}$		
Γ_{81}	$\bar{b} \rightarrow \bar{s}\gamma$	(	3.49 ± 0.19	) $\times 10^{-4}$		
Γ_{82}	$\bar{b} \rightarrow \bar{d}\gamma$	(	9.2 ± 3.0	) $\times 10^{-6}$		
Γ_{83}	$\bar{b} \rightarrow \bar{s}\text{gluon}$	<	6.8	%	CL=90%	
Γ_{84}	η anything	(	2.6 ± 0.5	-0.8	) $\times 10^{-4}$	
Γ_{85}	η' anything	(	4.2 ± 0.9	) $\times 10^{-4}$		
Γ_{86}	K^+ gluon (charmless)	<	1.87	$\times 10^{-4}$	CL=90%	
Γ_{87}	K^0 gluon (charmless)	(	1.9 ± 0.7	) $\times 10^{-4}$		

Light unflavored meson modes

Γ_{88}	$\rho\gamma$	(	1.39 ± 0.25	) $\times 10^{-6}$	S=1.2
Γ_{89}	$\rho/\omega\gamma$	(	1.30 ± 0.23	) $\times 10^{-6}$	S=1.2
Γ_{90}	$\pi^\pm$ anything	[f,i]	(	358 ± 7	) %
Γ_{91}	π^0 anything	(	235 ± 11	) %	
Γ_{92}	η anything	(	17.6 ± 1.6	) %	

Γ_{93}	ρ^0 anything	(21 $\pm$ 5) %	
Γ_{94}	ω anything	< 81 %	CL=90%
Γ_{95}	ϕ anything	(3.43 $\pm$ 0.12) %	
Γ_{96}	$\phi K^*(892)$	< 2.2 $\times 10^{-5}$	CL=90%
Γ_{97}	$\bar{b} \rightarrow \bar{d}$ gluon		
Γ_{98}	π^+ gluon (charmless)	(3.7 $\pm$ 0.8) $\times 10^{-4}$	

Baryon modes

Γ_{99}	$\Lambda_c^+ / \bar{\Lambda}_c^-$ anything	(3.55 $\pm$ 0.35) %	
Γ_{100}	Λ_c^+ anything	< 1.3 %	CL=90%
Γ_{101}	$\bar{\Lambda}_c^-$ anything	< 7 %	CL=90%
Γ_{102}	$\bar{\Lambda}_c^- \ell^+$ anything	< 9 $\times 10^{-4}$	CL=90%
Γ_{103}	$\bar{\Lambda}_c^- e^+$ anything	< 1.8 $\times 10^{-3}$	CL=90%
Γ_{104}	$\bar{\Lambda}_c^- \mu^+$ anything	< – 1.4 $\times 10^{-3}$	CL=90%
Γ_{105}	$\bar{\Lambda}_c^- p$ anything	(2.02 $\pm$ 0.32) %	
Γ_{106}	$\bar{\Lambda}_c^- p e^+ \nu_e$	< 8 $\times 10^{-4}$	CL=90%
Γ_{107}	$\bar{\Sigma}_c^{--}$ anything	(3.3 $\pm$ 1.7) $\times 10^{-3}$	
Γ_{108}	$\bar{\Sigma}_c^{--}$ anything	< 8 $\times 10^{-3}$	CL=90%
Γ_{109}	$\bar{\Sigma}_c^0$ anything	(3.6 $\pm$ 1.7) $\times 10^{-3}$	
Γ_{110}	$\bar{\Sigma}_c^0 N (N = p \text{ or } n)$	< 1.2 $\times 10^{-3}$	CL=90%
Γ_{111}	Ξ_c^0 anything, $\Xi_c^0 \rightarrow \Xi^- \pi^+$	(1.93 $\pm$ 0.30) $\times 10^{-4}$	S=1.1
Γ_{112}	$\Xi_c^+, \Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$	(4.5 $\pm$ 1.3 / 1.2) $\times 10^{-4}$	
Γ_{113}	$p/\bar{p}$ anything	[f] (8.0 $\pm$ 0.4) %	
Γ_{114}	$p/\bar{p}$ (direct) anything	[f] (5.5 $\pm$ 0.5) %	
Γ_{115}	$\bar{p} e^+ \nu_e$ anything	< 5.9 $\times 10^{-4}$	CL=90%
Γ_{116}	$\Lambda/\bar{\Lambda}$ anything	[f] (4.0 $\pm$ 0.5) %	
Γ_{117}	Λ anything	seen	
Γ_{118}	$\bar{\Lambda}$ anything	seen	
Γ_{119}	$\Xi^- / \bar{\Xi}^+$ anything	[f] (2.7 $\pm$ 0.6) $\times 10^{-3}$	
Γ_{120}	baryons anything	(6.8 $\pm$ 0.6) %	
Γ_{121}	$p\bar{p}$ anything	(2.47 $\pm$ 0.23) %	
Γ_{122}	$\Lambda\bar{p}/\bar{\Lambda}p$ anything	[f] (2.5 $\pm$ 0.4) %	
Γ_{123}	$\Lambda\bar{\Lambda}$ anything	< 5 $\times 10^{-3}$	CL=90%

**Lepton Family number (LF) violating modes or
 $\Delta B = 1$ weak neutral current (B1) modes**

Γ_{124}	$s e^+ e^-$	B1	(6.7 $\pm$ 1.7) $\times 10^{-6}$	S=2.0
Γ_{125}	$s \mu^+ \mu^-$	B1	(4.3 $\pm$ 1.0) $\times 10^{-6}$	
Γ_{126}	$s \ell^+ \ell^-$	B1 [b]	(5.8 $\pm$ 1.3) $\times 10^{-6}$	S=1.8
Γ_{127}	$\pi \ell^+ \ell^-$	B1	< 5.9 $\times 10^{-8}$	CL=90%
Γ_{128}	$\pi e^+ e^-$	B1	< 1.10 $\times 10^{-7}$	CL=90%
Γ_{129}	$\pi \mu^+ \mu^-$	B1	< 5.0 $\times 10^{-8}$	CL=90%
Γ_{130}	$K e^+ e^-$	B1	(4.4 $\pm$ 0.6) $\times 10^{-7}$	

Γ_{131}	$K^*(892) e^+ e^-$	$B1$	(	1.19 ± 0.20	) $\times 10^{-6}$	$S=1.2$
Γ_{132}	$K \mu^+ \mu^-$	$B1$	(	4.4 ± 0.4	) $\times 10^{-7}$	
Γ_{133}	$K^*(892) \mu^+ \mu^-$	$B1$	(	1.06 ± 0.09	) $\times 10^{-6}$	
Γ_{134}	$K \ell^+ \ell^-$	$B1$	(	4.8 ± 0.4	) $\times 10^{-7}$	
Γ_{135}	$K^*(892) \ell^+ \ell^-$	$B1$	(	1.05 ± 0.10	) $\times 10^{-6}$	
Γ_{136}	$K \nu \bar{\nu}$	$B1$	<	1.6	$\times 10^{-5}$	CL=90%
Γ_{137}	$K^* \nu \bar{\nu}$	$B1$	<	2.7	$\times 10^{-5}$	CL=90%
Γ_{138}	$\pi \nu \bar{\nu}$	$B1$	<	8	$\times 10^{-6}$	CL=90%
Γ_{139}	$\rho \nu \bar{\nu}$	$B1$	<	2.8	$\times 10^{-5}$	CL=90%
Γ_{140}	$s e^\pm \mu^\mp$	LF	[f] <	2.2	$\times 10^{-5}$	CL=90%
Γ_{141}	$\pi e^\pm \mu^\mp$	LF	<	9.2	$\times 10^{-8}$	CL=90%
Γ_{142}	$\rho e^\pm \mu^\mp$	LF	<	3.2	$\times 10^{-6}$	CL=90%
Γ_{143}	$K e^\pm \mu^\mp$	LF	<	3.8	$\times 10^{-8}$	CL=90%
Γ_{144}	$K^*(892) e^\pm \mu^\mp$	LF	<	5.1	$\times 10^{-7}$	CL=90%

[a] These values are model dependent.

[b] An ℓ indicates an e or a μ mode, not a sum over these modes.

[c] Here “anything” means at least one particle observed.

[d] This is a $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell)$ value.

[e] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.

[f] The value is for the sum of the charge states or particle/antiparticle states indicated.

[g] $D^{(*)} \bar{D}^{(*)}$ stands for the sum of $D^* \bar{D}^*$, $D^* \bar{D}$, $D \bar{D}^*$, and $D \bar{D}$.

[h] $X(3915)$ denotes a near-threshold enhancement in the $\omega J/\psi$ mass spectrum.

[i] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

$B^\pm/B^0$ ADMIXTURE BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{ anything})/\Gamma_{\text{total}}$

Γ_3/Γ

These branching fraction values are model dependent.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
10.82±0.15 OUR EVALUATION	(Produced by HFLAV)		
10.49±0.20 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
10.34±0.04±0.26	¹ LEES	17B	BABR $e^+ e^- \rightarrow \gamma(4S)$
10.28±0.18±0.24	² URQUIJO	07	BELL $e^+ e^- \rightarrow \gamma(4S)$
10.91±0.09±0.24	³ MAHMOOD	04	CLEO $e^+ e^- \rightarrow \gamma(4S)$
9.7 ±0.5 ±0.4	⁴ ALBRECHT	93H	ARG $e^+ e^- \rightarrow \gamma(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9.96±0.19±0.32	⁵ AUBERT,B	06Y	BABR Repl. by LEES 17B
10.85±0.21±0.36	⁶ OKABE	05	BELL Repl. by URQUIJO 07
10.83±0.16±0.06	⁷ AUBERT	04X	BABR Repl. by AUBERT,B 06Y

$10.36 \pm 0.06 \pm 0.23$	⁸ AUBERT,B	04A	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$10.87 \pm 0.18 \pm 0.30$	⁹ AUBERT	03	BABR	Repl. by AUBERT 04X
$10.90 \pm 0.12 \pm 0.49$	¹⁰ ABE	02Y	BELL	Repl. by OKABE 05
$10.49 \pm 0.17 \pm 0.43$	¹¹ BARISH	96B	CLE2	Repl. by MAHMOOD 04
$10.80 \pm 0.20 \pm 0.56$	¹² HENDERSON	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$10.0 \pm 0.4 \pm 0.3$	¹³ YANAGISAWA	91	CSB2	$e^+e^- \rightarrow \Upsilon(4S)$
$10.3 \pm 0.6 \pm 0.2$	¹⁴ ALBRECHT	90H	ARG	Direct e at $\Upsilon(4S)$
$10.0 \pm 0.6 \pm 0.2$	¹⁵ ALBRECHT	90H	ARG	Direct μ at $\Upsilon(4S)$
$11.7 \pm 0.4 \pm 1.0$	¹⁶ WACHS	89	CBAL	Direct e at $\Upsilon(4S)$
$12.0 \pm 0.7 \pm 0.5$	CHEN	84	CLEO	Direct e at $\Upsilon(4S)$
$10.8 \pm 0.6 \pm 1.0$	CHEN	84	CLEO	Direct μ at $\Upsilon(4S)$
$11.2 \pm 0.9 \pm 1.0$	LEVMAN	84	CUSB	Direct μ at $\Upsilon(4S)$
$13.2 \pm 0.8 \pm 1.4$	¹⁷ KLOPFEN...	83B	CUSB	Direct e at $\Upsilon(4S)$

¹ LEES 17B measurement is obtained from semileptonic decays to electrons. The result is averaged over $B^\pm$ and B^0 mesons, assuming lepton universality.

² URQUIJO 07 report a measurement of $(10.07 \pm 0.18 \pm 0.21)\%$ for the partial branching fraction of $B \rightarrow e\nu_e X_C$ decay with electron energy above 0.6 GeV. We converted the result to $B \rightarrow e\nu_e X$ branching fraction.

³ Uses charge and angular correlations in $\Upsilon(4S)$ events with a high-momentum lepton and an additional electron.

⁴ ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

⁵ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.074 \pm 0.041 \pm 0.026$.

⁶ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

⁷ The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

⁸ Uses the high-momentum lepton tag method and requires the electron energy above 0.6 GeV.

⁹ Uses the high-momentum lepton tag method. They also report $|V_{cb}| = 0.0423 \pm 0.0007(\text{exp}) \pm 0.0020(\text{theo.})$.

¹⁰ Uses the high-momentum lepton tag method. ABE 02Y also reports $|V_{cb}| = 0.0408 \pm 0.0010(\text{exp}) \pm 0.0025(\text{theo.})$. The second error is due to uncertainties of theoretical inputs.

¹¹ BARISH 96B analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

¹² HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.


¹³ YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\Upsilon(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.

¹⁴ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.

¹⁵ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.097 ± 0.006 is obtained using ISGUR 89B.

¹⁶ Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e\nu\text{up})/\sigma(B \rightarrow e\nu\text{charm}) < 0.065$ at 90% CL.

¹⁷ Ratio $\sigma(b \rightarrow e \nu \text{up})/\sigma(b \rightarrow e \nu \text{charm}) < 0.055$ at CL = 90%.



$\Gamma(e^+ \nu_e \text{ anything})/\Gamma(\mu^+ \nu_\mu \text{ anything})$ Γ_1/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
1.007±0.009±0.019	¹ AGGARWAL 23	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ The accompanying B meson is fully reconstructed in its hadronic decay modes.

$\Gamma(D^- \ell^+ \nu_\ell \text{ anything})/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_4/Γ_3

$\ell = e \text{ or } \mu.$

VALUE	DOCUMENT ID	TECN	COMMENT
0.26±0.07±0.04	¹ FULTON 91	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

¹ FULTON 91 uses $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(\bar{D}^0 \ell^+ \nu_\ell \text{ anything})/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_5/Γ_3

$\ell = e \text{ or } \mu.$

VALUE	DOCUMENT ID	TECN	COMMENT
0.67±0.09±0.10	¹ FULTON 91	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

¹ FULTON 91 uses $B(D^0 \rightarrow K^- \pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(\bar{D} \ell^+ \nu_\ell)/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_6/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.223±0.006±0.009	¹ AUBERT 10	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(D^{*-}\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$0.67 \pm 0.08 \pm 0.10$	ABDALLAH	04D	DLPH $e^+e^- \rightarrow Z^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.6 \pm 0.3 \pm 0.1$	¹ BARISH	95	CLE2 $e^+e^- \rightarrow \gamma(4S)$
¹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.			

 $\Gamma(D^{*0}\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.6 \pm 0.6 \pm 0.1$	¹ BARISH	95	CLE2 $e^+e^- \rightarrow \gamma(4S)$
¹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$, $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.			

 $\Gamma(\bar{D}^*e^+\nu_e)/\Gamma(\bar{D}^*\mu^+\nu_\mu)$ Γ_{10}/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.993 \pm 0.023 \pm 0.023$	¹ PRIM	23	BELL $e^+e^- \rightarrow \gamma(4S)$
¹ This is the lepton-flavor universality ratio $R_{e\mu}$ for the $B^+ \rightarrow \bar{D}^{*0}\ell^+\nu_\ell$ and $B^0 \rightarrow D^{*-}\ell^+\nu_\ell$ average.			

 $\Gamma(\bar{D}^{**}\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_{12}/Γ

D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances. $\ell = e$ or μ , not sum over e and μ modes.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$0.027 \pm 0.005 \pm 0.005$	63	¹	ALBRECHT	93	ARG $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.028	95	²	BARISH	95	CLE2 $e^+e^- \rightarrow \gamma(4S)$
¹ ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHKT model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0\pi^+) = 68.1\%$, $B(D^0 \rightarrow K^-\pi^+) = 3.65\%$, $B(D^0 \rightarrow K^-\pi^+\pi^-\pi^+) = 7.5\%$. We have taken their average e and μ value.					
² BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, assume all nonresonant channels are zero, and use GISW model for relative abundances of D^{**} states.					

 $\Gamma(\bar{D}_1(2420)\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0038 ± 0.0013 OUR AVERAGE	Error includes scale factor of 2.4.		
0.0033 ± 0.0006	¹ ABAZOV	05O	D0 $p\bar{p}$ at 1.96 TeV
0.0074 ± 0.0016	² BUSKULIC	97B	ALEP $e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	³ BUSKULIC	95B	ALEP Repl. by BUSKULIC 97B
¹ Assumes $B(D_1 \rightarrow D^*\pi) = 1$, $B(D_1 \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.397$.			
² BUSKULIC 97B assumes $B(D_1(2420) \rightarrow D^*\pi) = 1$, $B(D_1(2420) \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.378 \pm 0.022$.			
³ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}_1(2420)^0\ell^+\nu_\ell \text{ anything}) \times B(\bar{D}_1(2420)^0 \rightarrow \bar{D}^*(2010)^-\pi^+) = (2.04 \pm 0.58 \pm 0.34)10^{-3}$, where f_B is the production fraction for a single B charge state.			

$$[\Gamma(\bar{D}\pi\ell^+\nu_\ell\text{anything}) + \Gamma(\bar{D}^*\pi\ell^+\nu_\ell\text{anything})]/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ± 0.005 OUR AVERAGE	Error includes scale factor of 1.5.		
0.0340 ± 0.0052 ± 0.0032	¹ ABREU	00R	DLPH $e^+e^- \rightarrow Z$
0.0226 ± 0.0029 ± 0.0033	² BUSKULIC	97B	ALEP $e^+e^- \rightarrow Z$

¹ Assumes no contribution from B_s and b baryons. Further assumes contributions from single pion ($D\pi$ and $D^*\pi$) states only, allowing isospin conservation to relate the relative π^0 and π^+ rates.

² BUSKULIC 97B assumes $B(b \rightarrow B) = 0.378 \pm 0.022$ and uses isospin invariance by assuming that all observed $D^0\pi^+$, $D^{*0}\pi^+$, $D^+\pi^-$, and $D^{*+}\pi^-$ are from D^{**} states. A correction has been applied to account for the production of B_s^0 and Λ_b^0 .

$$\Gamma(\bar{D}\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0154 ± 0.0061	ABREU	00R	DLPH $e^+e^- \rightarrow Z$

$$\Gamma(\bar{D}^*\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0186 ± 0.0038	ABREU	00R	DLPH $e^+e^- \rightarrow Z$

$$\Gamma(\bar{D}_2^*(2460)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0044 ± 0.0016		¹ ABAZOV	05O	D0 $p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0065	95	² BUSKULIC	97B	ALEP $e^+e^- \rightarrow Z$
not seen		³ BUSKULIC	95B	ALEP $e^+e^- \rightarrow Z$

¹ Assumes $B(D_2^* \rightarrow D^*\pi^\pm) = 0.30 \pm 0.06$ and $B(b \rightarrow B) = 0.397$.

² A revised number based on BUSKULIC 97B which assumes $B(D_2^*(2460) \rightarrow D^*\pi^\pm) = 0.20$ and $B(b \rightarrow B) = 0.378 \pm 0.022$.

³ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}_2^*(2460)^0\ell^+\nu_\ell\text{anything}) \times B(\bar{D}_2^*(2460)^0 \rightarrow \bar{D}^*(2010)^-\pi^+) \leq 0.81 \times 10^{-3}$ at CL=95%, where f_B is the production fraction for a single B charge state.

$$\frac{\Gamma(B \rightarrow \bar{D}_2^*(2460)\ell^+\nu_\ell\text{anything}) \times B(D_2^*(2460) \rightarrow D^{*-}\pi^+)}{\Gamma(B \rightarrow \bar{D}_1^*(2420)\ell^+\nu_\ell\text{anything}) \times B(\bar{D}_1^*(2420) \rightarrow D^{*-}\pi^+)}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.39 ± 0.09 ± 0.12	ABAZOV	05O	D0 $p\bar{p}$ at 1.96 TeV

$$\Gamma(D^{*-}\pi^+\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

Includes resonant and nonresonant contributions.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
10.0 ± 2.7 ± 2.1	¹ BUSKULIC	95B	ALEP $e^+e^- \rightarrow Z$

¹ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}^*(2010)^-\pi^+\ell^+\nu_\ell\text{anything}) = (3.7 \pm 1.0 \pm 0.7)10^{-3}$. Above value assumes $f_B = 0.37 \pm 0.03$.

$$\Gamma(\bar{D}\pi^+\pi^-\ell^+\nu_\ell)/\Gamma(\bar{D}\ell^+\nu_\ell) \quad \Gamma_{19}/\Gamma_6$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
6.7 ± 1.0 ± 0.8	¹ LEES	16	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement used electrons and muons as leptons.

$\Gamma(\bar{D}^* \pi^+ \pi^- \ell^+ \nu_\ell) / \Gamma(\bar{D}^* \ell^+ \nu_\ell)$ Γ_{20}/Γ_9

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.5 \pm 0.4$	¹ LEES	16	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measurement used electrons and muons as leptons.

 $\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-3}$	90	¹ ALBRECHT	93E	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports < 0.012 from a measurement of $[\Gamma(B \rightarrow D_s^- \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

 $\Gamma(D_s^- \ell^+ \nu_\ell K^+ \text{ anything}) / \Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5 \times 10^{-3}$	90	¹ ALBRECHT	93E	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports < 0.008 from a measurement of $[\Gamma(B \rightarrow D_s^- \ell^+ \nu_\ell K^+ \text{ anything}) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

 $\Gamma(D_s^- \ell^+ \nu_\ell K^0 \text{ anything}) / \Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-3}$	90	¹ ALBRECHT	93E	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports < 0.012 from a measurement of $[\Gamma(B \rightarrow D_s^- \ell^+ \nu_\ell K^0 \text{ anything}) / \Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

 $\Gamma(X_c \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
10.63 ± 0.15 OUR EVALUATION	(Produced by HFLAV)		
10.29 ± 0.19 OUR AVERAGE			

$10.18 \pm 0.03 \pm 0.24$ ¹ LEES 17B BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$10.44 \pm 0.19 \pm 0.22$ ² URQUIJO 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.64 \pm 0.17 \pm 0.06$ ³ AUBERT 10A BABR Repl. by LEES 17B

$10.61 \pm 0.16 \pm 0.06$ ⁴ AUBERT 04X BABR Repl. by AUBERT 10A

¹ The measurement is obtained from semileptonic decays to electrons $B \rightarrow X_c e \nu$, and using a theoretical model (GAMBINO 07, GAMBINO 11) to predict the contribution from $B \rightarrow X_c \mu e \nu$. The result is averaged over $B^\pm$ and B^0 mesons, assuming lepton universality.

² Measured the independent B^+ and B^0 partial branching fractions with electron energy above 0.4 GeV.

³ Obtained from a combined fit to the moments of observed spectra in inclusive $B \rightarrow X_c \ell^+ \nu_\ell$ decay.

⁴ The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

$\Gamma(X_u \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{25} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.92 ± 0.21 OUR EVALUATION	(Produced by HFLAV)		
1.85 ± 0.08 ± 0.19	¹ CAO	21A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.665 ± 0.087 $^{+0.103}_{-0.094}$	² LEES	17B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
2.01 ± 0.15 ± 0.25	³ LEES	12R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
2.53 ± 0.24 ± 0.24	⁴ AUBERT,B	05X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
2.80 ± 0.52 ± 0.41	⁵ LIMOSANI	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.77 ± 0.29 ± 0.38	⁶ BORNHEIM	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.39 ± 0.14 ± 0.22	⁷ CAO	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.963 ± 0.173 ± 0.159	⁸ URQUIJO	10 BELL	Repl. by CAO 21A
1.18 ± 0.09 ± 0.07	⁹ AUBERT	08AS BABR	Repl. by LEES 12R
2.27 ± 0.26 $^{+0.37}_{-0.33}$	¹⁰ AUBERT	06H BABR	Repl. by LEES 17B
2.24 ± 0.27 ± 0.47	^{11,12} AUBERT	04I BABR	Repl. by AUBERT,B 05X

¹ Measures several partial branching fractions in different phase space regions. The most inclusive result of the full branching fraction is obtained in the region for lepton energy in B rest frame $E_\ell^* > 1$ GeV, where the measured partial branching fraction is $\Delta B = (1.59 \pm 0.07 \pm 0.16) \times 10^{-3}$. The acceptance in that region is reported to be 0.86.

² Obtained from the partial rate $\Delta B = (1.554 \pm 0.082^{+0.095}_{-0.086}) \times 10^{-3}$ for the electron momentum interval of 0.8–2.7 GeV/c based on GGOU1 method ($X_c \ell \nu$, m_c constraint fit of SF parameters).

³ Measures several partial branching fractions in different phase space regions. The most precise result on the full branching fraction is obtained in the region for lepton momentum in B rest frame $p_\ell^* > 1$ GeV/c, where the measured partial branching fraction is $\Delta B = (1.80 \pm 0.13 \pm 0.15) \times 10^{-3}$. The acceptance in that region is reported in a private communication by the Authors to be 0.894. The corresponding $|V_{ub}|$ from the BLNP method is $(4.28 \pm 0.15 \pm 0.18 \pm 0.19) \times 10^{-3}$, where the last uncertainty comes from theoretical prediction.

⁴ Determined from the partial rate $\Delta B = (4.41 \pm 0.42 \pm 0.42) \times 10^{-4}$ measured for electron energy > 2 GeV and hadronic mass squared < 3.5 GeV², and calculated acceptance 0.174 in that region. The V_{ub} is measured as $(4.41 \pm 0.30^{+0.65}_{-0.47} \pm 0.28) \times 10^{-3}$.

⁵ Uses electrons in the momentum interval 1.9–2.6 GeV/c in the center-of-mass frame. The V_{ub} is found to be $(5.08 \pm 0.47^{+0.49}_{-0.48}) \times 10^{-3}$.

⁶ BORNHEIM 02 uses the observed yield of leptons from semileptonic B decays in the end-point momentum interval 2.2–2.6 GeV/c with recent CLEO-2 data on $B \rightarrow X_S \gamma$. The V_{ub} is found to be $(4.08 \pm 0.34 \pm 0.53) \times 10^{-3}$.

⁷ Measurement requires lepton energy $E_\ell^* > 1$ GeV in the B rest frame. It is a part of the inclusive and exclusive $|V_{ub}|$ determination.

⁸ Uses a multivariate analysis method and requires lepton momentum in the B rest frame, $p_\ell^* > 1.0$ GeV/c.

⁹ Measures several partial branching fractions in different phase space regions. The most precise result is obtained in the region for hadronic mass $M_X < 1.55$ GeV/c², and is $\Delta B = (1.18 \pm 0.09 \pm 0.07) \times 10^{-3}$. The corresponding $|V_{ub}|$ from the BLNP method is $(4.27 \pm 0.16 \pm 0.13 \pm 0.30) \times 10^{-3}$, where the last uncertainty comes from the theoretical prediction of the partial rate in the given phase-space region.

- ¹⁰ Obtained from the partial rate $\Delta B = (0.572 \pm 0.041 \pm 0.065) \times 10^{-3}$ for the electron momentum interval of 2.0–2.6 GeV/c based on BLNP method.
- ¹¹ Used BaBar measurement of Semileptonic branching fraction $B(B \rightarrow X \ell \nu_\ell) = (10.87 \pm 0.18 \pm 0.30)\%$ to convert the ratio of rates to branching fraction.
- ¹² The third error includes the systematics and theoretical errors summed in quadrature.

$\Gamma(X_u \ell^+ \nu_\ell) / \Gamma(\ell^+ \nu_\ell \text{ anything})$

Γ_{25}/Γ_3

ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.06 \pm 0.25 \pm 0.42$			¹ AUBERT	04I	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
			² ALBRECHT	94C	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
		107	³ BARTELT	93B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
		77	⁴ ALBRECHT	91C	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
		41	⁵ ALBRECHT	90	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
		76	⁶ FULTON	90	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
<4.0	90		⁷ BEHREND	87	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
<4.0	90		CHEN	84	CLEO Direct e at $\Upsilon(4S)$
<5.5	90		KLOPFEN...	83B	CUSB Direct e at $\Upsilon(4S)$

- ¹ The third error includes the systematics and theoretical errors summed in quadrature.
- ² ALBRECHT 94C find $\Gamma(b \rightarrow c) / \Gamma(b \rightarrow \text{all}) = 0.99 \pm 0.02 \pm 0.04$.
- ³ BARTELT 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval 2.3–2.6 GeV/c which is attributed to $b \rightarrow u \ell \nu_\ell$. This corresponds to a model-dependent partial branching ratio ΔB_{ub} between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.
- ⁴ ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}/V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the 2.3–2.6 GeV momentum range.
- ⁵ ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}/V_{cb}| = 0.10 \pm 0.01$.
- ⁶ FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}/V_{cb}| = 0.1$ using $B(b \rightarrow c \ell \nu) = 10.2 \pm 0.2 \pm 0.7\%$.
- ⁷ The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

$\Gamma(X_u e^+ \nu_e) / \Gamma_{\text{total}}$

Γ_{26}/Γ

Requires $E_e^* > 1$ GeV, where E_e^* is e^+ energy in B rest frame.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.57 \pm 0.10 \pm 0.16$	¹ CAO	21A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ The correlation of 53% with $B(B \rightarrow X_u \mu^+ \nu_\mu)$ (lepton energy in B rest frame $E_{\mu^+}^* > 1$ GeV) is reported.

$\Gamma(X_u \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{27}/Γ Requires $E_\mu^* > 1$ GeV, where E_μ^* is μ^+ energy in B rest frame.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.62 ± 0.10 ± 0.18	¹ CAO	21A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ The correlation of 53% with $B(B \rightarrow X_u e^+ \nu_e)$ (lepton energy in B rest frame $E_{e^+}^* > 1$ GeV) is reported.

 $\Gamma(K^+ \ell^+ \nu_\ell \text{ anything})/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{28}/Γ_3 ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.58 ± 0.05 OUR AVERAGE			
0.594 ± 0.021 ± 0.056	ALBRECHT	94C	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
0.54 ± 0.07 ± 0.06	¹ ALAM	87B	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALAM 87B measurement relies on lepton-kaon correlations.

 $\Gamma(K^- \ell^+ \nu_\ell \text{ anything})/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{29}/Γ_3 ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.092 ± 0.035 OUR AVERAGE			
0.086 ± 0.011 ± 0.044	ALBRECHT	94C	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
0.10 ± 0.05 ± 0.02	¹ ALAM	87B	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALAM 87B measurement relies on lepton-kaon correlations.

 $\Gamma(K^0/\bar{K}^0 \ell^+ \nu_\ell \text{ anything})/\Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{30}/Γ_3 ℓ denotes e or μ , not the sum. Sum over K^0 and $\bar{K}^0$ states.

VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ± 0.05 OUR AVERAGE			
0.452 ± 0.038 ± 0.056	¹ ALBRECHT	94C	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
0.39 ± 0.06 ± 0.04	² ALAM	87B	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 94C assume a $K^0/\bar{K}^0$ multiplicity twice that of K_S^0 .

² ALAM 87B measurement relies on lepton-kaon correlations.

 $\Gamma(\bar{D} \tau^+ \nu_\tau)/\Gamma(\bar{D} \ell^+ \nu_\ell)$ Γ_{31}/Γ_6

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
34.2 ± 2.6 OUR EVALUATION	(Produced by HFLAV)		
35 ± 4 OUR AVERAGE	Error includes scale factor of 1.2.		
30.7 ± 3.7 ± 1.6	¹ CARIA	20	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
37.5 ± 6.4 ± 2.6	^{2,3} HUSCHLE	15	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
44.0 ± 5.8 ± 4.2	^{2,3} LEES	12D	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.16 ± 11.7 ± 5.2 ² AUBERT 08N BABR Repl. by LEES 12D

¹ The tag-side B meson is reconstructed in a semileptonic decay mode and the signal-side τ is reconstructed in a purely leptonic decay. The Belle combination of HUSCHLE 15 and CARIA 20 yields $R(D) = (32.6 \pm 3.4) \times 10^{-2}$.

² Uses a fully reconstructed B meson as a tag on the recoil side.

³ Uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ . Obtained from simultaneous fit to B^+ and B^0 assuming isospin symmetry.

$\Gamma(\bar{D}^* \tau^+ \nu_\tau) / \Gamma(\bar{D}^* \ell^+ \nu_\ell)$ Γ_{32}/Γ_9

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

28.6 $\pm$ 1.2 OUR EVALUATION (Produced by HFLAV) with $\rho^2=1.128 \pm 0.033$ and a correlation 0.747. The fitted χ^2 is 4.8 for 8 degrees of freedom.

29.1 $\pm$ 1.3 OUR AVERAGE

26.2 $^{+4.1+3.5}_{-3.9-3.2}$	¹ ADACHI	240	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
28.1 $\pm$ 1.8 $\pm$ 2.4	² AAIJ	23AR	LHCB	pp at 7 and 8 TeV
28.3 $\pm$ 1.8 $\pm$ 1.4	³ CARIA	20	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
27.0 $\pm$ 3.5 $^{+2.8}_{-2.5}$	⁴ HIROSE	17	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
29.3 $\pm$ 3.8 $\pm$ 1.5	⁵ HUSCHLE	15	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
33.2 $\pm$ 2.4 $\pm$ 1.8	⁵ LEES	12D	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

29.7 $\pm$ 5.6 $\pm$ 1.8	⁶ AUBERT	08N	BABR	Repl. by LEES 12D
--------------------------	---------------------	-----	------	-------------------

¹ Uses leptonic τ decays and a fully reconstructed B meson in hadronic final states as a tag on the recoil side.

² Uses $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and μ^+ as ℓ^+ . The measurement combines fully reconstructed D^{*+} sample with sample where only D^0 from D^* decays is reconstructed.

³ The tag-side B meson is reconstructed in a semileptonic decay mode and the signal-side τ is reconstructed in a purely leptonic decay. The Belle combination of HUSCHLE 15, HIROSE 17, and CARIA 20 yields $R(D^*) = 0.238 \pm 0.018$.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

⁵ Uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ . Obtained from simultaneous fit to B^+ and B^0 assuming isospin symmetry. Uses a fully reconstructed B meson as a tag on the recoil side.

⁶ Uses a fully reconstructed B meson as a tag on the recoil side. The results are normalized to the B^+ decay rate.

 $\langle n_c \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.10 $\pm$ 0.05 ¹ GIBBONS 97B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.98 $\pm$ 0.16 $\pm$ 0.12	² ALAM	87B	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
----------------------------	-------------------	-----	------	------------------------------------

¹ GIBBONS 97B from charm counting using $B(D_s^+ \rightarrow \phi \pi) = 0.036 \pm 0.009$ and $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.044 \pm 0.006$.

² From the difference between K^- and K^+ widths. ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.

 $\Gamma(D^\pm \text{ anything}) / \Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.231 $\pm$ 0.012 OUR AVERAGE

0.230 $\pm$ 0.012 $\pm$ 0.004	¹ GIBBONS	97B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.241 $\pm$ 0.037 $\pm$ 0.004	² BORTOLETTO92		CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.223 $\pm$ 0.051 $\pm$ 0.004	³ ALBRECHT	91H	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.203 $\pm$ 0.048 $\pm$ 0.003	20k	⁴ BORTOLETTO87	CLEO	Sup. by BORTOLETTO 92
-------------------------------	-----	---------------------------	------	-----------------------

¹ GIBBONS 97B reports $[\Gamma(B \rightarrow D^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0216 \pm 0.0008 \pm 0.00082$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BORTOLETTO 92 reports $[\Gamma(B \rightarrow D^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0226 \pm 0.0030 \pm 0.0018$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 91H reports $[\Gamma(B \rightarrow D^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.0209 \pm 0.0027 \pm 0.0040$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BORTOLETTO 87 reports $[\Gamma(B \rightarrow D^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.019 \pm 0.004 \pm 0.002$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.646 ± 0.021 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
0.6663 ± 0.0004 ± 0.0177		ZHUKOVA	23	BELL $e^+e^- \rightarrow \Upsilon(4S)$
0.636 ± 0.024 ± 0.005		¹ GIBBONS	97B	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
0.591 ± 0.047 ± 0.005		² BORTOLETTO	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
0.492 ± 0.074 ± 0.004		³ ALBRECHT	91H	ARG $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.532 ± 0.065 ± 0.004	21k	⁴ BORTOLETTO	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
0.608 ± 0.183 ± 0.005		⁵ GREEN	83	CLEO Repl. by BORTOLETTO 87


¹ GIBBONS 97B reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0251 \pm 0.0006 \pm 0.00075$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.945 \pm 0.030) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BORTOLETTO 92 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0233 \pm 0.0012 \pm 0.0014$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.945 \pm 0.030) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 91H reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0194 \pm 0.0015 \pm 0.0025$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.945 \pm 0.030) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BORTOLETTO 87 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.0210 \pm 0.0015 \pm 0.0021$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.945 \pm 0.030) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ GREEN 83 reports $[\Gamma(B \rightarrow D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(D^0 \rightarrow K^- \pi^+)] = 0.024 \pm 0.006 \pm 0.004$ which we divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.945 \pm 0.030) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(D^- \ell^+ \nu_\ell \text{ anything})/\Gamma(\bar{D}^0 \ell^+ \nu_\ell \text{ anything})$ Γ_4/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
0.359±0.006±0.009	¹ AAIJ	19AD LHCb	pp at 13 TeV

¹ AAIJ 19AD uses $D^0 \rightarrow K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$ modes.

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.225±0.015 OUR AVERAGE				
0.247±0.019±0.01		¹ GIBBONS 97B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.205±0.019±0.007		² ALBRECHT 96D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.230±0.028±0.009		³ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.283±0.053±0.002		⁴ ALBRECHT 91H	ARG	Sup. by ALBRECHT 96D
0.22 ±0.04 ^{+0.07} _{-0.04}	5200	⁵ BORTOLETTO87	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.27 ±0.06 ^{+0.08} _{-0.06}	510	⁶ CSORNA 85	CLEO	Repl. by BORTOLETTO 87

¹ GIBBONS 97B reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.239 \pm 0.015 \pm 0.014 \pm 0.009$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ALBRECHT 96D reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.196 \pm 0.019$ using CLEO measured $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.681 \pm 0.01 \pm 0.013$, $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$, $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = 0.081 \pm 0.005$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ³ BORTOLETTO 92 reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.25 \pm 0.03 \pm 0.04$ using MARK II $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ and $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.008$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ from a measurement of $[\Gamma(B \rightarrow D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.55 \pm 0.04$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^- \pi^+) = 0.0371 \pm 0.0025$.
- ⁵ BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^- \pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. The product branching ratio for $B(B \rightarrow D^*(2010)^+) B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.
- ⁶ V-A momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0 \pi^+) = 0.6^{+0.08}_{-0.15}$. The product branching fraction is $B(B \rightarrow D^{*+} X) \cdot B(D^{*+} \rightarrow \pi^+ D^0) \cdot B(D^0 \rightarrow K^- \pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

 $\Gamma(\overline{D}^*(2007)^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.260 ± 0.023 ± 0.015	¹ GIBBONS	97B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ GIBBONS 97B reports $B(B \rightarrow \overline{D}^*(2007)^0 \text{ anything}) = 0.247 \pm 0.012 \pm 0.018 \pm 0.018$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(D_s^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.106 ± 0.006 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		

0.1128 ± 0.0003 ± 0.0043		ZHUKOVA	23	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.089 ± 0.010 ± 0.008		¹ ARTUSO	05B	CLE2	$e^+ e^- \rightarrow \Upsilon(5S)$
0.087 ± 0.005 ± 0.008		² AUBERT	02G	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.065 ± 0.011 ± 0.006		³ ALBRECHT	92G	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.068 ± 0.010 ± 0.006	257	⁴ BORTOLETTO90		CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.085 ± 0.022 ± 0.008		⁵ HAAS	86	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.094 ± 0.007 ± 0.008		⁶ GIBAUT	96	CLE2	Repl. by ARTUSO 05B
0.094 ± 0.024 ± 0.008		⁷ ALBRECHT	87H	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ ARTUSO 05B reports $0.0905 \pm 0.0025 \pm 0.0140$ from a measurement of $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ² AUBERT 02G reports $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.00393 \pm 0.00007 \pm 0.00021$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times$

10^{-2} . Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.00292 \pm 0.00039 \pm 0.00031$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.


⁴ BORTOLETTO 90 reports $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.00306 \pm 0.00047$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ HAAS 86 reports $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0038 \pm 0.0010$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $64 \pm 22\%$ decays are 2-body.

⁶ GIBAUT 96 reports $0.1211 \pm 0.0039 \pm 0.0088$ from a measurement of $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷ ALBRECHT 87H reports $[\Gamma(B \rightarrow D_s^\pm \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0042 \pm 0.0009 \pm 0.0006$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $46 \pm 16\%$ of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.

WEIGHTED AVERAGE
 0.106 ± 0.006 (Error scaled by 1.7)



$\Gamma(D_s^{*\pm} \text{ anything})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.063 \pm 0.009 \pm 0.006$	¹ AUBERT	02G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
¹ AUBERT 02G reports $[\Gamma(B \rightarrow D_s^{*\pm} \text{ anything})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.00284 \pm 0.00029 \pm 0.00025$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(D_s^{*\pm} \bar{D}^{(*)})/\Gamma(D_s^{*\pm} \text{ anything})$
Sum over modes Γ_{39}/Γ_{38}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.533 \pm 0.037 \pm 0.037$	AUBERT	02G BABR	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(\bar{D} D_{s0}(2317))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ KROKOVNY	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
¹ The product branching ratio for $B(B \rightarrow \bar{D} D_{s0}(2317)^+) \times B(D_{s0}(2317)^+ \rightarrow D_s \pi^0)$ is measured to be $(8.5^{+2.1}_{-1.9} \pm 2.6) \times 10^{-4}$.			

 $\Gamma(\bar{D} D_{sJ}(2457))/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ KROKOVNY	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
¹ The product branching ratio for $B(B \rightarrow \bar{D} D_{sJ}(2457)^+) \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0, D_s^+ \gamma)$ are measured to be $(17.8^{+4.5}_{-3.9} \pm 5.3) \times 10^{-4}$ and $(6.7^{+1.3}_{-1.2} \pm 2.0) \times 10^{-4}$, respectively.			

 $[\Gamma(D^{(*)} \bar{D}^{(*)} K^0) + \Gamma(D^{(*)} \bar{D}^{(*)} K^\pm)]/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.071^{+0.025+0.010}_{-0.015-0.009}$	¹ BARATE	98Q ALEP	$e^+e^- \rightarrow Z$
¹ The systematic error includes the uncertainties due to the charm branching ratios.			

 $\Gamma(b \rightarrow c \bar{c} s)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.219 ± 0.037	¹ COAN	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
¹ COAN 98 uses D - ℓ correlation.			

 $\Gamma(D_s^{(*)} \bar{D}^{(*)})/\Gamma(D_s^\pm \text{ anything})$
Sum over modes. Γ_{44}/Γ_{37}

VALUE	DOCUMENT ID	TECN	COMMENT
0.469 ± 0.017 OUR AVERAGE			
$0.464 \pm 0.013 \pm 0.015$	AUBERT	02G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.56^{+0.21+0.09}_{-0.15-0.08}$	¹ BARATE	98Q ALEP	$e^+e^- \rightarrow Z$
$0.457 \pm 0.019 \pm 0.037$	GIBAUT	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$0.58 \pm 0.07 \pm 0.09$	ALBRECHT	92G ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.56 ± 0.10	BORTOLETTO	90 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
¹ BARATE 98Q measures $B(B \rightarrow D_s^{(*)} \bar{D}^{(*)}) = 0.056^{+0.021+0.009+0.019}_{-0.015-0.008-0.011}$, where the third error results from the uncertainty on the different D branching ratios and is			

dominated by the uncertainty on $B(D_s^+ \rightarrow \phi\pi^+)$. We divide $B(B \rightarrow D_s^{(*)}\bar{D}^{(*)})$ by our best value of $B(B \rightarrow D_s \text{ anything}) = 0.1 \pm 0.025$.

$\Gamma(D^* D^{*(2010)\pm})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-3}$	90	BARATE	98Q	ALEP $e^+e^- \rightarrow Z$

$[\Gamma(D D^{*(2010)\pm}) + \Gamma(D^* D^\pm)]/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-3}$	90	BARATE	98Q	ALEP $e^+e^- \rightarrow Z$

$\Gamma(D D^\pm)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-3}$	90	BARATE	98Q	ALEP $e^+e^- \rightarrow Z$

$\Gamma(D_s^{(*)\pm}\bar{D}^{(*)}\chi(n\pi^\pm))/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.094^{+0.040+0.034}_{-0.031-0.024}$	¹ BARATE	98Q	ALEP $e^+e^- \rightarrow Z$

¹ The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(\bar{D}^{*(2010)}\gamma)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	¹ LESIAK	92	CBAL $e^+e^- \rightarrow \gamma(4S)$

¹ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}$ Γ_{50}/Γ

Sum over modes.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-4}$	90	¹ ALEXANDER	93B	CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ from a measurement of $[\Gamma(B \rightarrow D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

$\Gamma(D_{s1}(2536)^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{51}/Γ

$D_{s1}(2536)^+$ is the narrow P -wave D_s^+ meson with $J^P = 1^+$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0095	90	¹ BISHAI	98	CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ Assuming factorization, the decay constant $f_{D_{s1}^+}$ is at least a factor of 2.5 times smaller than $f_{D_s^+}$.

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.094 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.1.		
$1.057 \pm 0.012 \pm 0.040$		¹ AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.121 \pm 0.013 \pm 0.042$		ANDERSON	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$1.29 \pm 0.45 \pm 0.01$	27	² MASCHMANN	90 CBAL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.24 \pm 0.27 \pm 0.01$	120	³ ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$1.35 \pm 0.24 \pm 0.01$	52	⁴ ALAM	86 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.12 \pm 0.06 \pm 0.01$	1489	⁵ BALEST	95B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$1.4 \begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix}$	7	⁶ ALBRECHT	85H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$1.1 \pm 0.21 \pm 0.23$	46	⁷ HAAS	85 CLEO	Repl. by ALAM 86

¹ AUBERT 03F also reports the momentum distribution and helicity of $J/\psi \rightarrow \ell^+ \ell^-$ in the $\Upsilon(4S)$ center-of-mass frame.

² MASCHMANN 90 reports $(1.12 \pm 0.33 \pm 0.25) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 87D reports $(1.07 \pm 0.16 \pm 0.22) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .

⁴ ALAM 86 reports $(1.09 \pm 0.16 \pm 0.21) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)]$ assuming $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.074 \pm 0.012$, which we rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ BALEST 95B reports $(1.12 \pm 0.04 \pm 0.06) \times 10^{-2}$ from a measurement of $[\Gamma(B \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0599 \pm 0.0025$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They measure $J/\psi(1S) \rightarrow e^+e^-$ and $\mu^+\mu^-$ and use PDG 1994 values for the branching fractions. The rescaling is the same for either mode so we use e^+e^- .

⁶ Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1$ GeV.

⁷ Dimuon and dielectron events used.

 $\Gamma(J/\psi(1S)(\text{direct})\text{anything})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0078 ± 0.0004 OUR AVERAGE	Error includes scale factor of 1.1.		
$0.00740 \pm 0.00023 \pm 0.00043$	¹ AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.00813 \pm 0.00017 \pm 0.00037$	² ANDERSON	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0080 ± 0.0008	³ BALEST	95B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ AUBERT 03F also reports the helicity of $J/\psi \rightarrow \ell^+ \ell^-$ produced directly in B decay.
- ² Also reports the measurement of $J/\psi \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.
- ³ BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. The $B \rightarrow J/\psi(1S) X$ branching ratio contains $J/\psi(1S)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow J/\psi(1S)$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)$, or $\chi_{c2}(1P) \rightarrow J/\psi(1S)$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow J/\psi(1S)(\text{direct}) X$ branching ratio.

$\Gamma(\psi(2S)\text{anything})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.00307 ± 0.00021 OUR AVERAGE

0.00297 ± 0.00020 ± 0.00020		AUBERT	03F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00316 ± 0.00014 ± 0.00028		¹ ANDERSON	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0046 ± 0.0017 ± 0.0011	8	ALBRECHT	87D	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0034 ± 0.0004 ± 0.0003	240	² BALEST	95B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	-----	---------------------	-----	------	------------------------------------

¹ Also reports the measurement of $\psi(2S) \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.

² BALEST 95B assume PDG 1994 values for sub mode branching ratios. They find $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow \ell^+ \ell^-) = 0.30 \pm 0.05 \pm 0.04$ and $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = 0.37 \pm 0.05 \pm 0.05$. Weighted average is quoted for $B(B \rightarrow \psi(2S)X)$.

$\Gamma(\chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.55 ± 0.27 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

3.33 ± 0.05 ± 0.24		¹ BHARDWAJ	16	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
3.67 ± 0.35 ± 0.44		AUBERT	03F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
4.35 ± 0.29 ± 0.40		ANDERSON	02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.63 ± 0.22 ± 0.34		² ABE	02L	BELL	Repl. by BHARDWAJ 16
3.30 ± 0.35 ± 0.12		³ CHEN	01	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
4.0 ± 0.6 ± 0.4	112	⁴ BALEST	95B	CLE2	Repl. by CHEN 01
10.5 ± 3.5 ± 2.5		⁵ ALBRECHT	92E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.


² ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

³ CHEN 01 reports $0.00414 \pm 0.00031 \pm 0.00040$ from a measurement of $[\Gamma(B \rightarrow \chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ BALEST 95B assume $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma) = (27.3 \pm 1.6) \times 10^{-2}$, the PDG 1994 value. Fit to ψ -photon invariant mass distribution allows for a $\chi_{c1}(1P)$ and a $\chi_{c2}(1P)$ component.

⁵ ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.

WEIGHTED AVERAGE
 3.55 ± 0.27 (Error scaled by 1.3)



$\Gamma(\chi_{c1}(1P)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$

Γ_{56}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.08 ± 0.19 OUR AVERAGE			
$3.03 \pm 0.05 \pm 0.24$	¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$3.41 \pm 0.35 \pm 0.42$	AUBERT 03F	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$3.1 \pm 0.4 \pm 0.1$	² CHEN 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$3.32 \pm 0.22 \pm 0.34$	³ ABE 02L	BELL	Repl. by BHARDWAJ 16
3.7 ± 0.7	⁴ BALEST 95B	CLE2	Repl. by CHEN 01

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² CHEN 01 reports $0.00383 \pm 0.00031 \pm 0.00040$ from a measurement of $[\Gamma(B \rightarrow \chi_{c1}(1P)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

⁴ BALEST 95B assume PDG 1994 values. $J/\psi(1S)$ mesons are reconstructed in the e^+e^- and $\mu^+\mu^-$ modes. The $B \rightarrow \chi_{c1}(1P)X$ branching ratio contains $\chi_{c1}(1P)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow \chi_{c1}(1P)(\text{direct})X$ branching ratio.


$\Gamma(\chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}$

Γ_{57}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
9.9 ± 1.7 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
$9.8 \pm 0.6 \pm 1.0$		¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$21.0 \pm 4.5 \pm 3.1$	AUBERT	03F	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$6.8 \pm 3.4 \pm 0.3$	² CHEN	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$18.0^{+2.3}_{-2.8} \pm 2.6$	³ ABE	02L	BELL	Repl. by BHARDWAJ 16
<38	90	⁴ BALEST	95B	CLE2 Repl. by CHEN 01

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
² CHEN 01 reports $(9.8 \pm 4.8 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(B \rightarrow \chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.5 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
³ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.
⁴ BALEST 95B assume $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma) = (13.5 \pm 1.1) \times 10^{-2}$, the PDG 1994 value. $J/\psi(1S)$ mesons are reconstructed in the e^+e^- and $\mu^+\mu^-$ modes, and PDG 1994 branching fractions are used. If interpreted as signal, the 35 ± 13 events correspond to $B(B \rightarrow \chi_{c2}(1P)X) = (0.25 \pm 0.10 \pm 0.03) \times 10^{-2}$.



$\Gamma(\chi_{c2}(1P)(\text{direct})\text{anything})/\Gamma_{\text{total}}$				Γ_{58}/Γ	
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
0.75 ± 0.11 OUR AVERAGE					
$0.70 \pm 0.06 \pm 0.10$	¹ BHARDWAJ	16	BELL	$e^+e^- \rightarrow \Upsilon(4S)$	
$1.90 \pm 0.45 \pm 0.29$	AUBERT	03F	BABR	$e^+e^- \rightarrow \Upsilon(4S)$	

- • • We do not use the following data for averages, fits, limits, etc. • • •

$1.53^{+0.23}_{-0.28} \pm 0.27$ ² ABE 02L BELL Repl. by BHARDWAJ 16

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

$\Gamma(\eta_c(1S)\text{anything})/\Gamma_{\text{total}}$

Γ_{59}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	¹ BALEST	95B	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+e^-$ and $J/\psi(1S) \rightarrow \mu^+\mu^-$. Search region $2960 < m_{\eta_c(1S)} < 3010$ MeV/ c^2 .

$\Gamma(K\chi_{c1}(3872))/\Gamma_{\text{total}}$

Γ_{60}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.9 ± 0.7 OUR AVERAGE			
$1.7 \pm 0.5 \pm 0.6$	¹ AUSHEV	10	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$2.2^{+0.7+1.2}_{-0.8-1.1}$	^{2,3} GOKHROO	06	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ AUSHEV 10 reports $[\Gamma(B \rightarrow K\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)] = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0) = (46 \pm 16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GOKHROO 06 reports $[\Gamma(B \rightarrow K\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)] = (1.22 \pm 0.31^{+0.23}_{-0.30}) \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0) = (55 \pm 28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Measure the near-threshold enhancements in the $(D^0\bar{D}^0\pi^0)$ system at a mass $3875.2 \pm 0.7^{+0.3}_{-1.6} \pm 0.8$ MeV/ c^2 .

$\Gamma(KX(3940), X \rightarrow D^{*0}D^0)/\Gamma_{\text{total}}$

Γ_{61}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.67	90	AUSHEV	10	BELL $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K\chi_{c0}(3915), \chi_{c0} \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$

Γ_{62}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 3.1$	¹ CHOI	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ CHOI 05 reports the observation of a near-threshold enhancement in the $\omega J/\psi$ mass spectrum in exclusive $B \rightarrow K\omega J/\psi$. The new state, denoted as $\chi_{c0}(3915)$, is measured to have a mass of $3943 \pm 11 \pm 13$ GeV/ c^2 and a width $\Gamma = 87 \pm 22 \pm 26$ MeV.

$\Gamma(K^\pm\text{anything})/\Gamma_{\text{total}}$

Γ_{63}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.789 ± 0.025 OUR AVERAGE			
$0.82 \pm 0.01 \pm 0.05$	ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
$0.775 \pm 0.015 \pm 0.025$	¹ ALBRECHT	93I	ARG $e^+e^- \rightarrow \Upsilon(4S)$
$0.85 \pm 0.07 \pm 0.09$	ALAM	87B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	² BRODY	82	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
seen	³ GIANNINI	82	CUSB	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93I value is not independent of the sum of $B \rightarrow K^+$ anything and $B \rightarrow K^-$ anything ALBRECHT 94C values.

² Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.

³ GIANNINI 82 at CESR-CUSB observed $1.58 \pm 0.35 K^0$ per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{64}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 $\pm$ 0.05	¹ ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.620 $\pm$ 0.013 $\pm$ 0.038	² ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
0.66 $\pm$ 0.05 $\pm$ 0.07	² ALAM	87B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.

² Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$

Γ_{65}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.13 $\pm$ 0.04	¹ ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.165 $\pm$ 0.011 $\pm$ 0.036	² ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
0.19 $\pm$ 0.05 $\pm$ 0.02	² ALAM	87B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.

² Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^0/\bar{K}^0 \text{ anything})/\Gamma_{\text{total}}$

Γ_{66}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 $\pm$ 0.04 OUR AVERAGE			
0.642 $\pm$ 0.010 $\pm$ 0.042	¹ ALBRECHT	94C	ARG $e^+e^- \rightarrow \Upsilon(4S)$
0.63 $\pm$ 0.06 $\pm$ 0.06	ALAM	87B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 94C assume a $K^0/\bar{K}^0$ multiplicity twice that of K_S^0 .

$\Gamma(K^*(892)^\pm \text{ anything})/\Gamma_{\text{total}}$

Γ_{67}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.182 $\pm$ 0.054 $\pm$ 0.024	ALBRECHT	94J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^0/\bar{K}^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$

Γ_{68}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.146 $\pm$ 0.016 $\pm$ 0.020	ALBRECHT	94J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$4.24 \pm 0.54 \pm 0.32$		¹ COAN	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150	90	² LESIAK	92	CBAL $e^+e^- \rightarrow \Upsilon(4S)$
< 24	90	ALBRECHT	88H	ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ An average of $B(B^+ \rightarrow K^*(892)^+\gamma)$ and $B(B^0 \rightarrow K^*(892)^0\gamma)$ measurements reported in COAN 00 by assuming full correlated systematic errors.

² LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(\eta K \gamma)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.5 \pm 1.3^{+1.2}_{-0.9}$	¹ NISHIDA	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ $m_{\eta K} < 2.4 \text{ GeV}/c^2$

$\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 12.7 \times 10^{-5}$	90	¹ COAN	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.6×10^{-3}	90	² LESIAK	92	CBAL $e^+e^- \rightarrow \Upsilon(4S)$
< 4.1×10^{-4}	90	ALBRECHT	88H	ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.66^{+0.59}_{-0.53} \pm 0.13$		¹ COAN	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<83	90	ALBRECHT	88H	ARG $e^+e^- \rightarrow \Upsilon(4S)$
-----	----	----------	-----	---------------------------------------

¹ COAN 00 obtains a fitted signal yield of $15.9^{+5.7}_{-5.2}$ events. A search for contamination by $K^*(1410)$ yielded a rate consistent with 0; the central value assumes no contamination.

$\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-3}$	90	¹ LESIAK	92	CBAL $e^+e^- \rightarrow \Upsilon(4S)$

¹ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.7 \times 10^{-5}$	90	¹ NISHIDA	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.0 \times 10^{-3}$ 90 ALBRECHT 88H ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$.

$\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	¹ LESIAK 92	CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(8.3^{+0.9}_{-0.8} \pm 0.7) \times 10^{-5}$	¹ RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)\eta'(958))/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$4.1^{+1.0}_{-0.9} \pm 0.5$		¹ AUBERT 07E	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<22 90 ¹ RICHICHI 00 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K\eta)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	¹ RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)\eta)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.80^{+0.49}_{-0.43} \pm 0.18) \times 10^{-5}$	¹ RICHICHI 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K\phi\phi)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.3^{+0.9}_{-0.8} \pm 0.3$	¹ HUANG 03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of charged and neutral B meson pairs and isospin symmetry.

$\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.49 ± 0.19 OUR AVERAGE			
$3.75 \pm 0.18 \pm 0.35$	^{1,2} SAITO 15	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$3.52 \pm 0.20 \pm 0.51$	^{1,3} LEES 12U	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$3.32 \pm 0.16 \pm 0.31$	^{1,4} LEES 12V	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$3.47 \pm 0.15 \pm 0.40$	^{1,5} LIMOSANI	09	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.90 \pm 0.91 \pm 0.64$	^{1,6} AUBERT	08O	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.29 \pm 0.44 \pm 0.29$	^{1,7} CHEN	01C	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.30 \pm 0.08 \pm 0.30$	⁸ DEL-AMO-SA..10M	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$	
$4.3 \pm 0.3 \pm 0.7$	⁹ AUBERT	09U	BABR	Repl. by DEL-AMO-SANCHEZ 10M
$3.92 \pm 0.31 \pm 0.47$	^{1,10} AUBERT,BE	06B	BABR	Repl. by LEES 12V
$3.49 \pm 0.20^{+0.59}_{-0.46}$	^{1,11} AUBERT,B	05R	BABR	Repl. by LEES 12U
$3.50 \pm 0.32 \pm 0.31$	^{1,12} KOPPENBURG04	BELL	Repl. by LIMOSANI 09	
$3.36 \pm 0.53^{+0.65}_{-0.68}$	¹³ ABE	01F	BELL	Repl. by SAITO 15
$2.32 \pm 0.57 \pm 0.35$	ALAM	95	CLE2	Repl. by CHEN 01C

¹ We extrapolate the measured value to $E_\gamma > 1.6$ GeV using the method of BUCH-MUELLER 06 (average of three theoretical models).

² SAITO 15 measured $(3.51 \pm 0.17 \pm 0.33) \times 10^{-4}$ using a sum-of-exclusive approach in which 38 of the hadronic final states with $m_{X_s} < 2.8$ GeV/ c^2 are reconstructed. The cut of minimum photon energy is $E_\gamma > 1.9$ GeV.

³ Reports $(3.29 \pm 0.19 \pm 0.48) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

⁴ Reports $(3.21 \pm 0.15 \pm 0.29 \pm 0.08) \times 10^{-4}$ for $1.8 < E_\gamma < 2.8$ GeV, where the last systematic uncertainty is for model dependency. Results with other cutoffs are also reported.

⁵ The measurement reported is $(3.45 \pm 0.15 \pm 0.40) \times 10^{-4}$ for $E_\gamma > 1.7$ GeV.

⁶ Uses a fully reconstructed B meson as a tag on the recoil side. The measurement reported is $(3.66 \pm 0.85 \pm 0.60) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

⁷ The measurement reported is $(3.21 \pm 0.43^{+0.32}_{-0.29}) \times 10^{-4}$ for $E_\gamma > 2.0$ GeV.

⁸ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)} \gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.5–2.0 GeV/ c^2 .

⁹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)} \gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.6–1.8 GeV/ c^2 .

¹⁰ The measurement reported is $(3.67 \pm 0.29 \pm 0.45) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

¹¹ The measurement reported is $(3.27 \pm 0.18^{+0.55}_{-0.42}) \times 10^{-4}$ for $E_\gamma > 1.9$ GeV.

¹² The measurement reported is $(3.55 \pm 0.32 \pm 0.32) \times 10^{-4}$ for $E_\gamma > 1.8$ GeV.

¹³ ABE 01F reports their systematic errors $(\pm 0.42^{+0.50}_{-0.54}) \times 10^{-4}$, where the second error is due to the theoretical uncertainty. We combine them in quadrature.

$\Gamma(\bar{b} \rightarrow \bar{d} \gamma) / \Gamma_{\text{total}}$

Γ_{82} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$9.2 \pm 2.0 \pm 2.3$	¹ DEL-AMO-SA..10M	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14 \pm 5 \pm 4$	² AUBERT	09U	BABR	Repl. by DEL-AMO-SANCHEZ 10M
------------------	---------------------	-----	------	------------------------------

¹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)} \gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.5–2.0 GeV/ c^2 .

² Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)} \gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.6–1.8 GeV/ c^2 .

$\Gamma(\bar{b} \rightarrow \bar{d}\gamma)/\Gamma(\bar{b} \rightarrow \bar{s}\gamma)$ Γ_{82}/Γ_{81}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.040 \pm 0.009 \pm 0.010$	¹ DEL-AMO-SA...10M	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.033 \pm 0.013 \pm 0.009$	² AUBERT	09U BABR	Repl. by DEL-AMO-SANCHEZ 10M

¹ Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.5–2.0 GeV/c².

² Measured using sums of seven exclusive final states $B \rightarrow X_{d(s)}\gamma$ where $X_{d(s)}$ is a nonstrange (strange) charmless hadronic system in mass range 0.6–1.8 GeV/c².

 $\Gamma(\bar{b} \rightarrow \bar{s}\text{gluon})/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.068	90		¹ COAN	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.08	2		² ALBRECHT	95D ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ COAN 98 uses D - ℓ correlation.

² ALBRECHT 95D use full reconstruction of one B decay as tag. Two candidate events for charmless B decay can be interpreted as either $b \rightarrow s\text{gluon}$ or $b \rightarrow u$ transition. If interpreted as $b \rightarrow s\text{gluon}$ they find a branching ratio of ~ 0.026 or the upper limit quoted above. Result is highly model dependent.

 $\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$2.61 \pm 0.30^{+0.44}_{-0.74}$		¹ NISHIMURA	10 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.69 \pm 0.29^{+0.36}_{-0.62}$		² NISHIMURA	10 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
<4.4	90	³ BROWDER	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $B \rightarrow \eta X_s$ with $0.4 < m_{X_s} < 2.6$ GeV/c².

² Uses $B \rightarrow \eta X_s$ with $1.8 < m_{X_s} < 2.6$ GeV/c².

³ BROWDER 98 search for high momentum $B \rightarrow \eta X_s$ between 2.1 and 2.7 GeV/c.

 $\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.9 OUR AVERAGE			
$3.9 \pm 0.8 \pm 0.9$	¹ AUBERT,B	04F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.6 \pm 1.1 \pm 0.6$	² BONVICINI	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$6.2 \pm 1.6^{+1.3}_{-2.0}$	³ BROWDER	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT,B 04F reports branching ratio $B \rightarrow \eta' X_s$ for high momentum η' between 2.0 and 2.7 GeV/c in the $\Upsilon(4S)$ center-of-mass frame. X_s represents a recoil system consisting of a kaon and zero to four pions.

² BONVICINI 03 observed a signal of 61.2 ± 13.9 events in $B \rightarrow \eta' X_{nc}$ production for high momentum η' between 2.0 and 2.7 GeV/c in the $\Upsilon(4S)$ center-of-mass frame. The

X_{nc} denotes “charmless” hadronic states recoiling against η' . The second error combines systematic and background subtraction uncertainties in quadrature.

³ BROWDER 98 observed a signal of 39.0 ± 11.6 events in high momentum $B \rightarrow \eta' X_S$ production between 2.0 and 2.7 GeV/c. The branching fraction is based on the interpretation of $b \rightarrow sg$, where the last error includes additional uncertainties due to the color-suppressed $b \rightarrow$ backgrounds.

$\Gamma(K^+ \text{ gluon (charmless)})/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.87	90	¹ DEL-AMO-SA...11	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ $B \rightarrow K^+ X$ with $m_X < 1.69 \text{ GeV}/c^2$.

$\Gamma(K^0 \text{ gluon (charmless)})/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.95^{+0.51}_{-0.45} \pm 0.50$	¹ DEL-AMO-SA...11	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ $B \rightarrow K^0 X$ with $m_X < 1.69 \text{ GeV}/c^2$.

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.39 ± 0.25 OUR AVERAGE	Error includes scale factor of 1.2.			
$1.73^{+0.34}_{-0.32} \pm 0.17$	1,2	AUBERT	08BH BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.21^{+0.24}_{-0.22} \pm 0.12$	1,2	TANIGUCHI	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.36^{+0.29}_{-0.27} \pm 0.10$	1,3	AUBERT	07L BABR	Repl. by AUBERT 08BH
< 1.9	90	1,3 AUBERT	04C BABR	Repl. by AUBERT 07L
<14	90	1,4 COAN	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$.

³ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.

⁴ COAN 00 reports $B(B \rightarrow \rho\gamma)/B(B \rightarrow K^*(892)\gamma) < 0.32$ at 90%CL and scaled by the central value of $B(B \rightarrow K^*(892)\gamma) = (4.24 \pm 0.54 \pm 0.32) \times 10^{-5}$.

$\Gamma(\rho\gamma)/\Gamma(K^*(892)\gamma)$ Γ_{88}/Γ_{69}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.02^{+0.60+0.26}_{-0.55-0.28}$	TANIGUCHI	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho/\omega\gamma)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.30 ± 0.23 OUR AVERAGE	Error includes scale factor of 1.2.			
$1.63^{+0.30}_{-0.28} \pm 0.16$	1,2,3	AUBERT	08BH BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.14 \pm 0.20^{+0.10}_{-0.12}$	1,3	TANIGUCHI	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$1.25^{+0.25}_{-0.24} \pm 0.09$	⁴ AUBERT	07L	BABR	Repl. by AUBERT 08BH
$1.32^{+0.34+0.10}_{-0.31-0.09}$	⁴ MOHAPATRA	06	BELL	Repl. by TANIGUCHI 08
$0.6 \pm 0.3 \pm 0.1$	⁴ AUBERT	05	BABR	Repl. by AUBERT 07L
<1.4	⁴ MOHAPATRA	05	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$.				
² Also reports $ V_{td}/V_{ts} = 0.233^{+0.025+0.022}_{-0.024-0.021}$.				
³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
⁴ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.				

$\Gamma(\rho/\omega\gamma)/\Gamma(K^*(892)\gamma)$

Γ_{89}/Γ_{69}

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
$2.84 \pm 0.50^{+0.27}_{-0.29}$		¹ TANIGUCHI	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

<3.5	90	MOHAPATRA	05	BELL Repl. by TANIGUCHI 08
¹ Also reports $ V_{td}/V_{ts} = 0.195^{+0.020}_{-0.019} \pm 0.015$.				

$\Gamma(\pi^\pm \text{ anything})/\Gamma_{\text{total}}$

Γ_{90}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$3.585 \pm 0.025 \pm 0.070$	¹ ALBRECHT	93I	ARG $e^+e^- \rightarrow \Upsilon(4S)$
¹ ALBRECHT 93 excludes $\pi^\pm$ from K_S^0 and Λ decays. If included, they find $4.105 \pm 0.025 \pm 0.080$.			

$\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$

Γ_{91}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$2.35 \pm 0.02 \pm 0.11$	¹ ABE	01J	BELL $e^+e^- \rightarrow \Upsilon(4S)$
¹ From fully inclusive π^0 yield with no corrections from decays of K_S^0 or other particles.			

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$

Γ_{92}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.176 \pm 0.011 \pm 0.012$	KUBOTA	96	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho^0 \text{ anything})/\Gamma_{\text{total}}$

Γ_{93}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.208 \pm 0.042 \pm 0.032$	ALBRECHT	94J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$

Γ_{94}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.81	90	ALBRECHT	94J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

Γ_{95}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0343 ± 0.0012 OUR AVERAGE			
$0.0353 \pm 0.0005 \pm 0.0030$	HUANG	07	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

$0.0341 \pm 0.0006 \pm 0.0012$	AUBERT	04S	BABR	$e^+e^- \rightarrow \gamma(4S)$
$0.0390 \pm 0.0030 \pm 0.0035$	ALBRECHT	94J	ARG	$e^+e^- \rightarrow \gamma(4S)$
$0.023 \pm 0.006 \pm 0.005$	BORTOLETTO	86	CLEO	$e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	¹ BERGFELD	98	CLE2

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\pi^+ \text{ gluon (charmless)})/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.72^{+0.50}_{-0.47} \pm 0.59$	¹ DEL-AMO-SA...11	BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ $B \rightarrow \pi^+ X$ with $m_X < 1.71 \text{ GeV}/c^2$.

$\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
$3.55 \pm 0.32 \pm 0.14$		¹ AUBERT	07C	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$6.4 \pm 0.8 \pm 0.8$		² CRAWFORD	92	CLEO $e^+e^- \rightarrow \gamma(4S)$
14 ± 9		³ ALBRECHT	88E	ARG $e^+e^- \rightarrow \gamma(4S)$
<11.2	90	⁴ ALAM	87	CLEO $e^+e^- \rightarrow \gamma(4S)$

¹ AUBERT 07C reports $0.045 \pm 0.003 \pm 0.012$ from a measurement of $[\Gamma(B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$, which we rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.35 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and $B^\pm$ decay are Λ_c .

³ ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.

⁴ Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.

$\Gamma(\Lambda_c^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ $\Gamma_{100}/\Gamma_{101}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.19 \pm 0.13 \pm 0.04$	¹ AMMAR	97	CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\bar{\Lambda}_c^- \mu^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ $\Gamma_{104}/\Gamma_{101}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$-2.0 \pm 2.0 \pm 1.9$	LEES	12	BABR $e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{\Lambda}_c^- \ell^+ \text{ anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})$ Γ_{102}/Γ_{99}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-2}$	90	¹ LEES	12	BABR $e^+e^- \rightarrow \gamma(4S)$

¹LEES 12 quotes also the measurement $\Gamma(B \rightarrow \bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^- \text{anything}) = (1.2 \pm 0.7 \pm 0.4) \times 10^{-2}$.

$\Gamma(\bar{\Lambda}_c^- e^+ \text{anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{anything})$ Γ_{103}/Γ_{99}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	¹ BONVICINI 98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

$\Gamma(\bar{\Lambda}_c^- e^+ \text{anything})/\Gamma(\bar{\Lambda}_c^- \text{anything})$ $\Gamma_{103}/\Gamma_{101}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 1.1 \pm 0.6$	¹ LEES 12	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹Uses the full reconstruction of the recoiling B in a hadronic decay as a tag.

$\Gamma(\bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(\bar{\Lambda}_c^- \text{anything})$ $\Gamma_{102}/\Gamma_{101}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.5 \times 10^{-2}$	90	¹ LEES 12	BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹LEES 12 quotes also the measurement $\Gamma(B \rightarrow \bar{\Lambda}_c^- \ell^+ \text{anything})/\Gamma(B \rightarrow \bar{\Lambda}_c^- \text{anything}) = (1.7 \pm 1.0 \pm 0.6) \times 10^{-2}$.

$\Gamma(\bar{\Lambda}_c^- p \text{anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{anything})$ Γ_{105}/Γ_{99}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.57 \pm 0.05 \pm 0.05$	BONVICINI 98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{\Lambda}_c^- p e^+ \nu_e)/\Gamma(\bar{\Lambda}_c^- p \text{anything})$ $\Gamma_{106}/\Gamma_{105}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	¹ BONVICINI 98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

$\Gamma(\bar{\Sigma}_c^{--} \text{anything})/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.0033 \pm 0.0017 \pm 0.0001$	77	¹ PROCARIO 94	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^{--} \text{anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.35 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^- \text{anything})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-3}$	90	¹ PROCARIO 94	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^- \text{anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] < 0.00048$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 6.35 \times 10^{-2}$.

$\Gamma(\bar{\Sigma}_c^0 \text{anything})/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.0036 \pm 0.0017 \pm 0.0001$	76	¹ PROCARIO 94	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

¹ PROCARIO 94 reports $[\Gamma(B \rightarrow \bar{\Sigma}_c^0 \text{ anything})/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$ which we divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.35 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^0 N(N = p \text{ or } n))/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻³	90	¹ PROCARIO 94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ PROCARIO 94 reports < 0.0017 from a measurement of $[\Gamma(B \rightarrow \bar{\Sigma}_c^0 N(N = p \text{ or } n))/\Gamma_{\text{total}}] \times [B(\Lambda_c^+ \rightarrow p K^- \pi^+)]$ assuming $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$, which we rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 6.35 \times 10^{-2}$.

$\Gamma(\Xi_c^0 \text{ anything}, \Xi_c^0 \rightarrow \Xi^- \pi^+)/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.030 OUR AVERAGE	Error includes scale factor of 1.1.		
0.211 ± 0.019 ± 0.025	¹ AUBERT, B	05M	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
0.144 ± 0.048 ± 0.021	² BARISH	97	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ The yield is obtained by requiring the momentum $P < 2.15$ GeV/c.

² BARISH 97 find $79 \pm 27 \Xi_c^0$ events.

$\Gamma(\Xi_c^+, \Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
0.453 ± 0.096^{+0.085}_{-0.065}	¹ BARISH	97	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ BARISH 97 find $125 \pm 28 \Xi_c^+$ events.

$\Gamma(p/\bar{p} \text{ anything})/\Gamma_{\text{total}}$ Γ_{113}/Γ

Includes p and $\bar{p}$ from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.004 OUR AVERAGE				
0.080 ± 0.005 ± 0.005		ALBRECHT 93I	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.080 ± 0.005 ± 0.003		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.082 ± 0.005 ^{+0.013} _{-0.010}	2163	¹ ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.021 ² ALAM 83B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 89K include direct and nondirect protons.

² ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and $\bar{p}$. Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$\Gamma(p/\bar{p} \text{ (direct) anything})/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.055 ± 0.005 OUR AVERAGE				
0.055 ± 0.005 ± 0.0035		ALBRECHT 93I	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.056 ± 0.006 ± 0.005		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.055 ± 0.016	1220	¹ ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$\Gamma(\bar{p}e^+\nu_e \text{ anything})/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.9 \times 10^{-4}$	90	¹ ADAM 03B	CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 16 \times 10^{-4}$	90	ALBRECHT 90H	ARG	$e^+e^- \rightarrow \gamma(4S)$

¹Based on $V-A$ model. $\Gamma(\Lambda/\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.005 OUR AVERAGE				
0.038 ± 0.004 ± 0.006	2998	CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
0.042 ± 0.005 ± 0.006	943	ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.022 ± 0.003 ± 0.0022		¹ ACKERSTAFF 97N	OPAL	$e^+e^- \rightarrow Z$
> 0.011		² ALAM 83B	CLEO	$e^+e^- \rightarrow \gamma(4S)$
¹ ACKERSTAFF 97N assumes $B(b \rightarrow B) = 0.868 \pm 0.041$, i.e., an admixture of B^0 , $B^\pm$, and B_s .				
² ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and $\bar{p}$. Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.				

 $\Gamma(\Lambda \text{ anything})/\Gamma(\bar{\Lambda} \text{ anything})$ $\Gamma_{117}/\Gamma_{118}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.43 ± 0.09 ± 0.07	¹ AMMAR 97	CLE2	$e^+e^- \rightarrow \gamma(4S)$
¹ AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).			

 $\Gamma(\Xi^-/\Xi^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027 ± 0.0006 OUR AVERAGE				
0.0027 ± 0.0005 ± 0.0004	147	CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
0.0028 ± 0.0014	54	ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.068 ± 0.005 ± 0.003	¹ ALBRECHT 92O	ARG	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.076 ± 0.014	² ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$
¹ ALBRECHT 92O result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{\Lambda}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.			
² ALBRECHT 89K obtain this result by adding their their measurements $(5.5 \pm 1.6)\%$ for direct protons and $(4.2 \pm 0.5 \pm 0.6)\%$ for inclusive Λ production. They then assume $(5.5 \pm 1.6)\%$ for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain $(7.6 \pm 1.4)\%$.			

 $\Gamma(p\bar{p} \text{ anything})/\Gamma_{\text{total}}$ Γ_{121}/Γ Includes p and $\bar{p}$ from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0247 ± 0.0023 OUR AVERAGE				
0.024 ± 0.001 ± 0.004		CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
0.025 ± 0.002 ± 0.002	918	ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$

$\Gamma(p\bar{p} \text{ anything})/\Gamma(p/\bar{p} \text{ anything})$ $\Gamma_{121}/\Gamma_{113}$ Includes p and $\bar{p}$ from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.30 \pm 0.02 \pm 0.05$	¹ CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
--------------------------	--------------------------	------	---------------------------------

¹ CRAWFORD 92 value is not independent of their $\Gamma(p\bar{p} \text{ anything})/\Gamma_{\text{total}}$ value. $\Gamma(\Lambda\bar{p}/\bar{\Lambda}p \text{ anything})/\Gamma_{\text{total}}$ Γ_{122}/Γ Includes p and $\bar{p}$ from Λ and $\bar{\Lambda}$ decay.

VALUE	EVT5	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.025 ± 0.004 OUR AVERAGE

$0.029 \pm 0.005 \pm 0.005$	CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
-----------------------------	-------------	------	---------------------------------

$0.023 \pm 0.004 \pm 0.003$	165 ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$
-----------------------------	------------------	-----	---------------------------------

 $\Gamma(\Lambda\bar{p}/\bar{\Lambda}p \text{ anything})/\Gamma(\Lambda/\bar{\Lambda} \text{ anything})$ $\Gamma_{122}/\Gamma_{116}$ Includes p and $\bar{p}$ from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.76 \pm 0.11 \pm 0.08$	¹ CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
--------------------------	--------------------------	------	---------------------------------

¹ CRAWFORD 92 value is not independent of their $[\Gamma(\Lambda\bar{p} \text{ anything}) + \Gamma(\bar{\Lambda}p \text{ anything})]/\Gamma_{\text{total}}$ value. $\Gamma(\Lambda\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE	CL%	EVT5	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

< 0.005	90	CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
-------------------	----	-------------	------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.0088	90	12 ALBRECHT 89K	ARG	$e^+e^- \rightarrow \gamma(4S)$
------------	----	-----------------	-----	---------------------------------

 $\Gamma(\Lambda\bar{\Lambda} \text{ anything})/\Gamma(\Lambda/\bar{\Lambda} \text{ anything})$ $\Gamma_{123}/\Gamma_{116}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.13	90	¹ CRAWFORD 92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
----------	----	--------------------------	------	---------------------------------

¹ CRAWFORD 92 value is not independent of their $\Gamma(\Lambda\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$ value. $\Gamma(se^+e^-)/\Gamma_{\text{total}}$ Γ_{124}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

6.7 ± 1.7 OUR AVERAGE Error includes scale factor of 2.0.

$7.69^{+0.82+0.71}_{-0.77-0.60}$	¹ LEES	14D	BABR	$e^+e^- \rightarrow \gamma(4S)$
----------------------------------	-------------------	-----	------	---------------------------------

$4.04 \pm 1.30^{+0.87}_{-0.83}$	² IWASAKI	05	BELL	$e^+e^- \rightarrow \gamma(4S)$
---------------------------------	----------------------	----	------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.0 \pm 1.7 \pm 1.3$	² AUBERT,B	04i	BABR	Repl. by LEES 14D
-----------------------	-----------------------	-----	------	-------------------

$5.0 \pm 2.3^{+1.3}_{-1.1}$	² KANEKO	03	BELL	Repl. by IWASAKI 05
-----------------------------	---------------------	----	------	---------------------

< 57	90	GLENN	98	CLEO $e^+e^- \rightarrow \gamma(4S)$
--------	----	-------	----	--------------------------------------

< 50000	90	BEBEK	81	CLEO $e^+e^- \rightarrow \gamma(4S)$
-----------	----	-------	----	--------------------------------------

¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, K_S^0 , $K_S^0\pi^0$, $K_S^0\pi^+$, $K_S^0\pi^+\pi^0$, and $K_S^0\pi + \pi^-$ corrected for unobserved modes.

² Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(s\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{125}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.3 $\pm$ 1.0 OUR AVERAGE				
4.41 ^{+1.31+0.63} _{-1.17-0.50}		¹ LEES	14D	BABR $e^+e^- \rightarrow \gamma(4S)$
4.13 $\pm$ 1.05 ^{+0.85} _{-0.81}		² IWASAKI	05	BELL $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.0 $\pm$ 2.8 $\pm$ 1.2		AUBERT,B	04i	BABR Repl. by LEES 14D
7.9 $\pm$ 2.1 ^{+2.1} _{-1.5}		KANEKO	03	BELL Repl. by IWASAKI 05
< 58	90	GLENN	98	CLEO $e^+e^- \rightarrow \gamma(4S)$
<17000	90	CHADWICK	81	CLEO $e^+e^- \rightarrow \gamma(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, K_S^0 , $K_S^0\pi^0$, $K_S^0\pi^+$, $K_S^0\pi^+\pi^0$, and $K_S^0\pi + \pi^-$ corrected for unobserved modes.

² Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.

$[\Gamma(se^+e^-) + \Gamma(s\mu^+\mu^-)]/\Gamma_{\text{total}}$

$(\Gamma_{124} + \Gamma_{125})/\Gamma$

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.2 $\times 10^{-5}$	90	GLENN	98	CLEO $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0024	90	¹ BEAN	87	CLEO Repl. by GLENN 98
<0.0062	90	² AVERY	84	CLEO Repl. by BEAN 87

¹ BEAN 87 reports $[(\mu^+\mu^-) + (e^+e^-)]/2$ and we converted it.

² Determine ratio of B^+ to B^0 semileptonic decays to be in the range 0.25–2.9.

$\Gamma(sl^+\ell^-)/\Gamma_{\text{total}}$

Γ_{126}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.8 $\pm$ 1.3 OUR AVERAGE	Error includes scale factor of 1.8.		
6.73 ^{+0.70+0.60} _{-0.64-0.56}	¹ LEES	14D	BABR $e^+e^- \rightarrow \gamma(4S)$
4.11 $\pm$ 0.83 ^{+0.85} _{-0.81}	² IWASAKI	05	BELL $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.6 $\pm$ 1.5 $\pm$ 1.3	³ AUBERT,B	04i	BABR Repl. by LEES 14D
6.1 $\pm$ 1.4 ^{+1.4} _{-1.1}	³ KANEKO	03	BELL Repl. by IWASAKI 05

¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, K_S^0 , $K_S^0\pi^0$, $K_S^0\pi^+$, $K_S^0\pi^+\pi^0$, and $K_S^0\pi + \pi^-$ corrected for unobserved modes.

² Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.

³ Requires $M_{e^+e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(\pi\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-8}$	90	¹ LEES	13M BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.2 \times 10^{-8}$	90	¹ WEI	08A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<9.1 \times 10^{-8}$	90	¹ AUBERT	07AG BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\pi e^+e^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.0 \times 10^{-8}$	90	¹ LEES	13M BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\pi\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0 \times 10^{-8}$	90	¹ LEES	13M BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K e^+e^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
4.4 ± 0.6 OUR AVERAGE				
$3.9^{+0.9}_{-0.8} \pm 0.2$		¹ AUBERT	09T BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$4.8^{+0.8}_{-0.7} \pm 0.3$		¹ WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.3^{+0.9}_{-0.8} \pm 0.2$		¹ AUBERT,B	06J BABR	Repl. by AUBERT 09T
$7.4^{+1.8}_{-1.6} \pm 0.5$		¹ AUBERT	03U BABR	Repl. by AUBERT,B 06J
$4.8^{+1.5}_{-1.3} \pm 0.3$		^{1,2} ISHIKAWA	03 BELL	Repl. by WEI 09A
<13	90	ABE	02 BELL	Repl. by ISHIKAWA 03

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The second error is a total of systematic uncertainties including model dependence.

 $\Gamma(K^*(892)e^+e^-)/\Gamma_{\text{total}}$ Γ_{131}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
11.9 ± 2.0 OUR AVERAGE		Error includes scale factor of 1.2.		
$9.9^{+2.3}_{-2.1} \pm 0.6$		¹ AUBERT	09T BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$13.9^{+2.3}_{-2.0} \pm 1.2$		¹ WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.7^{+3.0}_{-2.7} \pm 1.4$		¹ AUBERT,B	06J BABR	Repl. by AUBERT 09T
$9.8^{+5.0}_{-4.2} \pm 1.1$		¹ AUBERT	03U BABR	Repl. by AUBERT,B 06J

$14.9^{+5.2+1.2}_{-4.6-1.3}$	² ISHIKAWA	03	BELL	Repl. by WEI 09A
<56	90	ABE	02	BELL Repl. by ISHIKAWA 03

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{132}/Γ
 Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
4.4 ± 0.4 OUR AVERAGE			
$4.2 \pm 0.4 \pm 0.2$	AALTONEN	11A1 CDF	$p\bar{p}$ at 1.96 TeV
$4.1^{+1.3}_{-1.2} \pm 0.2$	¹ AUBERT	09T BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$5.0 \pm 0.6 \pm 0.3$	¹ WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$3.5^{+1.3}_{-1.1} \pm 0.3$	¹ AUBERT,B	06J BABR	Repl. by AUBERT 09T
$4.5^{+2.3}_{-1.9} \pm 0.4$	¹ AUBERT	03U BABR	Repl. by AUBERT,B 06J
$4.8^{+1.2}_{-1.1} \pm 0.4$	^{1,2} ISHIKAWA	03 BELL	Repl. by WEI 09A
$9.9^{+4.0+1.3}_{-3.2-1.0}$	ABE	02 BELL	Repl. by ISHIKAWA 03

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K\mu^+\mu^-)/\Gamma(K e^+ e^-)$ $\Gamma_{132}/\Gamma_{130}$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.01^{+0.19}_{-0.16}$ OUR AVERAGE			
$1.03^{+0.28}_{-0.24} \pm 0.01$	¹ CHOUDHURY 21	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.00^{+0.31}_{-0.25} \pm 0.07$	² LEES	12S BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.96^{+0.44}_{-0.34} \pm 0.05$	AUBERT	09T BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.03 \pm 0.19 \pm 0.06$	³ WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.06 \pm 0.48 \pm 0.08$	AUBERT,B	06J BABR	Repl. by AUBERT 09T

¹ For $1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$. Measurements in other q^2 bins are also reported.

² Measured in the union of $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $q^2 > 10.11 \text{ GeV}^2/c^4$.
 LEES 12S reports also individual measurements $\Gamma(B \rightarrow K\mu^+\mu^-)/\Gamma(B \rightarrow Ke^+e^-)$
 $= 0.74^{+0.40}_{-0.31} \pm 0.06$ for $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $\Gamma(B \rightarrow K\mu^+\mu^-)/\Gamma(B \rightarrow Ke^+e^-)$
 $= 1.43^{+0.65}_{-0.44} \pm 0.12$ for $q^2 > 10.11 \text{ GeV}^2/c^4$.

³ Superseded by CHOUDHURY 21.

$\Gamma(K^*(892)\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ
 Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
10.6 ± 0.9 OUR AVERAGE				
$10.1 \pm 1.0 \pm 0.5$		AALTONEN	11A1 CDF	$p\bar{p}$ at 1.96 TeV

$13.5^{+3.5}_{-3.3} \pm 1.0$	¹ AUBERT	09T	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$11.0^{+1.6}_{-1.4} \pm 0.8$	¹ WEI	09A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$8.8^{+3.5}_{-3.0} \pm 1.2$	¹ AUBERT,B	06J	BABR	Repl. by AUBERT 09T
$12.7^{+7.6}_{-6.1} \pm 1.6$	¹ AUBERT	03U	BABR	Repl. by AUBERT,B 06J
$11.7^{+3.6}_{-3.1} \pm 1.0$	² ISHIKAWA	03	BELL	Repl. by WEI 09A
<31	90	ABE	02	BELL Repl. by ISHIKAWA 03
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.				

$\Gamma(K^*(892)\mu^+\mu^-)/\Gamma(K^*(892)e^+e^-)$ $\Gamma_{133}/\Gamma_{131}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.98±0.15 OUR AVERAGE			
$1.13^{+0.34}_{-0.26} \pm 0.10$	¹ LEES	12S	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$1.37^{+0.53}_{-0.40} \pm 0.09$	AUBERT	09T	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$0.83 \pm 0.17 \pm 0.08$	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.91 \pm 0.45 \pm 0.06$	AUBERT,B	06J	BABR Repl. by AUBERT 09T

¹ Measured in the union of $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $q^2 > 10.11 \text{ GeV}^2/c^4$. LEES 12S reports also individual measurements $\Gamma(B \rightarrow K^*(892)\mu^+\mu^-)/\Gamma(B \rightarrow K^*(892)e^+e^-) = 1.06^{+0.48}_{-0.33} \pm 0.08$ for $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $\Gamma(B \rightarrow K^*(892)\mu^+\mu^-)/\Gamma(B \rightarrow K^*(892)e^+e^-) = 1.18^{+0.55}_{-0.37} \pm 0.11$ for $q^2 > 10.11 \text{ GeV}^2/c^4$.

$\Gamma(K\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{134}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
4.8±0.4 OUR AVERAGE				
$4.7 \pm 0.6 \pm 0.2$		LEES	12S	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$4.8^{+0.5}_{-0.4} \pm 0.3$		WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.9 \pm 0.7 \pm 0.2$		¹ AUBERT	09T	BABR Repl. by LEES 12S
$3.4 \pm 0.7 \pm 0.2$		¹ AUBERT,B	06J	BABR Repl. by AUBERT 09T
$6.5^{+1.4}_{-1.3} \pm 0.4$		² AUBERT	03U	BABR Repl. by AUBERT,B 06J
$4.8^{+1.0}_{-0.9} \pm 0.3$		³ ISHIKAWA	03	BELL Repl. by WEI 09A
$7.5^{+2.5}_{-2.1} \pm 0.6$		⁴ ABE	02	BELL Repl. by ISHIKAWA 03
< 5.1	90	¹ AUBERT	02L	BABR $e^+e^- \rightarrow \Upsilon(4S)$
< 17	90	⁵ ANDERSON	01B	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes all four $B \rightarrow K \ell^+ \ell^-$ modes having equal partial widths in the fit.

³ Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K \ell^+ \ell^-$, and $B(B \rightarrow K^*(892) \mu^+ \mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.

⁴ Assumes lepton universality.

⁵ The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892) \ell^+ \ell^-) / \Gamma_{\text{total}}$ Γ_{135} / Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

10.5 ± 1.0 OUR AVERAGE

10.2^{+1.4}_{-1.3} ± 0.5 LEES 12S BABR $e^+ e^- \rightarrow \gamma(4S)$

10.7^{+1.1}_{-1.0} ± 0.9 WEI 09A BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.1^{+1.9}_{-1.8} ± 0.7 ¹ AUBERT 09T BABR Repl. by LEES 12S

7.8^{+1.9}_{-1.7} ± 1.1 ¹ AUBERT,B 06J BABR Repl. by AUBERT 09T

8.8^{+3.3}_{-2.9} ± 1.0 ² AUBERT 03U BABR Repl. by AUBERT,B 06J

11.5^{+2.6}_{-2.4} ± 0.8 ³ ISHIKAWA 03 BELL Repl. by WEI 09A

<31 90 ^{1,4} AUBERT 02L BABR Repl. by AUBERT 03U

<33 90 ⁵ ANDERSON 01B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Assumes the partial width ratio of electron and muon modes to be $\Gamma(B \rightarrow K^*(892) e^+ e^-) / \Gamma(B \rightarrow K^*(892) \mu^+ \mu^-) = 1.33$.

³ Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K \ell^+ \ell^-$, and $B(B \rightarrow K^*(892) \mu^+ \mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.

⁴ For averaging $K^*(892) \mu^+ \mu^-$ and $K^*(892) e^+ e^-$ modes, AUBERT 02L assumed $B(B \rightarrow K^*(892) e^+ e^-) / B(B \rightarrow K^*(892) \mu^+ \mu^-) = 1.2$.

⁵ The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K \nu \bar{\nu}) / \Gamma_{\text{total}}$ Γ_{136} / Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.6 × 10⁻⁵ 90 ¹ GRYGIER 17 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 × 10⁻⁵ 90 ^{1,2} LEES 13I BABR $e^+ e^- \rightarrow \gamma(4S)$

<1.4 × 10⁻⁵ 90 ¹ DEL-AMO-SA..10Q BABR Repl. by LEES 13I

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² Also reported a limit < 3.2 × 10⁻⁵ at 90% CL obtained using a fully reconstructed hadronic B -tag evnets.

$\Gamma(K^* \nu \bar{\nu}) / \Gamma_{\text{total}}$ Γ_{137} / Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.7 × 10⁻⁵ 90 ¹ GRYGIER 17 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<7.6 \times 10^{-5}$	90	^{1,2} LEES	13I	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<8 \times 10^{-5}$	90	AUBERT	08BC	BABR	Repl. by LEES 13I

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Also reported a limit $< 7.9 \times 10^{-5}$ at 90% CL obtained using a fully reconstructed hadronic B -tag events.

$\Gamma(\pi\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.8 \times 10^{-5}$	90	¹ GRYGIER 17	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-5}$	90	¹ GRYGIER 17	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(se^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{140}/Γ

Test for lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	GLENN 98	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\pi e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{141}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.2 \times 10^{-8}$	90	¹ AUBERT 07AG	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.6 \times 10^{-6}$	90	¹ EDWARDS 02B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	--------------------------	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{142}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-6}$	90	¹ EDWARDS 02B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(Ke^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{143}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.38	90	¹ AUBERT,B 06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	¹ EDWARDS 02B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-------	----	--------------------------	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{144}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.1	90	¹ AUBERT,B 06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<62 90 ¹ EDWARDS 02B CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

CP VIOLATION

A_{CP} is defined as

$$\frac{B(\bar{B} \rightarrow \bar{f}) - B(B \rightarrow f)}{B(\bar{B} \rightarrow \bar{f}) + B(B \rightarrow f)},$$

the CP -violation charge asymmetry of inclusive $B^\pm$ and B^0 decay.

$A_{CP}(B \rightarrow K^*(892)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

−0.003±0.011 OUR AVERAGE

−0.004±0.014±0.003	¹ HORIGUCHI	17	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
−0.003±0.017±0.007	² AUBERT	09AO	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.08 ±0.13 ±0.03	³ COAN	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

−0.013±0.036±0.010	⁴ AUBERT,BE	04A	BABR	Repl. by AUBERT 09AO
−0.015±0.044±0.012	³ NAKAO	04	BELL	Repl. by HORIGUCHI 17
−0.044±0.076±0.012	⁵ AUBERT	02C	BABR	Repl. by AUBERT,BE 04A

¹ Uses $B(\Upsilon(4S) \rightarrow B^+B^-) = (51.4 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = (48.6 \pm 0.6)\%$.

² Corresponds to a 90% CL interval $-0.033 < A_{CP} < 0.028$.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ Corresponds to a 90% CL allowed region, $-0.074 < A_{CP} < 0.049$.

⁵ A 90% CL range is $-0.170 < A_{CP} < 0.082$.

$A_{CP}(B \rightarrow s\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.015 ±0.011 OUR AVERAGE

0.0144±0.0128±0.0011	¹ WATANUKI	19	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.017 ±0.019 ±0.010	² LEES	14K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

−0.011 ±0.030 ±0.014	³ AUBERT	08BJ	BABR	Repl. by LEES 14K
0.025 ±0.050 ±0.015	⁴ AUBERT,B	04E	BABR	Repl. by AUBERT 08BJ
0.002 ±0.050 ±0.030	⁵ NISHIDA	04	BELL	Repl. by WATANUKI 19

¹ Using a sum-of-exclusive technique with $m_{X_s} < 2.8 \text{ GeV}/c^2$.

² Measured with 16 exclusively reconstructed $B \rightarrow X_s\gamma$ decays with $0.6 < m_{X_s} < 2.0 \text{ GeV}/c^2$ (ten charged and six neutral self-tagging B modes).

³ Uses a sum of exclusively reconstructed $B \rightarrow X_s$ decay modes, with X_s mass between 0.6 and 2.8 GeV/c^2 .

⁴ Corresponds to $-0.06 < A_{CP} < 0.11$ at 90% CL.

⁵ This measurement is performed inclusively for recoil mass X_s less than 2.1 GeV , which corresponds to $-0.093 < A_{CP} < 0.096$ at 90% CL.

$A_{CP}(B \rightarrow (s+d)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.010 ± 0.031 OUR AVERAGE			
$0.022 \pm 0.039 \pm 0.009$	¹ PESANTEZ	15	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$0.057 \pm 0.060 \pm 0.018$	LEES	12V	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$-0.10 \pm 0.18 \pm 0.05$	² AUBERT	08O	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$-0.110 \pm 0.115 \pm 0.017$	AUBERT,BE	06B	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$-0.079 \pm 0.108 \pm 0.022$	³ COAN	01	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. Uses an opposite side lepton tag. Requires center-of-mass frame $E_\gamma > 2.1$ GeV.			
² Uses a fully reconstructed B meson as a tag on the recoil side. Requires $E_\gamma > 2.2$ GeV.			
³ Corresponds to $-0.27 < A_{CP} < 0.10$ at 90% CL.			

 $A_{CP}(B \rightarrow X_s \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.04 \pm 0.11 \pm 0.01$	¹ LEES	14D	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.22 \pm 0.26 \pm 0.02$	² AUBERT,B	04I	BABR Repl. by LEES 14D
¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, $K_S^0\pi^+$, and $K_S^0\pi^+\pi^0$.			
² The final state flavor is determined by the kaon and pion charges where modes with $X_S = K_S^0$, $K_S^0\pi^0$ or $K_S^0\pi^+\pi^-$ are not used.			

 $A_{CP}(B \rightarrow X_s \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.06 \pm 0.22 \pm 0.01$	¹ LEES	14D	BABR $e^+e^- \rightarrow \Upsilon(4S)$
¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, $K_S^0\pi^+$, and $K_S^0\pi^+\pi^0$.			

 $A_{CP}(B \rightarrow X_s \ell^+ \ell^-) (10.1 < q^2 < 12.9 \text{ or } q^2 > 14.2 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.19^{+0.18}_{-0.17} \pm 0.01$	¹ LEES	14D	BABR $e^+e^- \rightarrow \Upsilon(4S)$
¹ Measured from sum of exclusive modes through K^+ , $K^+\pi^0$, $K^+\pi^-$, $K^+\pi^-\pi^0$, $K^+\pi^-\pi^+$, $K_S^0\pi^+$, and $K_S^0\pi^+ (pi^-)^0$.			

 $A_{CP}(B \rightarrow K^* e^+ e^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.18 \pm 0.15 \pm 0.01$	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

 $A_{CP}(B \rightarrow K^* \mu^+ \mu^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.03 \pm 0.13 \pm 0.02$	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

 $A_{CP}(B \rightarrow K^* \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.04 ± 0.07 OUR AVERAGE			
$0.03 \pm 0.13 \pm 0.01$	¹ LEES	12S	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$+0.01^{+0.16}_{-0.15} \pm 0.01$	AUBERT	09T	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.10 \pm 0.10 \pm 0.01$	WEI	09A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Measured in the union of $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $q^2 > 10.11 \text{ GeV}^2/c^4$.
LEES 12S reports also individual measurements $A_{CP}(B \rightarrow K^* \ell^+ \ell^-) = -0.13^{+0.18}_{-0.19} \pm 0.01$ for $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$ and $A_{CP}(B \rightarrow K^* \ell^+ \ell^-) = 0.16^{+0.18}_{-0.19} \pm 0.01$ for $q^2 > 10.11 \text{ GeV}^2/c^4$.

$A_{CP}(B \rightarrow \eta \text{ anything})$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.13 \pm 0.04^{+0.02}_{-0.03}$	¹ NISHIMURA	10	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $B \rightarrow \eta X_s$ with $0.4 < m_{X_s} < 2.6 \text{ GeV}/c^2$.

$\Delta A_{CP}(X_s \gamma) = A_{CP}(B^\pm \rightarrow X_s \gamma) - A_{CP}(B^0 \rightarrow X_s \gamma)$

This is the isospin difference of the CP asymmetries.

VALUE	DOCUMENT ID	TECN	COMMENT
0.041 ± 0.023 OUR AVERAGE			
$0.0369 \pm 0.0265 \pm 0.0076$	¹ WATANUKI	19	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$0.050 \pm 0.039 \pm 0.015$	² LEES	14K	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Using a sum-of-exclusive technique with $m_{X_s} < 2.8 \text{ GeV}/c^2$.

² Measured with 16 exclusively reconstructed $B \rightarrow X_s \gamma$ decays with $0.6 < m_{X_s} < 2.0 \text{ GeV}/c^2$ (ten charged and six neutral self-tagging B modes).

$\overline{A}_{CP}(B \rightarrow X_s \gamma) = (A_{CP}(B^+ \rightarrow X_s \gamma) + A_{CP}(B^0 \rightarrow X_s \gamma))/2$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0091 \pm 0.0121 \pm 0.0013$	¹ WATANUKI	19	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Using a sum-of-exclusive technique with $m_{X_s} < 2.8 \text{ GeV}/c^2$.

$\Delta A_{CP}(B \rightarrow K^* \gamma) = A_{CP}(B^+ \rightarrow K^{*+} \gamma) - A_{CP}(B^0 \rightarrow K^{*0} \gamma)$

This is the isospin difference of the CP asymmetries.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.024 \pm 0.028 \pm 0.005$	¹ HORIGUCHI	17	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.4 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \overline{B}^0) = (48.6 \pm 0.6)\%$.

$\overline{A}_{CP}(B \rightarrow K^* \gamma) = (A_{CP}(B^+ \rightarrow K^{*+} \gamma) + A_{CP}(B^0 \rightarrow K^{*0} \gamma))/2$

This is the average CP asymmetry.

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.001 \pm 0.014 \pm 0.003$	¹ HORIGUCHI	17	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.4 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \overline{B}^0) = (48.6 \pm 0.6)\%$.

POLARIZATION IN B DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel ($\parallel$) or perpendicular ($\perp$) to each other with the parameters Γ_L/Γ , $\Gamma_\perp/\Gamma$, and the relative phases $\phi_\parallel$ and $\phi_\perp$. See the definitions in the note on “Polarization in B Decays” review in the B^0 Particle Listings.

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (q^2 > 0.1 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.63^{+0.18}_{-0.19} \pm 0.05$	¹ AUBERT,B	06J	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Results with different q^2 cuts are also reported.

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} < 2.5 \text{ GeV}/c^2)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.35 \pm 0.16 \pm 0.04$	AUBERT	09N	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} > 3.2 \text{ GeV}/c^2)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.71^{+0.20}_{-0.22} \pm 0.04$	AUBERT	09N	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (0.10 < q^2 < 0.98 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.263^{+0.045}_{-0.044} \pm 0.017$	AAIJ	16B	LHCB pp at 7, 8 TeV

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (1.1 < q^2 < 2.5 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.660^{+0.083}_{-0.077} \pm 0.022$	AAIJ	16B	LHCB pp at 7, 8 TeV

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.34^{+0.08}_{-0.07}$	OUR AVERAGE		

$0.37^{+0.10}_{-0.09} \pm 0.04$	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.30 \pm 0.16 \pm 0.02$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.29^{+0.21}_{-0.18} \pm 0.02$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.60^{+0.00}_{-0.28} \pm 0.19$	¹ CHATRCHYAN	13BL	CMS pp at 7 TeV
$0.00^{+0.13}_{-0.00} \pm 0.02$	AAIJ	12U	LHCB Repl. by AAIJ 13Y
$0.53^{+0.32}_{-0.34} \pm 0.07$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

¹ CHATRCHYAN 13BL uses, for this bin, $1.0 < q^2 < 2.0 \text{ GeV}^2/c^4$.

$F_L(B \rightarrow K^* \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.05 OUR AVERAGE			
$0.876^{+0.109}_{-0.097} \pm 0.017$	¹ AAIJ	16B LHCb	pp at 7, 8 TeV
$0.80 \pm 0.08 \pm 0.06$	KHACHATRY...16D	CMS	pp at 8 TeV
$0.74^{+0.10}_{-0.09} \pm 0.02_{-0.03}$	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.65 \pm 0.17 \pm 0.03$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.37^{+0.25}_{-0.24} \pm 0.10$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.71 \pm 0.24 \pm 0.05$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.77 \pm 0.15 \pm 0.03$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.40^{+0.32}_{-0.33} \pm 0.08$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

¹ Measured in $2.5 < q^2 < 4.0 \text{ GeV}^2/c^4$. **$F_L(B \rightarrow K^* \ell^+ \ell^-) (4.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$**

VALUE	DOCUMENT ID	TECN	COMMENT
$0.611^{+0.052}_{-0.053} \pm 0.017$	AAIJ	16B LHCb	pp at 7, 8 TeV

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (6.0 < q^2 < 8.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.579 \pm 0.046 \pm 0.015$	AAIJ	16B LHCb	pp at 7, 8 TeV

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (4.3 < q^2 < 8.6 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 ± 0.06 OUR AVERAGE			
$0.57 \pm 0.07 \pm 0.03$	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.81^{+0.13}_{-0.12} \pm 0.05$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.68^{+0.15}_{-0.17} \pm 0.09$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.64^{+0.23}_{-0.24} \pm 0.07$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.60^{+0.06}_{-0.07} \pm 0.01$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.82^{+0.19}_{-0.23} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $F_L(B \rightarrow K^* \ell^+ \ell^-) (10.09 < q^2 < 12.86 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.448 ± 0.033 OUR AVERAGE			
$0.493^{+0.049}_{-0.047} \pm 0.013$	¹ AAIJ	16B LHCb	pp at 7, 8 TeV
$0.39 \pm 0.05 \pm 0.04$	KHACHATRY...16D	CMS	pp at 8 TeV
$0.48^{+0.08}_{-0.09} \pm 0.03$	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.45^{+0.10}_{-0.11} \pm 0.04$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.47 \pm 0.14 \pm 0.03$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.17^{+0.17}_{-0.15} \pm 0.03$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.41 \pm 0.11 \pm 0.03$	AAIJ	12U	LHCB	Repl. by AAIJ 13Y
$0.31^{+0.19}_{-0.18} \pm 0.02$	AALTONEN	11L	CDF	Repl. by AALTONEN 12I

¹ Measured in $11.0 < q^2 < 12.5 \text{ GeV}^2/c^4$.

$F_L(B \rightarrow K^* \ell^+ \ell^-) (15.0 < q^2 < 17.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.349 \pm 0.039 \pm 0.009$	AAIJ	16B	LHCB pp at 7, 8 TeV

$F_L(B \rightarrow K^* \ell^+ \ell^-) (17.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$


VALUE	DOCUMENT ID	TECN	COMMENT
$0.354^{+0.049}_{-0.048} \pm 0.025$	AAIJ	16B	LHCB pp at 7, 8 TeV

$F_L(B \rightarrow K^* \ell^+ \ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.40 ± 0.06 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
$0.48^{+0.05}_{-0.06} \pm 0.04$	KHACHATRY...16D	CMS	pp at 8 TeV
$0.33^{+0.08+0.02}_{-0.07-0.03}$	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.53 \pm 0.12 \pm 0.03$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.29^{+0.14}_{-0.13} \pm 0.05$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$-0.15^{+0.27}_{-0.23} \pm 0.07$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37 \pm 0.09 \pm 0.05$	AAIJ	12U	LHCB	Repl. by AAIJ 13Y
$0.55^{+0.17}_{-0.18} \pm 0.02$	AALTONEN	11L	CDF	Repl. by AALTONEN 12I




$F_L(B \rightarrow K^* \ell^+ \ell^-) (16.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.350 ± 0.019 OUR AVERAGE			
0.345 ± 0.020 ± 0.007	¹ AAIJ	20Y LHCb	pp at 7, 8, 13 TeV
0.38 $^{+0.05}_{-0.06}$ ± 0.04	KHACHATRYAN 16D	CMS	pp at 8 TeV
0.38 $^{+0.09}_{-0.07}$ ± 0.03	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.44 ± 0.07 ± 0.03	CHATRCHYAN 13BL	CMS	pp at 7 TeV
0.20 $^{+0.19}_{-0.17}$ ± 0.05	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.12 $^{+0.15}_{-0.13}$ ± 0.02	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.344 $^{+0.028}_{-0.030}$ ± 0.008	¹ AAIJ	16B LHCb	Repl. by AAIJ 20Y
0.26 $^{+0.10}_{-0.08}$ ± 0.03	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.09 $^{+0.18}_{-0.14}$ ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 12I

¹ Measured in $15.0 < q^2 < 19.0 \text{ GeV}^2/c^4$. **$F_L(B \rightarrow K^* \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.67 ± 0.07 OUR AVERAGE Error includes scale factor of 2.8. See the ideogram below.			
0.700 ± 0.025 ± 0.013	¹ AAIJ	20Y LHCb	pp at 7, 8, 13 TeV
0.72 ± 0.06	KHACHATRYAN 16D	CMS	pp at 7, 8 TeV
0.24 $^{+0.09}_{-0.08}$ ± 0.02	² LEES	16C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.65 $^{+0.08}_{-0.07}$ ± 0.03	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.69 $^{+0.19}_{-0.21}$ ± 0.08	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
0.67 ± 0.23 ± 0.05	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.690 $^{+0.035}_{-0.036}$ ± 0.017	¹ AAIJ	16B LHCb	Repl. by AAIJ 20Y
0.68 ± 0.10 ± 0.02	CHATRCHYAN 13BL	CMS	Repl. by KHACHATRYAN 16D
0.55 ± 0.10 ± 0.03	AAIJ	12U LHCb	Repl. by AAIJ 13Y
0.50 $^{+0.27}_{-0.30}$ ± 0.03	AALTONEN	11L CDF	Repl. by AALTONEN 12I

¹ Measured in $1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$.² Measured by combining B^0 and B^+ with e and μ as leptons. Results are also provided separately for B^0 and B^+ .



$F_L(B \rightarrow K^* \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.33^{+0.14}_{-0.13} \pm 0.03$	AALTONEN 12I	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.47^{+0.23}_{-0.24} \pm 0.03$	AALTONEN 11L	CDF	Repl. by AALTONEN 12I

$A_T^{(2)}$ in $B \rightarrow K^* e^+ e^-$ (at low q^2)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.52 \pm 0.53 \pm 0.11$	1,2 FERLEWICZ 24	BELL	$e^+ e^- \rightarrow \gamma(4S)$
¹ Events are reconstructed in $B^0 \rightarrow K^{*0}(\rightarrow K^+ \pi^-) e^+ e^-$ and $B^+ \rightarrow K^{*+}(\rightarrow K_S^0 \pi^+) e^+ e^-$ decay modes. The effective dielectron invariant mass is required to be in the range $0.0008 < q^2 < 1.12 \text{ GeV}^2/c^4$.			
² A_T^{Im} and $A_T^{(2)}$ measure the ratio of the right- and left-handed Wilson coefficients, C_7'/C_7 , which constrains non-SM contributions at small q^2 .			

A_T^{Im} in $B \rightarrow K^* e^+ e^-$ (at low q^2)

VALUE	DOCUMENT ID	TECN	COMMENT
$-1.27 \pm 0.52 \pm 0.12$	1,2 FERLEWICZ 24	BELL	$e^+ e^- \rightarrow \gamma(4S)$

¹ Events are reconstructed in $B^0 \rightarrow K^{*0}(\rightarrow K^+\pi^-)e^+e^-$ and $B^+ \rightarrow K^{*+}(\rightarrow K_S^0\pi^+)e^+e^-$ decay modes. The effective dielectron invariant mass is required to be in the range $0.0008 < q^2 < 1.12 \text{ GeV}^2/c^4$.

² A_T^{Im} and $A_T^{(2)}$ measure the ratio of the right- and left-handed Wilson coefficients, C_7/C_7 , which constrains non-SM contributions at small q^2 .

$P_\tau(B \rightarrow D^*\tau^+\nu_\tau)$

Measures difference in decay widths with positive and negative τ^+ helicities normalized to the sum of those decay widths.

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.38 \pm 0.51^{+0.21}_{-0.16}$	¹ HIROSE	17	BELL $e^+e^- \rightarrow \gamma(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

Γ_L/Γ in $B \rightarrow \bar{D}^*\ell^+\nu_\ell$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.502 \pm 0.012 \pm 0.004$	¹ PRIM	23	BELL $e^+e^- \rightarrow \gamma(4S)$

¹ This is the $B^+ \rightarrow \bar{D}^{*0}\ell^+\nu_\ell$ and $B^0 \rightarrow D^{*-}\ell^+\nu_\ell$ average.

Γ_L/Γ in $B \rightarrow \bar{D}^*e^+\nu_e$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.485 \pm 0.018 \pm 0.005$	¹ PRIM	23	BELL $e^+e^- \rightarrow \gamma(4S)$

¹ This is the $B^+ \rightarrow \bar{D}^{*0}e^+\nu_e$ and $B^0 \rightarrow D^{*-}e^+\nu_e$ average.

Γ_L/Γ in $B \rightarrow \bar{D}^*\mu^+\nu_\mu$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.515 \pm 0.017 \pm 0.005$	¹ PRIM	23	BELL $e^+e^- \rightarrow \gamma(4S)$

¹ This is the $B^+ \rightarrow \bar{D}^{*0}\mu^+\nu_\mu$ and $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ average.

$\Delta(\Gamma_L/\Gamma)$ in $B \rightarrow \bar{D}^*(lepton)^+\nu_\ell$

$\Delta(\Gamma_L/\Gamma) = (\Gamma_L/\Gamma)^\mu - (\Gamma_L/\Gamma)^e$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.030 \pm 0.025 \pm 0.007$	¹ PRIM	23	BELL $e^+e^- \rightarrow \gamma(4S)$

¹ This is the B^+ and B^0 average.

PARTIAL BRANCHING FRACTIONS IN $B \rightarrow K^{(*)}\ell^+\ell^-$

$B(B \rightarrow K^*\ell^+\ell^-) (q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.68 ± 0.23 OUR AVERAGE			
$1.89^{+0.52}_{-0.46} \pm 0.06$	¹ LEES	12S	BABR $e^+e^- \rightarrow \gamma(4S)$
$1.73 \pm 0.33 \pm 0.10$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$1.46^{+0.40}_{-0.35} \pm 0.11$	WEI	09A	BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.98 \pm 0.40 \pm 0.09$ AALTONEN 11L CDF Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $0.1 < q^2 < 2.0 \text{ GeV}^2/c^2$.

$B(B \rightarrow K^* \ell^+ \ell^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

0.87 $\pm$ 0.17 OUR AVERAGE

$0.95^{+0.35}_{-0.30} \pm 0.04$	LEES	12S	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$0.82 \pm 0.26 \pm 0.06$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$0.86^{+0.31}_{-0.27} \pm 0.07$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.00 \pm 0.38 \pm 0.09$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

 $B(B \rightarrow K^* \ell^+ \ell^-)$ ($4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.67 $\pm$ 0.29 OUR AVERAGE

$1.82^{+0.56}_{-0.52} \pm 0.09$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$1.72 \pm 0.41 \pm 0.14$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$1.37^{+0.47}_{-0.42} \pm 0.39$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.69 \pm 0.57 \pm 0.15$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

¹ The value reported here from LEES 12S refers to $4.3 < q^2 < 8.12 \text{ GeV}^2/c^2$.

 $B(B \rightarrow K^* \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.93 $\pm$ 0.25 OUR AVERAGE

$1.86^{+0.52}_{-0.48} \pm 0.10$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$1.77 \pm 0.34 \pm 0.11$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$2.24^{+0.44}_{-0.40} \pm 0.19$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.97 \pm 0.47 \pm 0.17$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

¹ The value reported here from LEES 12S refers to $10.11 < q^2 < 12.89 \text{ GeV}^2/c^2$.

 $B(B \rightarrow K^* \ell^+ \ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.21 $\pm$ 0.17 OUR AVERAGE

$1.46^{+0.41}_{-0.36} \pm 0.06$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$1.21 \pm 0.24 \pm 0.07$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$1.05^{+0.29}_{-0.26} \pm 0.08$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$


• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.51 \pm 0.36 \pm 0.13$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

¹ The value reported here from LEES 12S refers to $14.21 < q^2 < 16.0 \text{ GeV}^2/c^2$.

$B(B \rightarrow K^* \ell^+ \ell^-)$ ($16.0 < q^2 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.3. See the ideogram below.		
$1.02^{+0.47}_{-0.42} \pm 0.06$	LEES	12S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.88 \pm 0.22 \pm 0.05$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$2.04^{+0.27}_{-0.24} \pm 0.16$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.35 \pm 0.37 \pm 0.12$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI



$B(B \rightarrow K^* \ell^+ \ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.64 ± 0.26 OUR AVERAGE			
$2.05^{+0.53}_{-0.48} \pm 0.07$	LEES	12S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.48 \pm 0.39 \pm 0.12$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$1.49^{+0.45}_{-0.40} \pm 0.12$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.60 \pm 0.54 \pm 0.14$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B \rightarrow K^* \ell^+ \ell^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$2.53 \pm 0.43 \pm 0.15$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.98 \pm 0.55 \pm 0.18$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B^+ \rightarrow K^* \mu^+ \mu^-) / B(B^+ \rightarrow K^* e^+ e^-)$ ($0.045 < q^2 < 1.1 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.52^{+0.36}_{-0.26} \pm 0.05$	WEHLE	21	BELL $e^+ e^- \rightarrow \gamma(4S)$

$B(B^+ \rightarrow K^* \mu^+ \mu^-) / B(B^+ \rightarrow K^* e^+ e^-)$ ($1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.96^{+0.45}_{-0.29} \pm 0.11$	WEHLE	21	BELL $e^+ e^- \rightarrow \gamma(4S)$

$B(B^+ \rightarrow K^* \mu^+ \mu^-) / B(B^+ \rightarrow K^* e^+ e^-)$ ($15.0 < q^2 < 19.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$1.18^{+0.52}_{-0.32} \pm 0.10$	WEHLE	21	BELL $e^+ e^- \rightarrow \gamma(4S)$


$B(B \rightarrow K \ell^+ \ell^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.16 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
$0.71^{+0.20}_{-0.18} \pm 0.02$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
$0.33 \pm 0.10 \pm 0.02$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
$0.81^{+0.18}_{-0.16} \pm 0.05$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.38 \pm 0.16 \pm 0.03$	AALTONEN	11L	CDF	Repl. by AALTONEN 11AI
--------------------------	----------	-----	-----	------------------------

¹ The value reported here from LEES 12S refers to $0.1 < q^2 < 2.0 \text{ GeV}^2/c^2$.



$B(B \rightarrow K \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

 $0.57^{+0.10}_{-0.09}$ OUR AVERAGE Error includes scale factor of 1.2.

$0.49^{+0.15}_{-0.13} \pm 0.01$	LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	------	-----	---------------------------------------

$0.77 \pm 0.14 \pm 0.05$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
--------------------------	----------	------	----------------------------

$0.46^{+0.14}_{-0.12} \pm 0.03$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.58 \pm 0.19 \pm 0.04$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

 $B(B \rightarrow K \ell^+ \ell^-) (4.3 < q^2 < 8.68 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

 1.00 ± 0.11 OUR AVERAGE

$0.94^{+0.20}_{-0.19} \pm 0.02$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	-------------------	-----	---------------------------------------

$1.05 \pm 0.17 \pm 0.07$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
--------------------------	----------	------	----------------------------

$1.00^{+0.19}_{-0.18} \pm 0.06$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.93 \pm 0.25 \pm 0.06$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

¹ The value reported here from LEES 12S refers to $4.3 < q^2 < 8.12 \text{ GeV}^2/c^2$. **$B(B \rightarrow K \ell^+ \ell^-) (10.09 < q^2 < 12.86 \text{ GeV}^2/c^4)$**

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

 0.57 ± 0.11 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

$0.90^{+0.20}_{-0.19} \pm 0.04$	¹ LEES	12S	BABR $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	-------------------	-----	---------------------------------------


$0.48 \pm 0.10 \pm 0.03$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV
--------------------------	----------	------	----------------------------

$0.55^{+0.16}_{-0.14} \pm 0.03$	WEI	09A	BELL $e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.72 \pm 0.17 \pm 0.05$	AALTONEN	11L	CDF Repl. by AALTONEN 11AI
--------------------------	----------	-----	----------------------------

¹ The value reported here from LEES 12S refers to $10.11 < q^2 < 12.89 \text{ GeV}^2/c^2$.




$B(B \rightarrow K \ell^+ \ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.49 ± 0.07 OUR AVERAGE			
$0.49^{+0.15}_{-0.14} \pm 0.02$	¹ LEES	12S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.52 \pm 0.09 \pm 0.03$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$0.38^{+0.19}_{-0.12} \pm 0.02$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.38 \pm 0.12 \pm 0.03$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

¹ The value reported here from LEES 12S refers to $14.21 < q^2 < 16.0 \text{ GeV}^2/c^2$.

$B(B \rightarrow K \ell^+ \ell^-) (16.0 < q^2 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.16 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.		
$0.67^{+0.23}_{-0.21} \pm 0.05$	LEES	12S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.38 \pm 0.09 \pm 0.02$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$0.98^{+0.20}_{-0.18} \pm 0.06$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.35 \pm 0.13 \pm 0.02$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI



$B(B \rightarrow K \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.33 ± 0.13 OUR AVERAGE			
$1.36^{+0.27}_{-0.24} \pm 0.03$	LEES	12S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.29 \pm 0.18 \pm 0.08$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$1.36^{+0.23}_{-0.21} \pm 0.08$	WEI	09A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.01 \pm 0.26 \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B \rightarrow K \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$1.07 \pm 0.17 \pm 0.07$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.96 \pm 0.25 \pm 0.06$	AALTONEN	11L CDF	Repl. by AALTONEN 11AI

$B(B \rightarrow X_S \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$1.60^{+0.41+0.25}_{-0.39-0.22}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \gamma(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi^-$ corrected for unobserved modes.

$B(B \rightarrow X_s e^+ e^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$1.93^{+0.47+0.28}_{-0.45-0.24}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi + \pi^-$ corrected for unobserved modes.

 $B(B \rightarrow X_s \mu^+ \mu^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.66^{+0.82+0.31}_{-0.76-0.25}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi + \pi^-$ corrected for unobserved modes.

 $B(B \rightarrow X_s \ell^+ \ell^-) (14.2 < q^2 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.57^{+0.16+0.03}_{-0.15-0.02}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi + \pi^-$ corrected for unobserved modes.

 $B(B \rightarrow X_s e^+ e^-) (14.2 < q^2 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.56^{+0.19+0.03}_{-0.18-0.03}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi + \pi^-$ corrected for unobserved modes.

 $B(B \rightarrow X_s \mu^+ \mu^-) (14.2 < q^2 \text{ GeV}^2/c^4)$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.60^{+0.31+0.05}_{-0.29-0.04}$	¹ LEES	14D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured from sum of exclusive modes through K^+ , $K^+ \pi^0$, $K^+ \pi^-$, $K^+ \pi^- \pi^0$, $K^+ \pi^- \pi^+$, K_S^0 , $K_S^0 \pi^0$, $K_S^0 \pi^+$, $K_S^0 \pi^+ \pi^0$, and $K_S^0 \pi + \pi^-$ corrected for unobserved modes.

LEPTON (HADRON) FORWARD-BACKWARD ASYMMETRY IN $B \rightarrow K^{(*)} \ell^+ \ell^-$ ($B \rightarrow K/\pi h^+ h^-$) DECAY

The forward-backward angular asymmetry of the lepton pair in $B \rightarrow K^{(*)} \ell^+ \ell^-$ ($B \rightarrow K/\pi h^+ h^-$) decay is defined as

$$A_{FB}(s) = \frac{N(\cos\theta > 0) - N(\cos\theta < 0)}{N(\cos\theta > 0) + N(\cos\theta < 0)},$$

where $s = q^2/m_B^2$, and θ is the angle of the ℓ^- (h^-) with respect to the flight direction of the B meson, measured in the dilepton (dihadron)


rest frame. In addition, the fraction of longitudinal polarization F_L of the K^* and F_S , the relative contribution from scalar and pseudoscalar penguin amplitudes in $B \rightarrow K \ell^+ \ell^-$, can be measured from the angular distribution of its decay products.

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (q^2 > 0.1 \text{ GeV}^2/c^4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$0.50 \pm 0.15 \pm 0.02$		¹ ISHIKAWA 06	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.55	95	² AUBERT,B 06J	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
¹ Using an unbinned max. likelihood fits to the M_{bc} distribution in five q^2 bins for $\cos \theta > 0$ and $\cos \theta < 0$.				
² Results with different q^2 cuts are also reported.				

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.14 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
$-0.02 \pm 0.12 \pm 0.01$	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$-0.35^{+0.26}_{-0.23} \pm 0.10$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.47^{+0.26}_{-0.32} \pm 0.03$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.29^{+0.37}_{-0.00} \pm 0.18$	¹ CHATRCHYAN 13BL	CMS	pp at 7 TeV
$-0.15 \pm 0.20 \pm 0.06$	AAIJ	12U LHCb	Repl. by AAIJ 13Y
$0.13^{+1.65}_{-0.75} \pm 0.25$	AALTONEN	11L CDF	Repl. by AALTONEN 12I



¹CHATRCHYAN 13BL uses, for this bin, $1.0 < q^2 < 2.0 \text{ GeV}^2/c^4$.

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} < 2.5 \text{ GeV}/c^2)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.24^{+0.18}_{-0.23} \pm 0.05$	AUBERT	09N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (m_{\ell\ell} > 3.2 \text{ GeV}/c^2)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.76^{+0.52}_{-0.32} \pm 0.07$	AUBERT	09N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (0.10 < q^2 < 0.98 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.003^{+0.058}_{-0.057} \pm 0.009$	AAIJ	16B LHCB	pp at 7, 8 TeV

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (1.1 < q^2 < 2.5 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.191^{+0.068}_{-0.080} \pm 0.012$	AAIJ	16B LHCB	pp at 7, 8 TeV

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.14 ± 0.05 OUR AVERAGE			
$-0.118^{+0.082}_{-0.090} \pm 0.007$	¹ AAIJ	16B LHCB	pp at 7, 8 TeV
$-0.12^{+0.15}_{-0.17} \pm 0.05$	KHACHATRY...16D	CMS	pp at 8 TeV
$-0.20 \pm 0.08 \pm 0.01$	AAIJ	13Y LHCB	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$-0.07 \pm 0.20 \pm 0.02$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.29^{+0.32}_{-0.35} \pm 0.15$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$0.11^{+0.31}_{-0.36} \pm 0.07$	WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.05^{+0.16}_{-0.20} \pm 0.04$	AAIJ	12U LHCB	Repl. by AAIJ 13Y
$0.19^{+0.40}_{-0.41} \pm 0.14$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

¹ Measured in $2.5 < q^2 < 4.0 \text{ GeV}^2/c^4$.

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.08^{+0.21}_{-0.20} \pm 0.05$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.21^{+0.31}_{-0.33} \pm 0.05$	AALTONEN	11L CDF	Repl. by AALTONEN 12I
---------------------------------	----------	---------	-----------------------

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-) (4.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.025^{+0.051}_{-0.052} \pm 0.004$	AAIJ	16B LHCB	pp at 7, 8 TeV

$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($6.0 < q^2 < 8.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.152^{+0.041}_{-0.040} \pm 0.008$	AAIJ	16B	LHCB pp at 7, 8 TeV

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.078 ± 0.022 OUR AVERAGE	Error includes scale factor of 1.1.		
$-0.073 \pm 0.021 \pm 0.002$	¹ AAIIJ	20Y	LHCB pp at 7, 8, 13 TeV
-0.12 ± 0.08	KHACHATRY...16D	CMS	pp at 7, 8 TeV
$0.21^{+0.10}_{-0.15} \pm 0.07$	² LEES	16C	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$-0.17 \pm 0.06 \pm 0.01$	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.29^{+0.20}_{-0.23} \pm 0.07$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.26^{+0.27}_{-0.30} \pm 0.07$	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.075^{+0.032}_{-0.034} \pm 0.007$	¹ AAIIJ	16B	LHCB Repl. by AAIIJ 20Y
0.55 ± 0.43	³ SATO	16	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$-0.07 \pm 0.12 \pm 0.01$	CHATRCHYAN 13BL	CMS	Repl. by KHACHATRYAN 16D
$-0.06^{+0.13}_{-0.14} \pm 0.07$	AAIJ	12U	LHCB Repl. by AAIIJ 13Y
$0.43^{+0.36}_{-0.37} \pm 0.06$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

¹ Measured in $1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$.

² Measured by combining B^0 and B^+ with e and μ as leptons. Results are also provided separately for B^0 and B^+ .

³ Uses $K^* \rightarrow K^- \pi^+$, $K^- \pi^0$, $K_S^0 \pi^-$ in the range $M(K\pi) < 1.1 \text{ GeV}/c^2$. Uncertainty is statistical only.

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($4.3 < q^2 < 8.6 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.13^{+0.06}_{-0.05}$ OUR AVERAGE	Error includes scale factor of 1.1.		
$0.16^{+0.06}_{-0.05} \pm 0.01$	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$-0.01 \pm 0.11 \pm 0.03$	CHATRCHYAN 13BL	CMS	pp at 7 TeV
$0.01 \pm 0.20 \pm 0.09$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.45^{+0.15}_{-0.21} \pm 0.15$	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.27^{+0.06}_{-0.08} \pm 0.02$	AAIJ	12U	LHCB Repl. by AAIIJ 13Y
$-0.06^{+0.30}_{-0.28} \pm 0.05$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.288 ± 0.034 OUR AVERAGE	Error includes scale factor of 1.2.		
$0.318^{+0.044}_{-0.040} \pm 0.009$	¹ AAIIJ	16B	LHCB pp at 7, 8 TeV
$0.16 \pm 0.06 \pm 0.01$	KHACHATRY...16D	CMS	pp at 8 TeV


0.28 ^{+0.07} _{-0.06} ± 0.02	AAIJ	13Y	LHCB	pp at 7 TeV, $K^{*0}\mu^+\mu^-$
0.40 ± 0.08 ± 0.05	CHATRCHYAN 13BL	CMS		pp at 7 TeV
0.38 ^{+0.16} _{-0.19} ± 0.09	AALTONEN	12I	CDF	$p\bar{p}$ at 1.96 TeV
0.43 ^{+0.18} _{-0.20} ± 0.03	WEI	09A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.27 ^{+0.11} _{-0.13} ± 0.02	AAIJ	12U	LHCB	Repl. by AAIJ 13Y
0.66 ^{+0.23} _{-0.20} ± 0.07	AALTONEN	11L	CDF	Repl. by AALTONEN 12I

¹ Measured in $11.0 < q^2 < 12.5 \text{ GeV}^2/c^4$.

$A_{FB}(B \rightarrow K^*\ell^+\ell^-)$ ($14.18 < q^2 < 16.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.43^{+0.05}_{-0.06} OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
0.39 ^{+0.04} _{-0.06} ± 0.01	KHACHATRY...16D	CMS	pp at 8 TeV
0.51 ^{+0.07} _{-0.05} ± 0.02	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0}\mu^+\mu^-$
0.29 ± 0.09 ± 0.05	CHATRCHYAN 13BL	CMS	pp at 7 TeV
0.44 ^{+0.18} _{-0.21} ± 0.10	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
0.70 ^{+0.16} _{-0.22} ± 0.10	WEI	09A	BELL $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.47 ^{+0.06} _{-0.08} ± 0.03	AAIJ	12U	LHCB Repl. by AAIJ 13Y
0.42 ± 0.16 ± 0.09	AALTONEN	11L	CDF Repl. by AALTONEN 12I

WEIGHTED AVERAGE
0.43+0.05-0.06 (Error scaled by 1.6)



$A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($15.0 < q^2 < 17.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.411^{+0.41}_{-0.037} \pm 0.008$	AAIJ	16B	LHCB pp at 7, 8 TeV

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($17.0 < q^2 < 19.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.305^{+0.049}_{-0.048} \pm 0.013$	AAIJ	16B	LHCB pp at 7, 8 TeV

 $A_{FB}(B \rightarrow K^* \ell^+ \ell^-)$ ($16.0 < q^2 < 19.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.362 ± 0.019 OUR AVERAGE			
$0.353 \pm 0.020 \pm 0.010$	¹ AAIJ	20Y	LHCB pp at 7, 8, 13 TeV
$0.35 \pm 0.07 \pm 0.01$	KHACHATRY...	16D	CMS pp at 8 TeV
$0.30 \pm 0.08 \pm^{+0.01}_{-0.02}$	AAIJ	13Y	LHCB pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
$0.41 \pm 0.05 \pm 0.03$	CHATRCHYAN	13BL	CMS pp at 7 TeV
$0.65 \pm^{+0.17}_{-0.18} \pm 0.16$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.66 \pm^{+0.11}_{-0.16} \pm 0.04$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.355 \pm 0.027 \pm 0.009$	¹ AAIJ	16B	LHCB Repl. by AAIJ 20Y
$0.16 \pm^{+0.11}_{-0.13} \pm 0.06$	AAIJ	12U	LHCB Repl. by AAIJ 13Y
$0.70 \pm^{+0.16}_{-0.25} \pm 0.10$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

¹ Measured in $15.0 < q^2 < 19.0 \text{ GeV}^2/c^4$. **$A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($q^2 > 0.1 \text{ GeV}^2/c^4$)**

VALUE	DOCUMENT ID	TECN	COMMENT
0.11 ± 0.12 OUR AVERAGE			
$0.15 \pm^{+0.21}_{-0.23} \pm 0.08$	¹ AUBERT,B	06J	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$0.10 \pm 0.14 \pm 0.01$	² ISHIKAWA	06	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Results with different q^2 cuts are also reported.² Using an unbinned max. likelihood fits to the M_{bc} distribution in five q^2 bins for $\cos \theta > 0$ and $\cos \theta < 0$. **$A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)**

VALUE	DOCUMENT ID	TECN	COMMENT
$0.00^{+0.06}_{-0.05}$ OUR AVERAGE			
$0.00 \pm^{+0.06}_{-0.05} \pm 0.03$	AAIJ	13H	LHCB pp at 7 TeV
$0.13 \pm^{+0.42}_{-0.43} \pm 0.07$	AALTONEN	12I	CDF $p\bar{p}$ at 1.96 TeV
$0.06 \pm^{+0.32}_{-0.35} \pm 0.02$	WEI	09A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.15 \pm^{+0.46}_{-0.39} \pm 0.08$	AALTONEN	11L	CDF Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.09^{+0.10}_{-0.07}$ OUR AVERAGE	Error includes scale factor of 1.4.		
$0.07^{+0.08}_{-0.05} \pm 0.02$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.32^{+0.15}_{-0.16} \pm 0.05$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$-0.43^{+0.38}_{-0.40} \pm 0.09$	WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.72^{+0.40}_{-0.35} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($0.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.16 \pm 0.04$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.36^{+0.24}_{-0.26} \pm 0.06$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.034^{+0.040}_{-0.029}$ OUR AVERAGE			
$0.02^{+0.05}_{-0.03} \pm 0.02$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.13 \pm 0.09 \pm 0.02$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$-0.04^{+0.13}_{-0.16} \pm 0.05$	WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.00 ± 0.13	¹ SATO	16 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.08^{+0.27}_{-0.22} \pm 0.07$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

¹ Statistical uncertainty only. **$A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($4.3 < q^2 < 8.6 \text{ GeV}^2/c^4$)**

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.04^{+0.04}_{-0.05}$ OUR AVERAGE			
$-0.02^{+0.03}_{-0.05} \pm 0.03$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV
$0.01^{+0.13}_{-0.10} \pm 0.01$	AALTONEN	12I CDF	$p\bar{p}$ at 1.96 TeV
$-0.20^{+0.12}_{-0.14} \pm 0.03$	WEI	09A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.20^{+0.17}_{-0.28} \pm 0.03$	AALTONEN	11L CDF	Repl. by AALTONEN 12I

 $A_{FB}(B \rightarrow K \ell^+ \ell^-)$ ($10.09 < q^2 < 12.86 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
-0.05 ± 0.06 OUR AVERAGE			
$-0.03 \pm 0.07 \pm 0.01$	AAIJ	13H LHCb	$p\bar{p}$ at 7 TeV

$-0.03^{+0.11}_{-0.10} \pm 0.04$ AALTONEN 12I CDF $p\bar{p}$ at 1.96 TeV

$-0.21^{+0.17}_{-0.15} \pm 0.06$ WEI 09A BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.10^{+0.17}_{-0.15} \pm 0.07$ AALTONEN 11L CDF Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K\ell^+\ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$-0.02^{+0.07}_{-0.05}$ OUR AVERAGE

$-0.01^{+0.12}_{-0.06} \pm 0.01$ AAIJ 13H LHCb pp at 7 TeV

$-0.05^{+0.09}_{-0.11} \pm 0.03$ AALTONEN 12I CDF $p\bar{p}$ at 1.96 TeV

$0.04^{+0.32}_{-0.26} \pm 0.05$ WEI 09A BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.03^{+0.49}_{-0.16} \pm 0.04$ AALTONEN 11L CDF Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow K\ell^+\ell^-) (16.0 < q^2 < 18.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$-0.09^{+0.07+0.02}_{-0.09-0.01}$ AAIJ 13H LHCb pp at 7 TeV

$A_{FB}(B \rightarrow K\ell^+\ell^-) (18.0 < q^2 < 22.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.02 \pm 0.11 \pm 0.01$ AAIJ 13H LHCb pp at 7 TeV

$A_{FB}(B \rightarrow K\ell^+\ell^-) (q^2 > 16.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.04^{+0.09}_{-0.07}$ OUR AVERAGE

$0.09^{+0.17}_{-0.13} \pm 0.03$ AALTONEN 12I CDF $p\bar{p}$ at 1.96 TeV

$0.02^{+0.11}_{-0.08} \pm 0.02$ WEI 09A BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.07^{+0.30}_{-0.23} \pm 0.02$ AALTONEN 11L CDF Repl. by AALTONEN 12I

$A_{FB}(B \rightarrow X_s\ell^+\ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.74 ± 0.54 ¹SATO 16 BELL $e^+e^- \rightarrow \gamma(4S)$

¹ Uses the sum of 10 exclusive X_s modes in the range $M(X_s) > 1.1 \text{ GeV}/c^2$. Uncertainty is statistical only.

$F_S(B \rightarrow K\ell^+\ell^-) (q^2 > 0.1 \text{ GeV}^2/c^4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.81^{+0.58}_{-0.61} \pm 0.46$ ¹AUBERT,B 06J BABR $e^+e^- \rightarrow \gamma(4S)$

¹ Results with different q^2 cuts are also reported.

$A_{FB}(B \rightarrow K p \bar{p}) (m_{p \bar{p}} < 2.85 \text{ GeV}/c^2)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.495 \pm 0.012 \pm 0.007$	¹ AAIJ	14AF LHCB	pp at 7, 8 TeV

¹ Measured in $B^+ \rightarrow K^+ p \bar{p}$ decays. **$A_{FB}(B \rightarrow \pi p \bar{p}) (m_{p \bar{p}} < 2.85 \text{ GeV}/c^2)$**

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.409 \pm 0.033 \pm 0.006$	¹ AAIJ	14AF LHCB	pp at 7, 8 TeV

¹ Measured in $B^+ \rightarrow \pi^+ p \bar{p}$ decays. **A_{FB} in $B \rightarrow \bar{D}^* e^+ \nu_e$**

VALUE	DOCUMENT ID	TECN	COMMENT
$0.227 \pm 0.020 \pm 0.006$	¹ PRIM	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ This is the B^+ and B^0 average. **A_{FB} in $B \rightarrow \bar{D}^* \mu^+ \nu_\mu$**

VALUE	DOCUMENT ID	TECN	COMMENT
$0.256 \pm 0.020 \pm 0.005$	¹ PRIM	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ This is the B^+ and B^0 average. **$\Delta(A_{FB}) = (A_{FB}^\mu - A_{FB}^e)$ in $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$**

VALUE	DOCUMENT ID	TECN	COMMENT
$0.028 \pm 0.028 \pm 0.008$	¹ PRIM	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ This is the B^+ and B^0 average.**ISOSPIN ASYMMETRY** Δ_{0-} is defined as

$$\frac{\Gamma(\bar{B}^0 \rightarrow f_d) - \Gamma(B^- \rightarrow f_u)}{\Gamma(\bar{B}^0 \rightarrow f_d) + \Gamma(B^- \rightarrow f_u)},$$

the isospin asymmetry of inclusive neutral and charged B decay.

 $\Delta_{0-}(B(B \rightarrow X_s \gamma))$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.005 ± 0.020 OUR AVERAGE			
$-0.0048 \pm 0.0149 \pm 0.0150$	¹ WATANUKI	19 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.006 \pm 0.058 \pm 0.026$	AUBERT,B	05R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Using a sum-of-exclusive technique with $m_{X_s} < 2.8 \text{ GeV}/c^2$. **$\Delta_{0-}(B(B \rightarrow X_{s+d} \gamma))$**

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.06 \pm 0.15 \pm 0.07$	¹ AUBERT	08o BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side. The result is for $E_\gamma > 2.2 \text{ GeV}$.

$\Delta_{0+}(B \rightarrow K^*(892)\gamma)$

Δ_{0+} describes the isospin asymmetry between $\Gamma(B^0 \rightarrow K^*(892)^0\gamma)$ and $\Gamma(B^+ \rightarrow K^*(892)^+\gamma)$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.063 ± 0.017 OUR AVERAGE			
$0.062 \pm 0.015 \pm 0.013$	¹ HORIGUCHI 17	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.066 \pm 0.021 \pm 0.022$	² AUBERT 09AO	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.050 \pm 0.045 \pm 0.037$	³ AUBERT,BE 04A	BABR	Repl. by AUBERT 09AO
$0.012 \pm 0.044 \pm 0.026$	NAKAO 04	BELL	Repl. by HORIGUCHI 17
¹ Uses $B(\Upsilon(4S) \rightarrow B^+B^-) = (51.4 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = (48.6 \pm 0.6)\%$.			
² Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays and the lifetime ratio $\tau_{B^+}/\tau_{B^0} = 1.071 \pm 0.009$. The 90% CL interval is $0.017 < \Delta_{0+} < 0.116$			
³ Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^{+/0} = 1.006 \pm 0.048$ and the lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$. The 90% CL interval is $-0.046 < \Delta_{0+} < 0.146$.			

 $\Delta_{\rho\gamma} = \Gamma(B^+ \rightarrow \rho^+\gamma) / (2 \cdot \Gamma(B^0 \rightarrow \rho^0\gamma)) - 1$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.46 ± 0.17 OUR AVERAGE			
$-0.43^{+0.25}_{-0.22} \pm 0.10$	AUBERT 08BH	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.48^{+0.21+0.08}_{-0.19-0.09}$	TANIGUCHI 08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Delta_{0-}(B(B \rightarrow K\ell^+\ell^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.15 ± 0.06 OUR AVERAGE	Error includes scale factor of 1.2.		
$-0.31^{+0.13}_{-0.11} \pm 0.01$	¹ CHOUDHURY 21	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.10^{+0.08}_{-0.09} \pm 0.02$	² AAIJ 14M	LHCB	pp at 7, 8 TeV
$-0.09^{+0.08}_{-0.08} \pm 0.02$	³ AAIJ 14M	LHCB	pp at 7, 8 TeV
$-0.58^{+0.29}_{-0.37} \pm 0.02$	⁴ LEES 12S	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.35^{+0.23}_{-0.27}$	⁵ AAIJ 12AH	LHCB	Repl. by AAIJ 14M
$-1.43^{+0.56}_{-0.85} \pm 0.05$	^{6,7} AUBERT 09T	BABR	Repl. by LEES 12S
$-0.31^{+0.17}_{-0.14} \pm 0.08$	^{8,9} WEI 09A	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
¹ For $1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$ using both $\mu^+\mu^-$ and e^+e^- as a lepton pair. Measurements in other q^2 bins are also reported.			
² For $1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$ using $\mu^+\mu^-$ as a lepton pair and assuming isospin symmetry for the $B \rightarrow J/\psi(1S)K$. Measurements in other q^2 bins are also reported.			
³ For $15.0 < q^2 < 19.0 \text{ GeV}^2/c^4$ using $\mu^+\mu^-$ as a lepton pair and assuming isospin symmetry for the $B \rightarrow J/\psi(1S)K$. Measurements in other q^2 bins are also reported.			
⁴ For $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$. Measurements in other q^2 bins are also reported.			
⁵ For $1 < q^2 < 6 \text{ GeV}^2/c^4$.			
⁶ For $0.1 < m_{\ell^+\ell^-}^2 < 7.02 \text{ GeV}^2/c^4$.			

⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁸ Superseded by CHOUDHURY 21.

⁹ For $q^2 < 8.68 \text{ GeV}^2/c^4$.

$\Delta_0(B(B \rightarrow K^* \ell^+ \ell^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.03^{+0.08}_{-0.07}$ OUR AVERAGE	Error includes scale factor of 1.2.		
$0.00^{+0.12}_{-0.10} \pm 0.02$	¹ AAIJ	14M LHCb	pp at 7, 8 TeV
$0.06^{+0.10}_{-0.09} \pm 0.02$	² AAIJ	14M LHCb	pp at 7, 8 TeV
$-0.25^{+0.20}_{-0.17} \pm 0.03$	³ LEES	12S BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.29 \pm 0.16 \pm 0.09$	⁴ WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.15 ± 0.16	⁵ AAIJ	12AH LHCb	Repl. by AAIJ 14M
$-0.56^{+0.17}_{-0.15} \pm 0.03$	^{6,7} AUBERT	09T BABR	Repl. by LEES 12S

¹ For $1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$ using $\mu^+ \mu^-$ as a lepton pair and assuming isospin symmetry for the $B(B \rightarrow J/\psi(1S) K^*(892))$. Measurements in other q^2 bins are also reported.

² For $15.0 < q^2 < 22.0 \text{ GeV}^2/c^4$ using $\mu^+ \mu^-$ as a lepton pair and assuming isospin symmetry for the $B(B \rightarrow J/\psi(1S) K^*(892))$. Measurements in other q^2 bins are also reported.

³ For $0.10 < q^2 < 8.12 \text{ GeV}^2/c^4$. Measurements in other q^2 bins are also reported.

⁴ For $q^2 < 8.68 \text{ GeV}^2/c^4$.

⁵ For $1 < q^2 < 6 \text{ GeV}^2/c^4$.

⁶ For $0.1 < m_{\ell^+ \ell^-}^2 < 7.02 \text{ GeV}^2/c^4$.

⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Delta_0(B(B \rightarrow K^{(*)} \ell^+ \ell^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.45 ± 0.17 OUR AVERAGE	Error includes scale factor of 1.7.		
$-0.64^{+0.15}_{-0.14} \pm 0.03$	^{1,2} AUBERT	09T BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.30^{+0.12}_{-0.11} \pm 0.08$	³ WEI	09A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
¹ For $0.1 < m_{\ell^+ \ell^-}^2 < 7.02 \text{ GeV}^2/c^4$. ² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. ³ For $q^2 < 8.68 \text{ GeV}^2/c^2$.			

$B \rightarrow X_c \ell \nu$ HADRONIC MASS MOMENTS

$\langle M_X^2 - \overline{M}_D^2 \rangle$ (First Moments)

VALUE (GeV^2)	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.08 OUR AVERAGE	Error includes scale factor of 1.8.		
$0.467 \pm 0.038 \pm 0.068$	¹ ACOSTA	05F CDF	$p\bar{p}$ at 1.96 TeV
$0.293 \pm 0.012 \pm 0.058$	² CSORNA	04 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.251 \pm 0.023 \pm 0.062$	³ CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;
² Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.
³ The leptons are required to have $P_\ell > 1.5$ GeV/c.

 $\langle M_X^2 \rangle$ (First Moments)

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
4.156 ± 0.029 OUR AVERAGE			
4.144 ± 0.028 ± 0.022	¹ SCHWANDA 07	BELL	$e^+e^- \rightarrow \gamma(4S)$
4.18 ± 0.04 ± 0.03	¹ AUBERT,B 04	BABR	$e^+e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.

 $\langle (M_X^2 - \overline{M}_X^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.55 ± 0.08 OUR AVERAGE			
0.515 ± 0.061 ± 0.064	¹ SCHWANDA 07	BELL	$e^+e^- \rightarrow \gamma(4S)$
0.629 ± 0.031 ± 0.143	² CSORNA 04	CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.05 ± 0.26 ± 0.13	³ ACOSTA 05F	CDF	$p\bar{p}$ at 1.96 TeV
0.576 ± 0.048 ± 0.168	¹ CRONIN-HEN..01B	CLE2	$e^+e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.
² Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.
³ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;

 $\langle (M_X^2 - \overline{M}_D^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.639 ± 0.056 ± 0.178	¹ CRONIN-HEN..01B	CLE2	$e^+e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_\ell > 1.5$ GeV/c.

 $B \rightarrow X_c \ell \nu$ LEPTON MOMENTUM MOMENTS **$R_0 (\Gamma_{E_l > 1.7 \text{ GeV}} / \Gamma_{E_l > 1.5 \text{ GeV}})$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.6187 ± 0.0014 ± 0.0016	¹ MAHMOOD 03	CLE2	$e^+e^- \rightarrow \gamma(4S)$


- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.

 $R_1 (\langle E_l \rangle_{E_l > 1.5 \text{ GeV}})$

VALUE	DOCUMENT ID	TECN	COMMENT
1.7797 ± 0.0018 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
1.7743 ± 0.0019 ± 0.0014	¹ AUBERT,B 04A	BABR	$e^+e^- \rightarrow \gamma(4S)$
1.7792 ± 0.0021 ± 0.0027	² MAHMOOD 04	CLEO	$e^+e^- \rightarrow \gamma(4S)$
1.7810 ± 0.0007 ± 0.0009	³ MAHMOOD 03	CLE2	$e^+e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.

- ² Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.
- ³ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.



$R_2 (\langle E_l^2 - \bar{E}_l^2 \rangle_{E_l > 1.5 \text{ GeV}})$

VALUE (10^{-3} GeV^2)	DOCUMENT ID	TECN	COMMENT
30.8 ± 0.8 OUR AVERAGE			
$30.3 \pm 0.9 \pm 0.5$	¹ AUBERT,B	04A	BABR $e^+ e^- \rightarrow \gamma(4S)$
$31.6 \pm 0.8 \pm 1.0$	² MAHMOOD	04	CLEO $e^+ e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.
- ² Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.

$R_3 (\langle E_l^3 - \bar{E}_l^3 \rangle_{E_l > 1.5 \text{ GeV}})$

VALUE (10^{-3} GeV^3)	DOCUMENT ID	TECN	COMMENT
$2.12 \pm 0.47 \pm 0.20$	¹ AUBERT,B	04A	BABR $e^+ e^- \rightarrow \gamma(4S)$

- ¹ The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.

$B \rightarrow X_s \gamma$ PHOTON ENERGY MOMENTS **$\langle E_\gamma \rangle$**

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
2.314$\pm$0.011 OUR AVERAGE			
2.346 $\pm$ 0.018 $^{+0.027}_{-0.022}$	^{1,2} LEES	12U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.304 $\pm$ 0.014 $\pm$ 0.017	^{2,3} LEES	12V BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.311 $\pm$ 0.009 $\pm$ 0.015	³ LIMOSANI	09 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.289 $\pm$ 0.058 $\pm$ 0.027	^{3,4} AUBERT	08O BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.309 $\pm$ 0.023 $\pm$ 0.023	^{2,3} SCHWANDA	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.288 $\pm$ 0.025 $\pm$ 0.023	³ AUBERT,BE	06B BABR	Repl. by LEES 12V

¹ LEES 12U uses $E_\gamma > 1.897$ GeV to calculate the moments; the moments are used to calculate the HQET parameters $m_b = 4.579^{+0.032}_{-0.029}$ GeV/ c^2 and $\mu_\pi^2 = 0.257^{+0.034}_{-0.039}$ GeV² in the shape function model. The same HQET parameters are also determined in the kinetic model.

² Results for different E_γ threshold values are also measured.

³ The result is for $E_\gamma > 1.9$ GeV.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

 $\langle E_\gamma^2 \rangle - \langle E_\gamma \rangle^2$

VALUE (10 ⁻² GeV ²)	DOCUMENT ID	TECN	COMMENT
3.03$\pm$0.25 OUR AVERAGE			
2.11 $\pm$ 0.57 $^{+0.55}_{-0.69}$	^{1,2} LEES	12U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
3.62 $\pm$ 0.33 $\pm$ 0.33	^{2,3} LEES	12V BABR	$e^+e^- \rightarrow \Upsilon(4S)$
3.02 $\pm$ 0.19 $\pm$ 0.30	³ LIMOSANI	09 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
3.34 $\pm$ 1.24 $\pm$ 0.62	^{3,4} AUBERT	08O BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.17 $\pm$ 0.60 $\pm$ 0.55	^{2,3} SCHWANDA	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.28 $\pm$ 0.40 $\pm$ 0.43	³ AUBERT,BE	06B BABR	Repl. by LEES 12V

¹ LEES 12U uses $E_\gamma > 1.897$ GeV to calculate the moments; the moments are used to calculate the HQET parameters $m_b = 4.579^{+0.032}_{-0.029}$ GeV/ c^2 and $\mu_\pi^2 = 0.257^{+0.034}_{-0.039}$ GeV² in the shape function model. The same HQET parameters are also determined in the kinetic model.

² Results for different E_γ threshold values are also measured.

³ The result is for $E_\gamma > 1.9$ GeV.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

$B^\pm/B^0$ ADMIXTURE REFERENCES

ADACHI	24O	PR D110 072020	I. Adachi <i>et al.</i>	(BELLE II Collab.)
FERLEWICZ	24	PR D110 072005	D. Ferlewicz <i>et al.</i>	(BELLE Collab.)
AAIJ	23AR	PRL 131 111802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AGGARWAL	23	PRL 131 051804	L. Aggarwal <i>et al.</i>	(BELLE II Collab.)
CAO	23	PRL 131 211801	L. Cao <i>et al.</i>	(BELLE Collab.)
PRIM	23	PR D108 012002	M.T. Prim <i>et al.</i>	(BELLE Collab.)
ZHUKOVA	23	JHEP 2308 131	V. Zhukova <i>et al.</i>	(BELLE Collab.)
CAO	21A	PR D104 012008	L. Cao <i>et al.</i>	(BELLE Collab.)
CHOUDHURY	21	JHEP 2103 105	S. Choudhury <i>et al.</i>	(BELLE Collab.)
WEHLE	21	PRL 126 161801	S. Wehle <i>et al.</i>	(BELLE Collab.)
AAIJ	20Y	PRL 125 011802	R. Aaij <i>et al.</i>	(LHCb Collab.)
CARIA	20	PRL 124 161803	G. Caria <i>et al.</i>	(BELLE Collab.)
AAIJ	19AD	PR D100 031102	R. Aaij <i>et al.</i>	(LHCb Collab.)
WATANUKI	19	PR D99 032012	S. Watanuki <i>et al.</i>	(BELLE Collab.)
GRYGIER	17	PR D96 091101	J. Grygier <i>et al.</i>	(BELLE Collab.)
HIROSE	17	PRL 118 211801	S. Hirose <i>et al.</i>	(BELLE Collab.)
Also		PR D97 012004	S. Hirose <i>et al.</i>	(BELLE Collab.)
HORIGUCHI	17	PRL 119 191802	T. Horiguchi <i>et al.</i>	(BELLE Collab.)
LEES	17B	PR D95 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	16B	JHEP 1602 104	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
KHACHATRYAN	16D	PL B753 424	V. Khachatryan <i>et al.</i>	(CMS Collab.)
LEES	16	PRL 116 041801	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	16C	PR D93 052015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SATO	16	PR D93 032008	Y. Sato <i>et al.</i>	(BELLE Collab.)
Also		PR D93 059901 (errat.)	Y. Sato <i>et al.</i>	(BELLE Collab.)
HUSCHLE	15	PR D92 072014	M. Huschle <i>et al.</i>	(BELLE Collab.)
PESANTEZ	15	PRL 114 151601	L. Pesantez <i>et al.</i>	(BELLE Collab.)
SAITO	15	PR D91 052004	T. Saito <i>et al.</i>	(BELLE Collab.)
AAIJ	14AF	PRL 113 141801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14M	JHEP 1406 133	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	14D	PRL 112 211802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	14K	PR D90 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	13H	JHEP 1302 105	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13Y	JHEP 1308 131	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHATRCHYAN	13BL	PL B727 77	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13M	PR D88 032012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12AH	JHEP 1207 133	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12U	PRL 108 181806	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	12I	PRL 108 081807	T. Aaltonen <i>et al.</i>	(CDF Collab.)
LEES	12	PR D85 011102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12D	PRL 109 101802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
Also		PR D88 072012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12R	PR D86 032004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12S	PR D86 032012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12U	PR D86 052012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12V	PRL 109 191801	J.P. Lees	(BABAR Collab.)
Also		PR D86 112008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AALTONEN	11AI	PRL 107 201802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11L	PRL 106 161801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
DEL-AMO-SANCHEZ	11	PR D83 031103	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
GAMBINO	11	JHEP 1109 055	P. Gambino	(LCGT)
AUBERT	10	PRL 104 011802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	10A	PR D81 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SANCHEZ	10M	PR D82 051101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SANCHEZ	10Q	PR D82 112002	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
NISHIMURA	10	PRL 105 191803	K. Nishimura <i>et al.</i>	(BELLE Collab.)
URQUIJO	10	PRL 104 021801	P. Urquijo <i>et al.</i>	(BELLE Collab.)
AUBERT	09AO	PRL 103 211802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09N	PR D79 031102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09T	PRL 102 091803	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		EPAPS Document No. E-PRLTAO-102-060910		(BABAR Collab.)
AUBERT	09U	PRL 102 161803	B. Aubert <i>et al.</i>	(BABAR Collab.)
LIMOSANI	09	PRL 103 241801	A. Limosani <i>et al.</i>	(BELLE Collab.)
WEI	09A	PRL 103 171801	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
Also		EPAPS Supplement	J.-T. Wei <i>et al.</i>	(BELLE Collab.)

AUBERT	08AS	PRL 100 171802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BC	PR D78 072007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BH	PR D78 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BJ	PRL 101 171804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08N	PRL 100 021801	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D79 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08O	PR D77 051103	B. Aubert <i>et al.</i>	(BABAR Collab.)
SCHWANDA	08	PR D78 032016	C. Schwanda <i>et al.</i>	(BELLE Collab.)
TANIGUCHI	08	PRL 101 111801	N. Taniguchi <i>et al.</i>	(BELLE Collab.)
WEI	08A	PR D78 011101	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
AUBERT	07AG	PRL 99 051801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07C	PR D75 012003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07E	PRL 98 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07L	PRL 98 151802	B. Aubert <i>et al.</i>	(BABAR Collab.)
GAMBINO	07	JHEP 0710 058	P. Gambino <i>et al.</i>	
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)
SCHWANDA	07	PR D75 032005	C. Schwanda <i>et al.</i>	(BELLE Collab.)
URQUIJO	07	PR D75 032001	P. Urquijo <i>et al.</i>	(BELLE Collab.)
AUBERT	06H	PR D73 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06J	PR D73 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	06Y	PR D74 091105	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06B	PRL 97 171803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BUCHMUEL...	06	PR D73 073008	O.L. Buchmueller, H.U. Flacher	(RHBL)
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	06	PRL 96 251801	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	06	PRL 96 221601	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
ABAZOV	05O	PRL 95 171803	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	05F	PR D71 051103	D. Acosta <i>et al.</i>	(CDF Collab.)
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05M	PRL 95 142003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05R	PR D72 052004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05X	PRL 95 111801	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PRL 97 019903 (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHOI	05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
IWASAKI	05	PR D72 092005	M. Iwasaki <i>et al.</i>	(BELLE Collab.)
LIMOSANI	05	PL B621 28	A. Limosani <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	05	PR D72 011101	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
NISHIDA	05	PL B610 23	S. Nishida <i>et al.</i>	(BELLE Collab.)
OKABE	05	PL B614 27	T. Okabe <i>et al.</i>	(BELLE Collab.)
ABDALLAH	04D	EPJ C33 213	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04I	PRL 92 071802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04S	PR D69 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04X	PRL 93 011803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04	PR D69 111103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04A	PR D69 111104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04E	PRL 93 021804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04F	PRL 93 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04I	PRL 93 081802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
CSORNA	04	PR D70 032002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
KOPPENBURG	04	PRL 93 061803	P. Koppenburg <i>et al.</i>	(BELLE Collab.)
MAHMOOD	04	PR D70 032003	A.H. Mahmodd <i>et al.</i>	(CLEO Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
NISHIDA	04	PRL 93 031803	S. Nishida <i>et al.</i>	(BELLE Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	03	PR D67 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	03F	PR D67 032002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BABAR Collab.)
BONVICINI	03	PR D68 011101	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
KANEKO	03	PRL 90 021801	J. Kaneko <i>et al.</i>	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)
MAHMOOD	03	PR D67 072001	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02L	PRL 89 011803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02Y	PL B547 181	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	02	PRL 89 282001	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT	02G	PR D65 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BABAR Collab.)
BORNHEIM	02	PRL 88 231803	A. Bornheim <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABE	01F	PL B511 151	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01J	PR D64 072001	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
CHEN	01	PR D63 031102	S. Chen <i>et al.</i>	(CLEO Collab.)
CHEN	01C	PRL 87 251807	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	01	PRL 86 5661	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HENNESSY	01B	PRL 87 251808	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PDG	01	Unofficial 2001 WWW edition		
ABREU	00R	PL B475 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	98	PR D57 3847	M. Bishai <i>et al.</i>	(CLEO Collab.)
BONVICINI	98	PR D57 6604	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BROWDER	98	PRL 81 1786	T.E. Browder <i>et al.</i>	(CLEO Collab.)
COAN	98	PRL 80 1150	T.E. Coan <i>et al.</i>	(CLEO Collab.)
GLENN	98	PRL 80 2289	S. Glenn <i>et al.</i>	(CLEO Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AMMAR	97	PR D55 13	R. Ammar <i>et al.</i>	(CLEO Collab.)
BARISH	97	PRL 79 3599	B. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97B	ZPHY C73 601	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBBONS	97B	PR D56 3783	L. Gibbons <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96D	PL B374 256	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
KUBOTA	96	PR D53 6033	Y. Kubota <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)
ALAM	95	PRL 74 2885	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95B	PR D52 2661	R. Balest <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95B	PL B345 103	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	94C	ZPHY C62 371	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94J	ZPHY C61 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
PROCARIO	94	PRL 73 1472	M. Procario <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93	ZPHY C57 533	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93H	PL B318 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93I	ZPHY C58 191	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	93	PL B311 307	M. Artuso	(SYRA)
BARTELT	93B	PRL 71 4111	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
HENDERSON	92	PR D45 2212	S. Henderson <i>et al.</i>	(CLEO Collab.)
LESIK	92	ZPHY C55 33	T. Lesiak <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91H	ZPHY C52 353	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
YANAGISAWA	91	PRL 66 2436	C. Yanagisawa <i>et al.</i>	(CUSB II Collab.)
ALBRECHT	90	PL B234 409	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90H	PL B249 359	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also		PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
FULTON	90	PRL 64 16	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
PDG	90	PL B239 1	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)
ALBRECHT	89K	ZPHY C42 519	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ISGUR	89B	PR D39 799	N. Isgur <i>et al.</i>	(TNT0, CIT)
WACHS	89	ZPHY C42 33	K. Wachs <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	88E	PL B210 263	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88H	PL B210 258	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KOERNER	88	ZPHY C38 511	J.G. Korner, G.A. Schuler	(MAINZ, DESY)

ALAM	87	PRL 59 22	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALAM	87B	PRL 58 1814	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87H	PL B187 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEAN	87	PR D35 3533	A. Bean <i>et al.</i>	(CLEO Collab.)
BEHREND	87	PRL 59 407	S. Behrends <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	87	PR D35 19	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BALTRUSAITIS...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BORTOLETTO	86	PRL 56 800	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
HAAS	86	PRL 56 2781	J. Haas <i>et al.</i>	(CLEO Collab.)
ALBRECHT	85H	PL 162B 395	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
CSORNA	85	PRL 54 1894	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
HAAS	85	PRL 55 1248	J. Haas <i>et al.</i>	(CLEO Collab.)
AVERY	84	PRL 53 1309	P. Avery <i>et al.</i>	(CLEO Collab.)
CHEN	84	PRL 52 1084	A. Chen <i>et al.</i>	(CLEO Collab.)
LEVMAN	84	PL 141B 271	G.M. Levman <i>et al.</i>	(CUSB Collab.)
ALAM	83B	PRL 51 1143	M.S. Alam <i>et al.</i>	(CLEO Collab.)
GREEN	83	PRL 51 347	J. Green <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83B	PL 130B 444	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALTARELLI	82	NP B208 365	G. Altarelli <i>et al.</i>	(ROMA, INFN, FRAS)
BRODY	82	PRL 48 1070	A.D. Brody <i>et al.</i>	(CLEO Collab.)
GIANNINI	82	NP B206 1	G. Giannini <i>et al.</i>	(CUSB Collab.)
BEBEK	81	PRL 46 84	C. Bebek <i>et al.</i>	(CLEO Collab.)
CHADWICK	81	PRL 46 88	K. Chadwick <i>et al.</i>	(CLEO Collab.)
ABRAMS	80	PRL 44 10	G.S. Abrams <i>et al.</i>	(SLAC, LBL)