

$J/\psi(1S)$

$$J^G(J^{PC}) = 0^-(1^{--})$$

$J/\psi(1S)$ MASS

VALUE (MeV)	EPTS	DOCUMENT ID	TECN	COMMENT
3096.900±0.006 OUR AVERAGE				
3096.900±0.002±0.006		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3096.66 ±0.19 ±0.02	6.1k	⁴ AAIJ 15BI	LHCB	$pp \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ±0.3		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3096.93 ±0.09	502	⁵ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ±1		⁶ BRANDELIK 79C	DASP	e^+e^-

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

$J/\psi(1S)$ WIDTH

VALUE (keV)	EPTS	DOCUMENT ID	TECN	COMMENT
92.6 ± 1.7 OUR AVERAGE		Error includes scale factor of 1.1.		
92.45± 1.40±1.48		¹ ANASHIN 20	KEDR	e^+e^-
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI 95B	BES	e^+e^-
91 ±11 ±6		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 $\begin{smallmatrix} +6.1 \\ -5.8 \end{smallmatrix}$		⁴ HSUEH 92	RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
92.94± 1.83		^{5,6} ANASHIN 18A	KEDR	e^+e^-
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there.² Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.³ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

⁴Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

⁵Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.

⁶Superseded by ANASHIN 20 that is based on the same dataset.

⁷Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	$(87.7 \pm 0.5) \%$	
Γ_2 virtual $\gamma \rightarrow$ hadrons	$(13.46 \pm 0.07) \%$	
Γ_3 ggg	$(64.1 \pm 1.0) \%$	
Γ_4 $\gamma g g$	$(8.8 \pm 1.1) \%$	
Γ_5 e^+e^-	$(5.971 \pm 0.032) \%$	
Γ_6 $e^+e^- \gamma$	[a] $(8.8 \pm 1.4) \times 10^{-3}$	
Γ_7 $\mu^+\mu^-$	$(5.961 \pm 0.033) \%$	
Γ_8 $e^+e^-e^+e^-$	$(5.5 \pm 0.5) \times 10^{-5}$	
Γ_9 $e^+e^- \mu^+\mu^-$	$(3.53 \pm 0.26) \times 10^{-5}$	
Γ_{10} $\mu^+\mu^- \mu^+\mu^-$	$(1.11 \pm 0.11) \times 10^{-6}$	

Decays involving hadronic resonances

Γ_{11} $\rho\pi$	$(1.88 \pm 0.12) \%$	S=2.6
Γ_{12} $\rho^0\pi^0$	$(6.2 \pm 0.6) \times 10^{-3}$	
Γ_{13} $a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0$	$(2.8 \pm 0.6) \times 10^{-3}$	
Γ_{14} $a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow 2(\pi^+\pi^-)\pi^0$	$(3.7 \pm 0.7) \times 10^{-3}$	
Γ_{15} $a_2(1320)\rho$	$(1.09 \pm 0.22) \%$	
Γ_{16} $\eta\pi^+\pi^-$	$(3.8 \pm 0.7) \times 10^{-4}$	
Γ_{17} $\eta\rho$	$(1.93 \pm 0.23) \times 10^{-4}$	
Γ_{18} $\eta\pi^+\pi^-\pi^0$	$(1.17 \pm 0.20) \%$	
Γ_{19} $\eta\pi^+\pi^-3\pi^0$	$(4.9 \pm 1.0) \times 10^{-3}$	
Γ_{20} $\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	
Γ_{21} $\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0$	$< 2.52 \times 10^{-4}$	CL=90%
Γ_{22} ηK^+K^-	$(8.6 \pm 3.0) \times 10^{-4}$	
Γ_{23} $\eta K^\pm K_S^0 \pi^\mp$	[b] $(2.2 \pm 0.4) \times 10^{-3}$	
Γ_{24} $\eta K^*(892)^0 \bar{K}^*(892)^0$	$(1.15 \pm 0.26) \times 10^{-3}$	
Γ_{25} $\rho\eta'(958)$	$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6
Γ_{26} $\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0$	$(2.8 \pm 0.8) \%$	
Γ_{27} $\rho^+ \rho^- \pi^+ \pi^- \pi^0$	$(6 \pm 4) \times 10^{-3}$	
Γ_{28} $\rho^+ K^+ K^- \pi^- + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	$(3.5 \pm 0.8) \times 10^{-3}$	

Γ_{29}	$\rho^\mp K^\pm K_S^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{30}	$h_1(1415)\eta' \rightarrow \gamma\eta\eta'$		
Γ_{31}	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$		
Γ_{32}	$\rho(1450)\pi$	seen	
Γ_{33}	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	$(2.2 \pm 1.1) \times 10^{-4}$	
Γ_{34}	$\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.3 \pm 0.6) \times 10^{-4}$	
Γ_{35}	$\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	$(2.7 \pm 0.6) \times 10^{-4}$	
Γ_{36}	$\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958)$	$(3.3 \pm 0.7) \times 10^{-6}$	
Γ_{37}	$\rho(1700)\pi$	seen	
Γ_{38}	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 1.1) \times 10^{-4}$	
Γ_{39}	$\rho(2150)\pi$	seen	
Γ_{40}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	$(10 \pm 40) \times 10^{-6}$	
Γ_{41}	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$		
Γ_{42}	$\omega\pi^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
Γ_{43}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 0.7) \times 10^{-5}$	
Γ_{44}	$\omega\pi^+\pi^-$	$(8.5 \pm 1.0) \times 10^{-3}$	S=1.3
Γ_{45}	$\omega\pi^0\pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$	
Γ_{46}	$\omega 3\pi^0$	$(1.9 \pm 0.6) \times 10^{-3}$	
Γ_{47}	$\omega f_2(1270)$	$(4.3 \pm 0.6) \times 10^{-3}$	
Γ_{48}	$\omega\eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
Γ_{49}	$\omega\pi^+\pi^-\pi^0$	$(4.0 \pm 0.7) \times 10^{-3}$	
Γ_{50}	$\omega\pi^0\eta$	$(3.4 \pm 1.7) \times 10^{-4}$	
Γ_{51}	$\omega\pi^+\pi^+\pi^-\pi^-$	$(8.5 \pm 3.4) \times 10^{-3}$	
Γ_{52}	$\omega\pi^+\pi^-2\pi^0$	$(3.3 \pm 0.5) \%$	
Γ_{53}	$\omega\eta'\pi^+\pi^-$	$(1.12 \pm 0.13) \times 10^{-3}$	
Γ_{54}	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$	
Γ_{55}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{56}	$\omega f_0(1710) \rightarrow \omega K \bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$	
Γ_{57}	$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$	
Γ_{58}	$\omega f_2'(1525)$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{59}	$\omega X(1835) \rightarrow \omega p \bar{p}$	$< 3.9 \times 10^{-6}$	CL=95%
Γ_{60}	$\omega K^+ K^- \eta$	$(3.33 \pm 0.12) \times 10^{-4}$	
Γ_{61}	$\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-$	$< 6.2 \times 10^{-5}$	
Γ_{62}	$\omega K^+ K^-$	$(1.52 \pm 0.31) \times 10^{-3}$	
Γ_{63}	$\omega K^\pm K_S^0 \pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$	
Γ_{64}	$\omega K \bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{65}	$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	
Γ_{66}	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$	
Γ_{67}	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$	
Γ_{68}	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$	
Γ_{69}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$	
Γ_{70}	$\eta' h_1(1415) \rightarrow \gamma \eta' \eta'$	$(4.7 \pm \frac{1.1}{2.0}) \times 10^{-7}$	

Γ_{71}	$\bar{K} K^*(892) + \text{c.c.}$	seen	
Γ_{72}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(4.8 \pm 0.5) \times 10^{-3}$	
Γ_{73}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm \frac{0.8}{1.0}) \times 10^{-3}$	S=2.9
Γ_{74}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm \frac{0.13}{0.20}) \times 10^{-3}$	
Γ_{75}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{76}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	
Γ_{77}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	
Γ_{78}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$	
Γ_{79}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	
Γ_{80}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$	
Γ_{81}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$	
Γ_{82}	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$	
Γ_{83}	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm \frac{0.6}{0.5}) \times 10^{-6}$	
Γ_{84}	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$	
Γ_{85}	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm \frac{1.0}{0.6}) \times 10^{-3}$	
Γ_{86}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$	
Γ_{87}	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm \frac{0.22}{0.40}) \times 10^{-3}$	
Γ_{88}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	
Γ_{89}	$K^*(1410) \bar{K} + \text{c.c.}$	seen	
Γ_{90}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$	
Γ_{91}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(8 \pm 5) \times 10^{-5}$	
Γ_{92}	$K_2^*(1430) \bar{K} + \text{c.c.}$	seen	
Γ_{93}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$	
Γ_{94}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{95}	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%
Γ_{96}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm \frac{0.25}{0.19}) \times 10^{-4}$	
Γ_{97}	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$	
Γ_{98}	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	
Γ_{99}	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$	

Γ_{100}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	
Γ_{101}	$K_2^*(1430)^- K^*(892)^+ +$ $\text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- +$ c.c.	$(4 \pm 4) \times 10^{-4}$	
Γ_{102}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%
Γ_{103}	$\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	
Γ_{104}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.10 \pm_{-0.14}^{+0.60}) \times 10^{-5}$	
Γ_{105}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(6.2 \pm_{-1.6}^{+2.9}) \times 10^{-6}$	
Γ_{106}	$K_1(1270)^\pm K^\mp$	$< 3.0 \times 10^{-3}$	CL=90%
Γ_{107}	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$	
Γ_{108}	$a_2(1320)^\pm \pi^\mp$	$[b] < 4.3 \times 10^{-3}$	CL=90%
Γ_{109}	$\phi \pi^0$	$3 \times 10^{-6} \text{ or } 1 \times 10^{-7}$	
Γ_{110}	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7
Γ_{111}	$\phi \pi^0 \pi^0$	$(4.9 \pm 1.0) \times 10^{-4}$	
Γ_{112}	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$	
Γ_{113}	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4}$	S=1.2
Γ_{114}	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2
Γ_{115}	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$	
Γ_{116}	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{117}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.58 \pm 0.34) \times 10^{-4}$	
Γ_{118}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.7 \pm 0.5) \times 10^{-4}$	
Γ_{119}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$	
Γ_{120}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$	
Γ_{121}	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{122}	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$		
Γ_{123}	$\phi(1680)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(6.7 \pm 1.1) \times 10^{-6}$	
Γ_{124}	$X(2000)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(1.70 \pm_{-0.23}^{+0.50}) \times 10^{-6}$	
Γ_{125}	$h_1(1900)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(8.4 \pm_{-1.3}^{+1.4}) \times 10^{-6}$	
Γ_{126}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$	
Γ_{127}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{128}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$	
Γ_{129}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$	
Γ_{130}	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$	
Γ_{131}	$\phi f_2'(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7
Γ_{132}	$\phi X(1835) \rightarrow \phi p \bar{p}$	$< 2.1 \times 10^{-7}$	CL=90%
Γ_{133}	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	$< 2.8 \times 10^{-4}$	CL=90%
Γ_{134}	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	$< 6.13 \times 10^{-5}$	CL=90%

Γ_{135}	$\phi K \bar{K}$		$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3
Γ_{136}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$		$(3.6 \pm 0.6) \times 10^{-4}$	
Γ_{137}	$\phi K^+ K^-$		$(8.2 \pm 1.1) \times 10^{-4}$	
Γ_{138}	$\phi K_S^0 K_S^0$		$(5.8 \pm 1.5) \times 10^{-4}$	
Γ_{139}	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{140}	$\phi K^*(892) \bar{K} + \text{c.c.}$		$(2.18 \pm 0.23) \times 10^{-3}$	
Γ_{141}	$b_1(1235)^\pm \pi^\mp$	[b]	$(3.0 \pm 0.5) \times 10^{-3}$	
Γ_{142}	$b_1(1235)^0 \pi^0$		$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{143}	$f'_2(1525) K^+ K^-$		$(1.04 \pm 0.35) \times 10^{-3}$	
Γ_{144}	$\Delta(1232)^+ \bar{p}$		$< 1 \times 10^{-4}$	CL=90%
Γ_{145}	$\Delta(1232)^{++} \bar{p} \pi^-$		$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{146}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{147}	$\bar{\Sigma}(1385)^0 p K^-$		$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{148}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$		$< 8.2 \times 10^{-6}$	CL=90%
Γ_{149}	$\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.}$	[b]	$(3.0 \pm 0.7) \times 10^{-4}$	
Γ_{150}	$\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.}$		$(3.3 \pm 0.8) \times 10^{-4}$	
Γ_{151}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.}$	[b]	$(1.08 \pm 0.06) \times 10^{-3}$	
Γ_{152}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.}$		$(1.25 \pm 0.07) \times 10^{-3}$	
Γ_{153}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$		$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{154}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$		$< 4.1 \times 10^{-6}$	CL=90%
Γ_{155}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$		$< 1.80 \times 10^{-3}$	CL=90%
Γ_{156}	$\Xi^0 \Xi^0$		$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{157}	$\Xi(1530)^- \Xi^+ + \text{c.c.}$		$(3.18 \pm 0.08) \times 10^{-4}$	
Γ_{158}	$\Xi(1530)^0 \Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{159}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	[c]	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{160}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[c]	$< 2.1 \times 10^{-5}$	CL=90%
Γ_{161}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[c]	$< 1.6 \times 10^{-5}$	CL=90%
Γ_{162}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[c]	$< 5.6 \times 10^{-5}$	CL=90%
Γ_{163}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[c]	$< 1.1 \times 10^{-5}$	CL=90%

Decays into stable hadrons

Γ_{164}	$2(\pi^+ \pi^-) \pi^0$		$(4.2 \pm 0.4) \%$	S=2.1
Γ_{165}	$3(\pi^+ \pi^-) \pi^0$		$(2.9 \pm 0.6) \%$	
Γ_{166}	$\pi^+ \pi^- 3\pi^0$		$(1.9 \pm 0.9) \%$	
Γ_{167}	$\rho^\pm \pi^\mp \pi^0 \pi^0$		$(1.41 \pm 0.22) \%$	
Γ_{168}	$\rho^+ \rho^- \pi^0$		$(6.0 \pm 1.1) \times 10^{-3}$	
Γ_{169}	$\pi^+ \pi^- 4\pi^0$		$(6.5 \pm 1.3) \times 10^{-3}$	
Γ_{170}	$\pi^+ \pi^- \pi^0$		$(2.00 \pm 0.07) \%$	S=2.0
Γ_{171}	$2(\pi^+ \pi^- \pi^0)$		$(1.61 \pm 0.20) \%$	
Γ_{172}	$\pi^+ \pi^- \pi^0 K^+ K^-$		$(1.52 \pm 0.27) \%$	S=1.4
Γ_{173}	$\pi^+ \pi^-$		$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{174}	$2(\pi^+ \pi^-)$		$(3.20 \pm 0.25) \times 10^{-3}$	S=1.2
Γ_{175}	$3(\pi^+ \pi^-)$		$(4.3 \pm 0.4) \times 10^{-3}$	

Γ_{176}	$2(\pi^+\pi^-)3\pi^0$	$(6.2 \pm 0.9) \%$	
Γ_{177}	$4(\pi^+\pi^-)\pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{178}	$2(\pi^+\pi^-)\eta$	$(2.29 \pm 0.28) \times 10^{-3}$	
Γ_{179}	$3(\pi^+\pi^-)\eta$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{180}	$2(\pi^+\pi^-\pi^0)\eta$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{181}	$\pi^+\pi^-\pi^0\pi^0\eta$	$(2.4 \pm 0.5) \times 10^{-3}$	
Γ_{182}	$\rho^\pm\pi^\mp\pi^0\eta$	$(1.9 \pm 0.8) \times 10^{-3}$	
Γ_{183}	K^+K^-	$(3.06 \pm 0.05) \times 10^{-4}$	
Γ_{184}	$K_S^0K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
Γ_{185}	$K_S^0K_S^0$	$< 1.4 \times 10^{-8}$	CL=95%
Γ_{186}	$K\bar{K}\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{187}	$K^+K^-\pi^0$	$(2.88 \pm 0.12) \times 10^{-3}$	
Γ_{188}	$K_S^0K^\pm\pi^\mp$	$(5.3 \pm 0.5) \times 10^{-3}$	
Γ_{189}	$K_S^0K_L^0\pi^0$	$(2.06 \pm 0.26) \times 10^{-3}$	
Γ_{190}	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
Γ_{191}	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	
Γ_{192}	$K^+K^-\pi^+\pi^-$	$(7.0 \pm 1.0) \times 10^{-3}$	
Γ_{193}	$K^+K^-\pi^0\pi^0$	$(2.13 \pm 0.22) \times 10^{-3}$	
Γ_{194}	$K^+K^-\pi^0\pi^0\pi^0$	$(1.61 \pm 0.29) \times 10^{-3}$	
Γ_{195}	$K_S^0K^\pm\pi^\mp\pi^0\pi^0$	$(5.3 \pm 0.7) \times 10^{-3}$	
Γ_{196}	$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	$(6.3 \pm 0.4) \times 10^{-3}$	
Γ_{197}	$K_S^0K^\pm\rho(770)^\pm\pi^0$	$(2.9 \pm 0.8) \times 10^{-3}$	
Γ_{198}	$K_S^0K_L^0\pi^+\pi^-$	$(3.8 \pm 0.6) \times 10^{-3}$	
Γ_{199}	$K_S^0K_L^0\pi^0\pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{200}	$K_S^0K_L^0\eta$	$(1.45 \pm 0.33) \times 10^{-3}$	
Γ_{201}	$K_S^0K_S^0\pi^+\pi^-$	$(1.68 \pm 0.19) \times 10^{-3}$	
Γ_{202}	$K^\mp K_S^0\pi^\pm\pi^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
Γ_{203}	$K_S^0K^\pm\pi^\mp\rho(770)^0$	$(3.1 \pm 0.5) \times 10^{-3}$	
Γ_{204}	$K^+K^-2(\pi^+\pi^-)$	$(3.1 \pm 1.3) \times 10^{-3}$	
Γ_{205}	$K^+K^-\pi^+\pi^-\eta$	$(4.7 \pm 0.7) \times 10^{-3}$	
Γ_{206}	$2(K^+K^-)$	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{207}	$K^+K^-K_S^0K_S^0$	$(4.2 \pm 0.7) \times 10^{-4}$	
Γ_{208}	$K_S^0K^*(892)^0\pi^+\pi^-$	$(1.7 \pm 0.6) \times 10^{-3}$	
Γ_{209}	$K_S^0K^*(892)^0\pi^0\pi^0$	$(1.01 \pm 0.18) \times 10^{-3}$	
Γ_{210}	$K^\mp K^*(892)^\pm\pi^+\pi^-$	$(3.4 \pm 1.2) \times 10^{-3}$	
Γ_{211}	$K^*(892)^\pm K^*(892)^0\pi^\mp$	$(4.8 \pm 1.0) \times 10^{-3}$	
Γ_{212}	$K^\mp K^*(892)^\pm\pi^0\pi^0$	$(1.57 \pm 0.32) \times 10^{-3}$	
Γ_{213}	$K^*(892)^+K^*(892)^-\pi^0$	$(1.12 \pm 0.23) \%$	
Γ_{214}	$\rho\bar{\rho}$	$(2.120 \pm 0.029) \times 10^{-3}$	
Γ_{215}	$\rho\bar{\rho}\pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1

Γ_{216}	$p\bar{p}\pi^+\pi^-$		$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{217}	$p\bar{p}\pi^+\pi^-\pi^0$	[d]	$(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{218}	$p\bar{p}\eta$		$(2.00 \pm 0.12) \times 10^{-3}$	
Γ_{219}	$p\bar{p}\rho$		$< 3.1 \times 10^{-4}$	CL=90%
Γ_{220}	$p\bar{p}\omega$		$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{221}	$p\bar{p}\eta'(958)$		$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
Γ_{222}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$		$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{223}	$p\bar{p}\phi$		$(5.19 \pm 0.33) \times 10^{-5}$	
Γ_{224}	$p\bar{n}\pi^-$		$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{225}	$n\bar{n}$		$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{226}	$n\bar{n}\pi^+\pi^-$		$(4 \pm 4) \times 10^{-3}$	
Γ_{227}	$nN(1440)$		seen	
Γ_{228}	$nN(1520)$		seen	
Γ_{229}	$nN(1535)$		seen	
Γ_{230}	$\Lambda\bar{\Lambda}$		$(1.88 \pm 0.08) \times 10^{-3}$	S=2.6
Γ_{231}	$\Lambda\bar{\Lambda}\pi^0$		$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{232}	$\Lambda\bar{\Lambda}\pi^+\pi^-$		$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{233}	$\Lambda\bar{\Lambda}\eta$		$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{234}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	[b]	$(1.26 \pm 0.05) \times 10^{-3}$	S=1.2
Γ_{235}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$		$(1.21 \pm 0.07) \times 10^{-3}$	S=1.8
Γ_{236}	$pK^-\bar{\Lambda} + \text{c.c.}$		$(8.6 \pm 1.1) \times 10^{-4}$	
Γ_{237}	$pK^-\bar{\Sigma}^0$		$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{238}	$pK_S^0\bar{\Sigma}^- + \text{c.c.}$		$(2.73 \pm 0.05) \times 10^{-4}$	
Γ_{239}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$		$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{240}	$\Lambda\bar{\Sigma} + \text{c.c.}$		$(2.83 \pm 0.23) \times 10^{-5}$	
Γ_{241}	$\Sigma^+\bar{\Sigma}^-$		$(1.07 \pm 0.04) \times 10^{-3}$	
Γ_{242}	$\Sigma^0\bar{\Sigma}^0$		$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
Γ_{243}	$\Sigma^+\bar{\Sigma}^-\eta$		$(6.3 \pm 0.4) \times 10^{-5}$	
Γ_{244}	$\Xi^-\bar{\Xi}^+$		$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4

Radiative decays

Γ_{245}	$\gamma\eta_c(1S)$		$(1.41 \pm 0.14) \%$	S=1.3
Γ_{246}	$\gamma\eta_c(1S) \rightarrow 3\gamma$		seen	
Γ_{247}	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$		seen	
Γ_{248}	3γ		$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{249}	4γ		$< 9 \times 10^{-6}$	CL=90%
Γ_{250}	5γ		$< 1.5 \times 10^{-5}$	CL=90%
Γ_{251}	$\gamma\pi^0$		$(3.39 \pm 0.08) \times 10^{-5}$	
Γ_{252}	$\gamma\pi^0\pi^0$		$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{253}	$\gamma 2\pi^+ 2\pi^-$		$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{254}	$\gamma f_2(1270)f_2(1270)$		$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{255}	$\gamma f_2(1270)f_2(1270)$ (non resonant)		$(8.2 \pm 1.9) \times 10^{-4}$	
Γ_{256}	$\gamma\pi^+\pi^-\pi^0$		$(8.3 \pm 3.1) \times 10^{-3}$	

$\Gamma_{257} \gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	
$\Gamma_{258} \gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
$\Gamma_{259} \gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
$\Gamma_{260} \gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
$\Gamma_{261} \gamma \eta$	$(1.090 \pm 0.013) \times 10^{-3}$	
$\Gamma_{262} \gamma \eta \pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	
$\Gamma_{263} \gamma f_0(500) \rightarrow \gamma \pi \pi$		
$\Gamma_{264} \gamma f_0(500) \rightarrow \gamma K \bar{K}$		
$\Gamma_{265} \gamma f_0(500) \rightarrow \gamma \eta \eta$		
$\Gamma_{266} \gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0$	$< 2.5 \times 10^{-6}$	CL=95%
$\Gamma_{267} \gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0$	$< 6.6 \times 10^{-6}$	CL=95%
$\Gamma_{268} \gamma \eta \pi \pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
$\Gamma_{269} \gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	
$\Gamma_{270} \gamma \eta'(958)$	$(5.28 \pm 0.06) \times 10^{-3}$	S=1.3
$\Gamma_{271} \gamma f_0(980) \rightarrow \gamma \pi \pi$		
$\Gamma_{272} \gamma f_0(980) \rightarrow \gamma K \bar{K}$		
$\Gamma_{273} \gamma \rho \rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
$\Gamma_{274} \gamma \rho \omega$	$< 5.4 \times 10^{-4}$	CL=90%
$\Gamma_{275} \gamma \rho \phi$	$< 8.8 \times 10^{-5}$	CL=90%
$\Gamma_{276} \gamma \omega \omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
$\Gamma_{277} \gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
$\Gamma_{278} \gamma \eta(1405/1475) \rightarrow \gamma K \bar{K} \pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
$\Gamma_{279} \gamma \eta(1405/1475) \rightarrow \gamma \gamma \rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
$\Gamma_{280} \gamma \eta(1405/1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
$\Gamma_{281} \gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3
$\Gamma_{282} \gamma \eta(1405/1475) \rightarrow \gamma \gamma \phi$	$< 8.2 \times 10^{-5}$	CL=95%
$\Gamma_{283} \gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^+ \pi^- \pi^0$	$(1.50 \pm 0.16) \times 10^{-5}$	
$\Gamma_{284} \gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^0 \pi^0 \pi^0$	$(7.1 \pm 1.1) \times 10^{-6}$	
$\Gamma_{285} \gamma \eta(1405) \rightarrow \gamma \gamma \gamma$	$< 2.63 \times 10^{-6}$	CL=90%
$\Gamma_{286} \gamma \eta(1475) \rightarrow \gamma \gamma \gamma$	$< 1.86 \times 10^{-6}$	CL=90%
$\Gamma_{287} \gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
$\Gamma_{288} \gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
$\Gamma_{289} \gamma \eta(1760) \rightarrow \gamma \gamma \gamma$	$< 4.80 \times 10^{-6}$	CL=90%
$\Gamma_{290} \gamma \eta(2225)$	$(3.14 \pm_{-0.19}^{+0.50}) \times 10^{-4}$	
$\Gamma_{291} \gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3
$\Gamma_{292} \gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm_{-0.22}^{+0.60}) \times 10^{-5}$	
$\Gamma_{293} \gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	
$\Gamma_{294} \gamma f_0(1370) \rightarrow \gamma \pi \pi$		
$\Gamma_{295} \gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$	
$\Gamma_{296} \gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$	
$\Gamma_{297} \gamma f_0(1370) \rightarrow \gamma \eta \eta$		

Γ_{298}	$\gamma f_0(1370) \rightarrow \gamma \eta \eta'$		
Γ_{299}	$\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi$	$(7.9 \pm 1.3) \times 10^{-4}$	
Γ_{300}	$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	$(1.09 \pm 0.24) \times 10^{-4}$	
Γ_{301}	$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	$(1.7 \pm_{-1.4}^{0.6}) \times 10^{-5}$	
Γ_{302}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm_{-0.60}^{0.24}) \times 10^{-5}$	
Γ_{303}	$\gamma f_0(1500) \rightarrow \gamma \eta \eta'$		
Γ_{304}	$\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	
Γ_{305}	$\gamma f_2'(1525)$	$(5.7 \pm_{-0.5}^{0.8}) \times 10^{-4}$	S=1.5
Γ_{306}	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm_{-0.5}^{0.7}) \times 10^{-5}$	
Γ_{307}	$\gamma f_2'(1525) \rightarrow \gamma \eta \eta$	$(3.4 \pm 1.4) \times 10^{-5}$	
Γ_{308}	$\gamma f_2(1565) \rightarrow \gamma \eta \eta'$		
Γ_{309}	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$	
Γ_{310}	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.8 \pm 0.5) \times 10^{-4}$	
Γ_{311}	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(9.5 \pm_{-0.5}^{1.0}) \times 10^{-4}$	S=1.5
Γ_{312}	$\gamma f_0(1710) \rightarrow \gamma \omega \omega$	$(3.1 \pm 1.0) \times 10^{-4}$	
Γ_{313}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$(2.4 \pm_{-0.7}^{1.2}) \times 10^{-4}$	
Γ_{314}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta'$		
Γ_{315}	$\gamma f_0(1710) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$	
Γ_{316}	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm_{-0.33}^{0.20}) \times 10^{-5}$	
Γ_{317}	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \pm_{-2.4}^{3.5}) \times 10^{-5}$	
Γ_{318}	$\gamma \eta_1(1855) \rightarrow \gamma \eta \eta'$	$(2.7 \pm_{-0.5}^{0.4}) \times 10^{-6}$	
Γ_{319}	$\gamma f_0(1770) \rightarrow \gamma \eta \eta'$		
Γ_{320}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$	
Γ_{321}	$\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$	
Γ_{322}	$\gamma f_2(2010) \rightarrow \gamma \eta \eta'$		
Γ_{323}	$\gamma f_0(2020) \rightarrow \gamma \pi \pi$		
Γ_{324}	$\gamma f_0(2020) \rightarrow \gamma K \bar{K}$		
Γ_{325}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta$		
Γ_{326}	$\gamma f_0(2020) \rightarrow \gamma \eta' \eta'$	$(2.63 \pm_{-0.50}^{0.32}) \times 10^{-4}$	
Γ_{327}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta'$		
Γ_{328}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{329}	$\gamma f_4(2050) \rightarrow \gamma \eta \eta'$		
Γ_{330}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \pm_{-0.30}^{0.60}) \times 10^{-4}$	
Γ_{331}	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$		
Γ_{332}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$	
Γ_{333}	$\gamma f_0(2200)$	seen	

Γ_{334}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$	
Γ_{335}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm_{-0.50}^{+0.19}) \times 10^{-4}$	
Γ_{336}	$\gamma f_0(2200) \rightarrow \gamma \pi \pi$		
Γ_{337}	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$		
Γ_{338}	$\gamma f_J(2220)$	seen	
Γ_{339}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	< 3.9	$\times 10^{-5}$ CL=90%
Γ_{340}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	< 4.1	$\times 10^{-5}$ CL=90%
Γ_{341}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$	
Γ_{342}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$	
Γ_{343}	$\gamma f_0(2330) \rightarrow \gamma \pi \pi$		
Γ_{344}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta$		
Γ_{345}	$\gamma f_0(2330) \rightarrow \gamma \eta' \eta'$	$(6.1 \pm_{-1.8}^{+4.0}) \times 10^{-6}$	
Γ_{346}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta'$		
Γ_{347}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \pm_{-2.2}^{+2.4}) \times 10^{-5}$	
Γ_{348}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm_{-1.5}^{+4.0}) \times 10^{-5}$	
Γ_{349}	$\gamma f_2(2340) \rightarrow \gamma \eta' \eta'$	$(8.7 \pm_{-1.8}^{+0.9}) \times 10^{-6}$	
Γ_{350}	$\gamma f_0(2470) \rightarrow \gamma \eta' \eta'$	$(8.2 \pm_{-2.8}^{+4.0}) \times 10^{-7}$	
Γ_{351}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.7 \pm_{-0.8}^{+0.6}) \times 10^{-4}$	S=1.6
Γ_{352}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \pm_{-0.9}^{+1.5}) \times 10^{-5}$	
Γ_{353}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm_{-1.3}^{+2.0}) \times 10^{-5}$	
Γ_{354}	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$		
Γ_{355}	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	< 3.56	$\times 10^{-6}$ CL=90%
Γ_{356}	$\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \pm_{-0.8}^{+0.7}) \times 10^{-5}$	
Γ_{357}	$\gamma \eta(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$	
Γ_{358}	$\gamma \eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$	
Γ_{359}	$\gamma \eta(2370) \rightarrow \gamma \eta \eta \eta'$	< 9.2	$\times 10^{-6}$ CL=90%
Γ_{360}	$\gamma D^0 + \text{c.c.}$	< 9.1	$\times 10^{-8}$ CL=90%
Γ_{361}	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{362}	$\gamma p \bar{p} \pi^+ \pi^-$	< 7.9	$\times 10^{-4}$ CL=90%
Γ_{363}	$\gamma \Lambda \bar{\Lambda}$	< 1.3	$\times 10^{-4}$ CL=90%
Γ_{364}	$\gamma A^0 \rightarrow \gamma \text{invisible}$	[e] < 1.7	$\times 10^{-6}$ CL=90%
Γ_{365}	$\gamma A^0 \rightarrow \gamma \gamma \gamma$	< 4.9	$\times 10^{-7}$ CL=95%
Γ_{366}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] < 7.8	$\times 10^{-7}$ CL=90%

Dalitz decays

Γ_{367}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$
Γ_{368}	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$
Γ_{369}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$

Γ_{370}	$\eta(1405)e^+e^- \rightarrow$ $f_0(980)\pi^0 e^+e^- \rightarrow$ $\pi^+\pi^-\pi^0 e^+e^-$	$(2.04 \pm 0.22) \times 10^{-7}$	
Γ_{371}	$X(1835)e^+e^-, X \rightarrow \pi^+\pi^-\eta'$	$(3.58 \pm 0.25) \times 10^{-6}$	
Γ_{372}	$X(2120)e^+e^-, X \rightarrow \pi^+\pi^-\eta'$	$(8.2 \pm 1.3) \times 10^{-7}$	
Γ_{373}	$\eta(2370)e^+e^-, \eta \rightarrow \pi^+\pi^-\eta'$	$(1.08 \pm 0.17) \times 10^{-6}$	
Γ_{374}	$\eta U \rightarrow \eta e^+e^-$	$[g] < 9.11$	$\times 10^{-7}$ CL=90%
Γ_{375}	$\eta'(958)U \rightarrow \eta'(958)e^+e^-$	$[g] < 2.0$	$\times 10^{-7}$ CL=90%
Γ_{376}	ϕe^+e^-	< 1.2	$\times 10^{-7}$ CL=90%

Weak decays

Γ_{377}	$D^- e^+ \nu_e + \text{c.c.}$	< 7.1	$\times 10^{-8}$ CL=90%
Γ_{378}	$D^- \mu^+ \nu_\mu + \text{c.c.}$	< 5.6	$\times 10^{-7}$ CL=90%
Γ_{379}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	< 8.5	$\times 10^{-8}$ CL=90%
Γ_{380}	$D_s^- e^+ \nu_e + \text{c.c.}$	< 1.3	$\times 10^{-6}$ CL=90%
Γ_{381}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	< 1.8	$\times 10^{-6}$ CL=90%
Γ_{382}	$D^- \pi^+ + \text{c.c.}$	< 7.0	$\times 10^{-8}$ CL=90%
Γ_{383}	$D^- \rho^+ + \text{c.c.}$	< 6.0	$\times 10^{-7}$ CL=90%
Γ_{384}	$\bar{D}^0 \pi^0 + \text{c.c.}$	< 4.7	$\times 10^{-7}$ CL=90%
Γ_{385}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$ CL=90%
Γ_{386}	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$ CL=90%
Γ_{387}	$\bar{D}^0 \eta + \text{c.c.}$	< 6.8	$\times 10^{-7}$ CL=90%
Γ_{388}	$\bar{D}^0 \rho^0 + \text{c.c.}$	< 5.2	$\times 10^{-7}$ CL=90%
Γ_{389}	$D_s^- \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$ CL=90%
Γ_{390}	$D_s^- \rho^+ + \text{c.c.}$	< 1.3	$\times 10^{-5}$ CL=90%

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

Γ_{391}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$ CL=90%
Γ_{392}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$ CL=90%
Γ_{393}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$ CL=90%
Γ_{394}	$e^\pm \tau^\mp$	LF	< 7.5	$\times 10^{-8}$ CL=90%
Γ_{395}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$ CL=90%
Γ_{396}	$\Lambda_c^+ e^- + \text{c.c.}$		< 6.9	$\times 10^{-8}$ CL=90%

Other decays

Γ_{397}	invisible	< 7	$\times 10^{-4}$ CL=90%
Γ_{398}	$\mu^+ \mu^- X^0 \rightarrow \mu^+ \mu^- + \text{invisible}$		

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] $\Theta(1540)$ is a hypothetical pentaquark state of $1.54 \text{ GeV}/c^2$ mass and a width of less than $25 \text{ MeV}/c^2$.

- [d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.
[e] For a narrow state A with mass less than 960 MeV.
[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.
[g] For a dark photon U with mass between 100 and 2100 MeV.

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and $B^\pm$ with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

$J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$81.37 \pm 1.36 \pm 1.30$	¹ ANASHIN 20	KEDR	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
74.1 ± 8.1	BAI	95B	BES e^+e^-
59 ± 24	BALDINI-...	75	FRAG e^+e^-
59 ± 14	BOYARSKI	75	MRK1 e^+e^-
50 ± 25	ESPOSITO	75B	FRAM e^+e^-

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there.

$\Gamma(e^+e^-)$

Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.53 ± 0.10	OUR AVERAGE			
$5.550 \pm 0.056 \pm 0.089$		^{1,2} ANASHIN 18A	KEDR	e^+e^-
$5.36^{+0.29}_{-0.28}$		³ HSUEH 92	RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.58 \pm 0.05 \pm 0.08$		⁴ ABLIKIM 16Q	BES3	$3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71 ± 0.16	13k	⁵ ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ± 0.19	7.8k	⁵ AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ± 0.39		BAI	95B	BES e^+e^-
4.72 ± 0.35		ALEXANDER 89	RVUE	See Υ mini-review
4.4 ± 0.6		³ BRANDELIK 79C	DASP	e^+e^-
4.6 ± 0.8		⁶ BALDINI-... 75	FRAG	e^+e^-
4.8 ± 0.6		BOYARSKI 75	MRK1	e^+e^-
4.6 ± 1.0		ESPOSITO 75B	FRAM	e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+\mu^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.13 ± 0.52	BAI	95B	BES e^+e^-
4.8 ± 0.6	BOYARSKI	75	MRK1 e^+e^-
5 ± 1	ESPOSITO	75B	FRAM e^+e^-

$\Gamma(\gamma\gamma)$ Γ_{391}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP e^+e^-

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(I) in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$4.884 \pm 0.048 \pm 0.078$	^{1,2} ANASHIN	18A	KEDR e^+e^-
4 ± 0.8	³ BALDINI-...	75	FRAG e^+e^-
3.9 ± 0.8	³ ESPOSITO	75B	FRAM e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$333.1 \pm 6.6 \pm 4.0$	^{1,2} ANASHIN	18A	KEDR e^+e^-
$332.3 \pm 6.4 \pm 4.8$	ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow e^+e^-$
350 ± 20	BRANDELIK	79C	DASP e^+e^-
320 ± 70	³ BALDINI-...	75	FRAG e^+e^-
340 ± 90	³ ESPOSITO	75B	FRAM e^+e^-
360 ± 100	³ FORD	75	SPEC e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 ± 4 OUR AVERAGE				
$333.4 \pm 2.5 \pm 4.4$		ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
$331.8 \pm 5.2 \pm 6.3$		ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow \mu^+\mu^-$
$338.4 \pm 5.8 \pm 7.1$	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$330.1 \pm 7.7 \pm 7.3$	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90	DASP	75	DASP	e^+e^-
380 ± 50	¹ ESPOSITO	75B	FRAM	e^+e^-

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 OUR AVERAGE				
2.34 ± 0.43 ± 0.16	49	LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$
2.22 ± 0.96 ± 0.02	9	¹ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
64.8 ± 11.1 ± 0.4	200	¹ LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.9 ± 5.7 ± 0.1	101	¹ LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-\pi^0\gamma\gamma)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
4.76 ± 1.64 ± 0.03	¹ LEES	23	BABR $e^+e^- \rightarrow \gamma_{ISR}$ hadrons

¹ LEES 23 reports $[\Gamma(J/\psi(1S) \rightarrow \eta K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 1.55 \pm 0.51 \pm 0.16$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 ± 1.4 ± 0.4	44	LEES	17D	BABR $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
155 ± 26 ± 36	14k	LEES	21	BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{27}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
32±13±15	14k	LEES 21	BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$

$$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{29}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4±1.0±1.9	130	LEES 17D	BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{44}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	¹ AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	¹ LEES 18E	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega 3\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{46}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±3.1±0.1	89	¹ LEES 21C	BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{48}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
16.9±7.6±0.2	¹ LEES 21C	BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

¹ Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best values $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$$\Gamma(\omega \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{49}\Gamma_5/\Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.3±0.2	170	AUBERT 06D	BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \pi^0 \gamma$

$$\Gamma(\omega \pi^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{50}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	¹ LEES 18E	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$

¹LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{52}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
185±30±1	14k	¹ LEES	21 BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

¹LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 165 \pm 9 \pm 25 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{64}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70±1.98±0.03	24	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega K^+ K^- \gamma$

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 3.3 \pm 1.3 \pm 1.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{73}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
29.0±1.7±1.3	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+ K^*(892)^- \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{74}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.96±0.85±0.70	155	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+ K^- \pi^0 \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{75}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.76±1.70±1.00	89	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{76}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
26.6±2.5±1.5	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{77}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17.70±1.70±1.00	94	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{78}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42.6±4.8±7.2	99	¹ LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0)=1/6$.

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{79}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8±2.8±6.8	80	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{80}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0±2.8 OUR AVERAGE				
9.2±1.2±3.2	64	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	² LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{81}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±1.2±0.3	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{84}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60±0.75±2.25	34	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{86}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28±0.34±0.07	47±12	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28±0.40±0.11 25±8 ^{1,2} AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by (2/3)² to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K \pi)$.

² Superseded by LEES 12F.

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{87}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80±0.48±0.32	1±5	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by (1/4)² to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{98}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1±9.8±0.5	35	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{99} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

25.8 ± 1.4 ± 0.6	710	1,2,3 LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
-------------------------	-----	------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1	317	2,4 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------	-----	------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{100} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

18.6 ± 16.1 ± 0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
--------------------------	-------	----------	----------	--

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{101} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.32 ± 2.00 ± 0.08	8 ± 8	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
---------------------------	-------	-------------------	----------	--

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{103} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.8 ± 0.4 ± 0.3	110 ± 14	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------------------	----------	---------------------	-----------	---

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{110} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.41 ± 0.34 OUR AVERAGE

4.39 ± 0.48 ± 0.05	181	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------	-----	-------------------	----------	---

4.44 ± 0.48 ± 0.05	254 ± 23	² SHEN	09 BELL	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------	----------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 5.2 $\pm 0.7 \pm 0.1$ 103 ³ AUBERT,BE 06D BABR 10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
- ¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{111}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.73$\pm 0.56 \pm 0.03$	45	¹ LEES 12F	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 3.09 $\pm 0.86 \pm 0.03$ 23 ² AUBERT,BE 06D BABR 10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
- ¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{112}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.94$\pm 0.19 \pm 0.01$	35	¹ AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow \phi 2(\pi^+\pi^-)\gamma$

- ¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{113}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.6± 1.4 OUR AVERAGE				

- 4.1 $\pm 1.6 \pm 0.4$ ¹ LEES 23 BABR $e^+e^- \rightarrow \gamma_{ISR}$ hadrons
- 6.1 $\pm 2.7 \pm 0.4$ 6 ² AUBERT 07AU BABR 10.6 $e^+e^- \rightarrow \phi\eta\gamma$
- ¹ LEES 23 quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi^0) = 0.64 \pm 0.26 \pm 0.06$ eV.
- ² AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{117} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.42 ± 0.18 OUR AVERAGE

1.38 ± 0.24 ± 0.02	57 ± 9	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
--------------------	--------	-------------------	----------	---

1.48 ± 0.27 ± 0.09	60 ± 11	² SHEN	09 BELL	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------	---------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.00 ± 0.23 ± 0.01	20 ± 5	³ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
--------------------	--------	---------------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Multiplied by 2/3 to take into account the $\phi \pi^+ \pi^-$ mode only. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{118} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.96 ± 0.26 ± 0.01	16 ± 4	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
---------------------------	--------	-------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94 ± 0.39 ± 0.01	7.0 ± 2.8	² AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
--------------------	-----------	---------------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{126} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.79 ± 0.32^{+0.02}_{-0.06}	61	^{1,2,3} LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
--	----	-----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.08 ± 0.73 ^{+0.05} _{-0.14}	44	^{2,4} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
---	----	-----------------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28 \text{ eV}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{131}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.1±3.2±0.2	11	1,2 LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

²LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3 \text{ eV}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{137}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.53±0.61±0.05	163	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow K^+ K^- K^+ K^- \gamma$

¹LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{138}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.21±0.83±0.03	29	¹ LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f'_2(1525) K^+ K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{143}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.

²LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525) K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20 \text{ eV}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{164}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0 \gamma$

$$\Gamma(\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{166}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
100 ± 50	OUR AVERAGE	Error includes scale factor of 4.3.		
55 ± 16 ± 1	14k	¹ LEES	21	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ
¹ LEES 21 reports [Γ(J/ψ(1S) → π ⁺ π ⁻ 3π ⁰) × Γ(J/ψ(1S) → e ⁺ e ⁻)/Γ _{total}] × [Γ(ψ(2S) → J/ψ(1S)π ⁺ π ⁻)/Γ _{total}] = 19.2 ± 4.5 ± 3.2 eV which we divide by our best value Γ(ψ(2S) → J/ψ(1S)π ⁺ π ⁻)/Γ _{total} = 0.3469 ± 0.0034. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$$\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{169}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8 ± 4.4 ± 5.4	340	LEES	21C	BABR e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 4π ⁰)

$$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{167}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0 ± 9.0 ± 8.0	1.2k	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ

$$\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{168}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0 ± 5.0 ± 3.3	529	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{170}\Gamma_5/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.1248 ± 0.0019 ± 0.0026	LEES	21B	BABR 10.5 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.122 ± 0.005 ± 0.008	AUBERT,B	04N	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{171}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9 ± 0.5 ± 1.0	761	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰)γ

$$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{172}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0 ± 4.3 ± 6.4	768	AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{174}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4 ± 0.9 ± 0.4		LEES	12E	BABR 10.6 e ⁺ e ⁻ → 2π ⁺ 2π ⁻ γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
19.5 ± 1.4 ± 1.3	270	¹ AUBERT	05D	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)γ
¹ Superseded by LEES 12E.				

$$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{175}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.16 ± 0.14	496	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → 3(π ⁺ π ⁻)γ

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{176}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
345±10±50	14k	LEES 21	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{178}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{180}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES 21	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{181}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3	14k	¹ LEES 21	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
12.8± 1.8±2.0	203	LEES 18E	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 6 \pm 4 \pm 1$ eV which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{182}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES 18E	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{183}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78±0.11±0.05	462	¹ LEES 15J	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
1.94±0.11±0.05	462	² LEES 15J	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
1.42±0.23±0.08	51	³ LEES 13Q	BABR	$e^+e^- \rightarrow K^+K^-\gamma$

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{189}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES 17A	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0\gamma$

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{190}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.9±0.4	106	LEES 17A	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0\gamma$

$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{191} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.4 \pm 0.7 \pm 0.1$	37	LEES	17A	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{192} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$37.94 \pm 0.81 \pm 1.10$	3.1k	LEES	12F	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$36.3 \pm 1.3 \pm 2.1$	1.5k	¹ AUBERT	07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$33.6 \pm 2.7 \pm 2.7$	233	² AUBERT	05D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹Superseded by LEES 12F.²Superseded by AUBERT 07AK.

$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{193} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$11.75 \pm 0.81 \pm 0.90$	388	LEES	12F	BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$13.6 \pm 1.1 \pm 1.3$	203	¹ AUBERT	07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

¹Superseded by LEES 12F.

$\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{194} \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
$8.9 \pm 1.3 \pm 0.9$	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{195} \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
$29.3 \pm 2.6 \pm 2.9$	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{196} \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
$34.6 \pm 1.4 \pm 1.8$	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K_S^0 K^\pm \rho(770)^\pm \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{197} \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
$16.0 \pm 4.1 \pm 1.6$	LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{198} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$20.8 \pm 2.3 \pm 2.1$	248	LEES	14H	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$

$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{199} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$10.3 \pm 2.3 \pm 0.5$	47	LEES	17A	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{200} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$8.0 \pm 1.8 \pm 0.4$	45	LEES	17A	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

$$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{201} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±0.9±0.5	133	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{202} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7±1.9±1.8	393	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{203} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
17.3±2.1±1.7	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{204} \Gamma_5 / \Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75±0.23±0.17	205	AUBERT	06D	BABR $10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

$$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{205} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{206} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00±0.33±0.29	287 ± 24	LEES	12F	BABR $10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.11±0.39±0.30	156 ± 15	¹ AUBERT	07AK	BABR $10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D	BABR $10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{207} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.4±0.1	29	LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

$$\Gamma(K_S^0 K^*(892)^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{208} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
9.45±3.15±0.90	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K_S^0 K^*(892)^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{209} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
5.59±0.79±0.55	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K^\mp K^*(892)^\pm \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{210}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
18.6±6.3±1.8	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K^*(892)^\pm K^*(892)^0 \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{211}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
26.6±4.5±2.7	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{212}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
8.67±1.56±0.84	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$


$$\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{213}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
62.1±10.8±6.30	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{214}\Gamma_5/\Gamma$$

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
11.3±0.4±0.3	821	1 LEES	13O	BABR $e^+ e^- \rightarrow p\bar{p}\gamma$
12.9±0.4±0.4	918	2 LEES	13Y	BABR $e^+ e^- \rightarrow p\bar{p}\gamma$
9.7±1.7		3 ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
12.0±0.6±0.5	438	4 AUBERT	06B	BABR $e^+ e^- \rightarrow p\bar{p}\gamma$

WEIGHTED AVERAGE
11.9±0.6 (Error scaled by 1.8)



¹ISR photon reconstructed in the detector²ISR photon undetected³Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.⁴Superseded by LEES 130 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{230}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7±0.9±0.7	AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

 $\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{241}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
6.8±1.5±0.8	GONG	23 BELL	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

 $\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{242}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.2±1.5±0.6	GONG	23 BELL	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
6.4±1.2±0.6	AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

 $J/\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ above.

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.877±0.005 OUR AVERAGE			
0.878±0.005	BAI	95B BES	e^+e^-
0.86 ±0.02	BOYARSKI	75 MRK1	e^+e^-

 $\Gamma(\text{virtual}\gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.1346±0.0007	¹ LIAO	23 RVUE	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.135 ±0.003	^{2,3} SETH	04 RVUE	e^+e^-
0.17 ±0.02	² BOYARSKI	75 MRK1	e^+e^-

¹Using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (5.967 \pm 0.023)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

²Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

³Using $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

 $\Gamma(ggg)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
64.1±1.0	6 M	¹ BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$

¹Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+\ell^-)$, $B(\text{virtual}\gamma \rightarrow \text{hadrons})$, and $B(\gamma\eta_c)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.79$\pm$1.05	200 k	¹ BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
13.7$\pm$0.1$\pm$0.7	6 M	BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.971$\pm$0.032 OUR AVERAGE				
5.983 $\pm$ 0.007 $\pm$ 0.037	720k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945 $\pm$ 0.067 $\pm$ 0.042	15k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 $\pm$ 0.05 $\pm$ 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 $\pm$ 0.33		BAI	95B BES	$e^+ e^-$
5.92 $\pm$ 0.15 $\pm$ 0.20		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 $\pm$ 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

 $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.8$\pm$1.3$\pm$0.4	¹ ARMSTRONG	96 E760	$\bar{p} p \rightarrow e^+ e^- \gamma$

¹ For $E_\gamma > 100$ MeV.

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.961$\pm$0.033 OUR AVERAGE				
5.973 $\pm$ 0.007 $\pm$ 0.038	770k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960 $\pm$ 0.065 $\pm$ 0.050	17k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 $\pm$ 0.06 $\pm$ 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 $\pm$ 0.33		BAI	95B BES	$e^+ e^-$
5.90 $\pm$ 0.15 $\pm$ 0.19		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 $\pm$ 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

 $\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$ Γ_5/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
1.0016$\pm$0.0031 OUR AVERAGE			
1.0022 $\pm$ 0.0044 $\pm$ 0.0048	¹ AULCHENKO	14 KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017 $\pm$ 0.0017 $\pm$ 0.0033	² ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 $\pm$ 0.021 $\pm$ 0.013	³ ANASHIN	10 KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 $\pm$ 0.012 $\pm$ 0.006	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.011 $\pm$ 0.013 $\pm$ 0.016	BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 $\pm$ 0.07	BAI	95B BES	$e^+ e^-$
1.00 $\pm$ 0.05	BOYARSKI	75 MRK1	$e^+ e^-$

0.91 ± 0.15	ESPOSITO	75B	FRAM	e^+e^-
0.93 ± 0.10	FORD	75	SPEC	e^+e^-

¹ From 235.3k $J/\psi \rightarrow e^+e^-$ and 156.6k $J/\psi \rightarrow \mu^+\mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$.

$\Gamma(e^+e^-e^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.48$\pm 0.31\pm 0.45$	700	¹ ABLIKIM	24L	BES3 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

¹ $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

$\Gamma(e^+e^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53$\pm 0.22\pm 0.13$	354	¹ ABLIKIM	24L	BES3 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

¹ $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

$\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.3 $\pm 1.1\pm 0.1$	452	¹ AAIJ	24AE	LHCB	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$
<16	90	3.4	² ABLIKIM	24L	BES3 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
10.1 $^{+3.3}_{-2.8}\pm 0.1$	12	³ HAYRAPETY...24A	CMS		$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$

¹ AAIJ 24AE reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (1.89 \pm 0.17 \pm 0.09) \times 10^{-5}$ which we multiply by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Measured with $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$, $J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$. $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

³ HAYRAPETYAN 24A reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (16.9^{+5.5}_{-4.6} \pm 0.6) \times 10^{-6}$ which we multiply by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma(\mu^+\mu^-)$ Γ_{10}/Γ_7

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
18.7± 1.8 OUR AVERAGE				


18.9 $\pm 1.7\pm 0.9$	452	¹ AAIJ	24AE	LHCB	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$
16.9 $^{+5.5}_{-4.6}\pm 0.6$	12	HAYRAPETY...24A	CMS		$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$

¹ Includes prompt production and inclusive decays of b -hadrons.

HADRONIC DECAYS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$			Γ_{11}/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.88 ±0.12	OUR AVERAGE	Error includes scale factor of 2.6. See the ideogram below.			
2.072±0.017±0.062	19.8k	¹ ANASHIN	23 KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
2.18 ±0.19		^{2,3} AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$	
2.184±0.005±0.201	220k	^{3,4} BAI	04H BES	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
2.091±0.021±0.116		^{3,5} BAI	04H BES	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	
1.21 ±0.20		BAI	96D BES	$e^+e^- \rightarrow \rho\pi$	
1.42 ±0.01 ±0.19		COFFMAN	88 MRK3	e^+e^-	
1.3 ±0.3	150	FRANKLIN	83 MRK2	e^+e^-	
1.6 ±0.4	183	ALEXANDER	78 PLUT	e^+e^-	
1.33 ±0.21		BRANDELIK	78B DASP	e^+e^-	
1.0 ±0.2	543	BARTEL	76 CNTR	e^+e^-	
1.3 ±0.3	153	JEAN-MARIE	76 MRK1	e^+e^-	

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.
² From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).
³ Not independent of their $B(\pi^+\pi^-\pi^0)$.
⁴ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.
⁵ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{11}/Γ_{170}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.142±0.011±0.026	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.331±0.033	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.² From a Dalitz plot analysis in a Veneziano model.
 $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$ Γ_{12}/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
0.328±0.005±0.027	COFFMAN	88	MRK3 e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ±0.08	ALEXANDER	78	PLUT e^+e^-
0.32 ±0.08	BRANDELIK	78B	DASP e^+e^-
0.39 ±0.11	BARTEL	76	CNTR e^+e^-
0.37 ±0.09	JEAN-MARIE	76	MRK1 e^+e^-

 $\Gamma(a_2(1320)^0\pi^+\pi^-\rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
2.84±0.08±0.60	1317	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

 $\Gamma(a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.67±0.09±0.73	1628	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

 $\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.2 OUR AVERAGE				
11.7±0.7±2.5	7584	AUGUSTIN	89	DM2 $J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
8.4±4.5	36	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.68	471	¹ ABLIKIM	19Q	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$

¹ From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$.
 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^+\pi^-\eta$

 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
1.20±0.14±0.37	471	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C	BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

$$\Gamma(\eta K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}} \quad \Gamma_{23}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$21.8 \pm 2.2 \pm 3.4$	232 ± 23	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$

$$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{24}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM	10C	BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

$$\Gamma(\rho\eta'(958)) / \Gamma_{\text{total}} \quad \Gamma_{25}/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.8 OUR AVERAGE	Error	includes scale factor of 1.6.		
$7.90 \pm 0.19 \pm 0.49$	3476	¹ ABLIKIM	17AK	BES3 $J/\psi \rightarrow \pi^+ \pi^- \eta'$
$8.3 \pm 3.0 \pm 1.2$	19	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
$11.4 \pm 1.4 \pm 1.6$		COFFMAN	88	MRK3 $J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

$$\Gamma(\rho^+ K^+ K^- \pi^- + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{28}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.53 \pm 0.16 \pm 0.81$	485	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$$\Gamma(h_1(1415)\eta' \rightarrow \gamma\eta\eta') / \Gamma_{\text{total}} \quad \Gamma_{30}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$0.08 \pm 0.01^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	------	---

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ *P*-wave.

$$\Gamma(h_1(1595)\eta' \rightarrow \gamma\eta\eta') / \Gamma_{\text{total}} \quad \Gamma_{31}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$0.16 \pm 0.02^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	------	---

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ *P*-wave.

$$\Gamma(\rho(1450)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.2 \pm 0.2 \pm 1.1$	19.8k	¹ ANASHIN	23	KEDR $e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0 \pi^0$, $J/\psi \rightarrow \rho^+ \pi^-$, $J/\psi \rightarrow \rho^- \pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.

$$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{33}/\Gamma_{170}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.9 \pm 1.7 \pm 2.7$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
0.80 ± 0.27	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{34}/\Gamma_{188}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$6.3 \pm 0.8 \pm 0.6$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
---	----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0)/\Gamma(K^+ K^- \pi^0) \quad \Gamma_{35}/\Gamma_{187}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$9.3 \pm 2.0 \pm 0.6$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$
---	----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}} \quad \Gamma_{36}/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.28 \pm 0.55 \pm 0.44$	119	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
--	-----	----------------------	-----------	--------------------------------------

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

$$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{38}/\Gamma_{170}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$8 \pm 2 \pm 5$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
-----------------------------------	-----	-------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 6	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{40}/\Gamma_{170}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4 \pm 1 \pm 20$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
------------------------------------	-----	-------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

600 ± 250	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{41}/\Gamma_{170}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0 ± 0.8	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in a Veneziano model.


$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

Γ_{42}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
$0.538 \pm 0.012 \pm 0.065$	2090	¹ ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\pi^0$
$0.360 \pm 0.028 \pm 0.054$	222	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
$0.482 \pm 0.019 \pm 0.064$		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

WEIGHTED AVERAGE
 0.45 ± 0.05 (Error scaled by 1.4)



$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_{43}/Γ_{170}

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8 \pm 3 \pm 2$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$


¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 1.0 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$10.6 \pm 1.2 \pm 0.1$	3531	¹ ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0 ± 1.6	18058	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8 ± 1.6	215	BURMESTER	77D	PLUT e^+e^-
6.8 ± 1.9	348	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.




$\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_{45}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.4 \pm 0.3 \pm 0.7$	509	AUGUSTIN	89	DM2	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					Γ_{47}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.3 ± 0.6 OUR AVERAGE					
$4.3 \pm 0.2 \pm 0.6$	5860	AUGUSTIN	89	DM2	e^+e^-
4.0 ± 1.6	70	BURMESTER	77D	PLUT	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.9 ± 0.8	81	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{48}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.74 ± 0.20 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
2.352 ± 0.273	5k	¹ ABLIKIM	06F	BES2	$J/\psi \rightarrow \omega\eta$
$1.44 \pm 0.40 \pm 0.14$	13	² AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
$1.43 \pm 0.10 \pm 0.21$	378	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
$1.71 \pm 0.08 \pm 0.20$		COFFMAN	88	MRK3	$e^+e^- \rightarrow 3\pi\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.
² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

WEIGHTED AVERAGE
 1.74 ± 0.20 (Error scaled by 1.6)



$\Gamma(\omega\pi^+\pi^-\pi^-\pi^-)/\Gamma_{\text{total}}$

Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85 ± 34	140	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{53}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.12 \pm 0.02 \pm 0.13$	14k	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$

Γ_{54}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.18 OUR AVERAGE				
$2.08 \pm 0.30 \pm 0.14$	137	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26 ± 0.43	218	² ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
$1.8^{+1.0}_{-0.8} \pm 0.3$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$1.66 \pm 0.17 \pm 0.19$		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

² Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$

Γ_{55}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.41 \pm 0.27 \pm 0.47$	¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
$4.8 \pm 1.1 \pm 0.3$	1,2	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios. $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.8^{+1.9}_{-1.6} \pm 1.7$	111^{+31}_{-26}	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.2 \times 10^{-4}$	90	¹ VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 2.8 \times 10^{-4}$	90	¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$. $\Gamma(\omega X(1835) \rightarrow \omega p \bar{p})/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-6}$	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma \pi^0 p \bar{p}$

 $\Gamma(\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$< 6.2 \times 10^{-5}$	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$

¹ Using the decays $\omega \rightarrow \pi^+ \pi^- \pi^0$ and $\eta' \rightarrow \eta \pi^+ \pi^-$. $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.52 \pm 0.30 \pm 0.01$	276	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.33 \pm 0.02 \pm 0.12$	ABLIKIM	24BQ BES3	$e^+ e^- \rightarrow J/\psi(1S)$

 $\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
$37.7 \pm 0.8 \pm 5.8$	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
$29.5 \pm 1.4 \pm 7.0$	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4 OUR AVERAGE				
19.8 ± 2.1 ± 3.9		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	e^+e^-

¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.13 OUR AVERAGE			
1.50 ± 0.02 ± 0.19	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
1.47 ± 0.03 ± 0.17	² ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

² From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.66 ± 0.03 ± 0.21	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.16 ± 0.12 ± 0.29	1.1k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.51 ± 0.09 ± 0.21	1.0k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

$\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
4.69 ± 0.80 ^{+0.74}_{-1.82}	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{72} / \Gamma_{188}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
90.5 ± 0.9 ± 3.8	4k	¹ LEES	17c BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{73} / Γ


VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ^{+0.8}_{-1.0} OUR AVERAGE		Error includes scale factor of 2.9. See the ideogram below.		

8.07 ± 0.04 ^{+0.38} _{-0.61}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

WEIGHTED AVERAGE
6.0+0.8-1.0 (Error scaled by 2.9)



$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}}$ Γ_{74} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
2.69 ± 0.01 ^{+0.13}_{-0.20}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0) \quad \Gamma_{74} / \Gamma_{187}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
92.4 ± 1.5 ± 3.4	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{76} / \Gamma$$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88	MRK3 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6	45	VANNUCCI	77	MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
-----------	----	----------	----	---

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{78} / \Gamma$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
5.73 ± 0.14 ± 0.82	¹ ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ² ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

¹ Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value 2/3 for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

² A $K_S^0(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{82} / \Gamma$$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.81 ± 0.10 ± 0.54	1559	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$$\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{83} / \Gamma$$

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
6.28^{+0.16+0.59}_{-0.17-0.52}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$$\Gamma(K^*(892)^\pm K^*(700)^\mp) / \Gamma_{\text{total}} \quad \Gamma_{85} / \Gamma$$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.18^{+0.94}_{-0.54}	655	ABLIKIM	10E	BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{86} / \Gamma$$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5 90 VANNUCCI 77 MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) / \Gamma_{\text{total}} \quad \Gamma_{87} / \Gamma$$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.19^{+0.11}_{-0.32}	323	ABLIKIM	10E	BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT
$3.8 \pm 0.8 \pm 1.2$	¹	BAI	99C	BES $e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

 $\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{90}/Γ_{187}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 1.1 \pm 0.7$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{91}/Γ_{188}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{93}/Γ_{187}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{94}/Γ_{188}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(\bar{K}_2^*(1430)K + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 40 \times 10^{-4}$	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 66 \times 10^{-4}$	90	BRAUNSCH...	76	DASP $e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
-----------------------	----	-------------	----	---

 $\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.04^{+0.25}_{-0.19}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.65 \pm 0.80 \pm 0.44$	1094	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ± 2.6	40	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
---------------	----	----------	----	--

$$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}} \quad \Gamma_{102}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI	77	MRK1 $e^+e^- \rightarrow \pi^+\pi^- K^+K^-$

$$\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{104}/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{105}/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}} \quad \Gamma_{106}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	¹ BAI	99C	BES e^+e^-

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$$\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{107}/\Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$$\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}} \quad \Gamma_{108}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43 \times 10^{-4}$	90	BRAUNSCH...	76	DASP e^+e^-

$$\Gamma(\phi \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{109}/\Gamma$$

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM	15K	BES3 $e^+e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$
$0.124 \pm 0.033 \pm 0.030$		35 ± 9	² ABLIKIM	15K	BES3 $e^+e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	³ ABLIKIM	05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \phi \gamma \gamma$
<6.8	90	COFFMAN	88	MRK3	$e^+e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94±0.15 OUR AVERAGE		Error includes scale factor of 1.7.		
1.09±0.02±0.13		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ±0.9	23	FELDMAN	77 MRK1	e^+e^-

 $\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
16.0±1.0±3.0	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{113}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.74 ±0.06 OUR AVERAGE		Error includes scale factor of 1.2.		
0.71 ±0.10 ±0.05	99±14	¹ ZHU	23 BELL	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
0.898±0.024±0.089		ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.64 ±0.04 ±0.11	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.661±0.045±0.078		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta$

¹ From a fit to the combined $\phi\eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

 $\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ±0.5 OUR AVERAGE			Error includes scale factor of 2.2. See the ideogram below.		
5.10±0.03±0.32	31k		ABLIKIM	16M BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46±0.31±0.56			ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ±0.3 ±0.8	167		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
3.08±0.34±0.36			COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 13	90		VANNUCCI	77 MRK1	e^+e^-

WEIGHTED AVERAGE
 4.6 ± 0.5 (Error scaled by 2.2)



$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.32 \pm 0.06 \pm 0.16$	2.2k	¹ ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
$4.6 \pm 0.4 \pm 0.8$		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.50 \pm 0.80 \pm 0.61$	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.50 \pm 0.24$	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_0(980)\eta \rightarrow \eta \phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
$3.23 \pm 0.75 \pm 0.73$	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$

 $\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.24 \pm 0.20^{+0.52}_{-0.22}$	¹ ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$
$2.74 \pm 0.13^{+0.15}_{-0.16}$	² ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$
4.37 ± 1.35	^{3,4} ABLIKIM	18D BES3	$J/\psi \rightarrow \phi \eta \pi^0$
$5.0 \pm 2.7 \pm 2.5$	⁵ ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

¹ $J/\psi \rightarrow \phi a_0(980)$ electromagnetic decay.

² $J/\psi \rightarrow \phi f_0(980)$, $\phi a_0(980)$ mixing.

³ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

⁴ Superseded by ABLIKIM 24CB.

⁵ Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

 $\Gamma(\phi(1680)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.66 \pm 0.26^{+1.1}_{-1.0}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(X(2000)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$1.70 \pm 0.19^{+0.48}_{-0.13}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(h_1(1900)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.44 \pm 0.35^{+1.4}_{-1.2}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 $\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.6 ± 0.5 OUR AVERAGE

$3.4 \pm 1.8 \pm 1.5$	1.1k	¹ ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
$3.2 \pm 0.6 \pm 0.4$		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
$2.1 \pm 0.5 \pm 0.4$	25	² JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6 \pm 0.2 \pm 0.1$	16	BECKER	87 MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$
-----------------------	----	--------	---------	---

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.36 \pm 2.31 \pm 1.54$	78	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi 3\pi^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.08 \pm 1.63 \pm 1.47$	9	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$		172	¹ ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ With 3.6 σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE		Error includes scale factor of 2.7.		
$12.3 \pm 0.6 \pm 2.0$		^{1,2} FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$
4.8 ± 1.8	46	¹ GIDAL	81	MRK2 $J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

² Including interference with $f_0(1710)$.

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-7}$	90	¹ ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0, p\bar{p}K^+ K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

$\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{134}/Γ


VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.13 \times 10^{-5}$	90	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$

Γ_{135}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$16.6 \pm 1.9 \pm 1.2$	163 ± 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
$21.4 \pm 0.4 \pm 2.2$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$48^{+20}_{-16} \pm 6$	$9.0^{+3.7}_{-3.0}$	^{1,2} HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
$14.6 \pm 0.8 \pm 2.1$		³ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$

¹We have multiplied $K^+ K^-$ measurement by 2 to obtain $K \bar{K}$.
²Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.
³Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.



$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$

Γ_{136}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 0.2 \pm 0.6$	^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹Including interference with $f'_2(1525)$.
²Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$

Γ_{139}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.8 OUR AVERAGE				
$7.4 \pm 0.6 \pm 1.4$	227 ± 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
$7.4 \pm 0.9 \pm 1.1$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
$7 \pm 0.6 \pm 1.0$	163 ± 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.3 OUR AVERAGE				
$20.8 \pm 2.7 \pm 3.9$	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
$29.6 \pm 3.7 \pm 4.7$	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
$20.7 \pm 2.4 \pm 3.0$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
$20 \pm 3 \pm 3$	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
30 ± 5 OUR AVERAGE				
31 ± 6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29 ± 7	87	BURMESTER	77D PLUT	$e^+ e^-$

 $\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$23 \pm 3 \pm 5$	229	AUGUSTIN	89 DM2	$e^+ e^-$

 $\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.1 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$

 $\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.58 \pm 0.23 \pm 0.40$	332	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10 \pm 0.09 \pm 0.28$	233	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\bar{\Sigma}(1385)^0 p K^-)/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.51 \pm 0.26 \pm 0.18$	89	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.82 \times 10^{-5}$	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.2 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$
------------------------	------	---------	--------	-----------

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.07 OUR AVERAGE				
$0.30 \pm 0.03 \pm 0.08$	74 ± 8	HENRARD	87 DM2	$e^+ e^-$
$0.29 \pm 0.11 \pm 0.10$	26	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.08 OUR AVERAGE				
$0.34 \pm 0.04 \pm 0.08$	77	HENRARD	87	DM2 $e^+ e^-$
$0.31 \pm 0.11 \pm 0.11$	28	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.06 OUR AVERAGE				
$1.096 \pm 0.012 \pm 0.071$	43k	ABLIKIM	16L	BES3 $e^+ e^-$
$1.23 \pm 0.07 \pm 0.30$	0.8k	ABLIKIM	12P	BES2 $e^+ e^-$
$1.00 \pm 0.04 \pm 0.21$	0.6k	HENRARD	87	DM2 $e^+ e^-$
$0.86 \pm 0.18 \pm 0.22$	56	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.25 ± 0.07 OUR AVERAGE				
$1.258 \pm 0.014 \pm 0.078$	53k	ABLIKIM	16L	BES3 $e^+ e^-$
$1.50 \pm 0.08 \pm 0.38$	1k	ABLIKIM	12P	BES2 $e^+ e^-$
$1.19 \pm 0.04 \pm 0.25$	0.7k	HENRARD	87	DM2 $e^+ e^-$
$1.03 \pm 0.24 \pm 0.25$	68	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.071 \pm 0.009 \pm 0.082$	103k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons

 $\Gamma(\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.1 \times 10^{-6}$	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda \bar{\Lambda} \gamma$

 $\Gamma(\bar{\Lambda}(1520) \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.80 \times 10^{-3}$	90	LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17 ± 0.04 OUR AVERAGE				
$1.165 \pm 0.004 \pm 0.043$	135k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons
$1.20 \pm 0.12 \pm 0.21$	206	ABLIKIM	080	BES2 $e^+ e^- \rightarrow J/\psi$

 $\Gamma(\Xi(1530)^- \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.318 ± 0.008 OUR AVERAGE				
$0.317 \pm 0.002 \pm 0.008$	70k	ABLIKIM	20	BES3 $e^+ e^- \rightarrow J/\psi$
$0.59 \pm 0.09 \pm 0.12$	75	HENRARD	87	DM2 $e^+ e^-$

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.32 \pm 0.12 \pm 0.07$	24 ± 9	HENRARD	87 DM2	$e^+ e^-$

 $\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 $\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 $\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 $\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 $\Gamma(\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

————— STABLE HADRONS —————


 $\Gamma(2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
$5.44 \pm 0.07 \pm 0.33$	23K	ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
$3.5 \pm 0.8 \pm 0.1$	14k	¹ LEES	21 BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$
4.73 ± 0.44	228k	² ABLIKIM	19Q BES3	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER	77D PLUT	$e^+ e^-$
4 ± 1	675	JEAN-MARIE	76 MRK1	$e^+ e^-$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^0 \pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3}$ keV which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^0 \pi^0) = (18.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$, assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

WEIGHTED AVERAGE
 4.2 ± 0.4 (Error scaled by 2.1)



$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$			Γ_{165}/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.029 ± 0.006	OUR AVERAGE				
0.028 ± 0.009	11	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$	
0.029 ± 0.007	181	JEAN-MARIE 76	MRK1	e^+e^-	

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				Γ_{170}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
20.0 ± 0.7	OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.			
$18.78 \pm 0.13 \pm 0.51$	19.8k	¹ ANASHIN	23 KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
$21.37 \pm 0.04^{+0.64}_{-0.62}$	1.8M	² ABLIKIM	12H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
$23.0 \pm 2.0 \pm 0.4$	256	³ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow J/\psi \pi^+\pi^-\gamma$	
$21.84 \pm 0.05 \pm 2.01$	220k	^{4,5} BAI	04H BES	e^+e^-	
$20.91 \pm 0.21 \pm 1.16$		^{5,6} BAI	04H BES	e^+e^-	
15 ± 2	168	FRANKLIN	83 MRK2	e^+e^-	

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.


² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.014$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁶ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52±0.27 OUR AVERAGE		Error includes scale factor of 1.4.		
1.74±0.08±0.24	2616	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.2 ±0.3	309	VANNUCCI	77 MRK1	$e^+ e^-$

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{173}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	¹ METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+ \pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	$e^+ e^-$
1.0 ±0.5	5	BRANDELIK	78B DASP	$e^+ e^-$
1.6 ±1.6	1	VANNUCCI	77 MRK1	$e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{174}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.20±0.25 OUR AVERAGE		Error includes scale factor of 1.2.		
2.88±0.14±0.24	2654	ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-)$

$3.53 \pm 0.12 \pm 0.29$	1107	¹ ABLIKIM	05H	BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $J/\psi \pi^+ \pi^-, J/\psi \rightarrow$ $2(\pi^+ \pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1	e^+e^-

¹ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{175}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
40 ± 20	32	JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90 ± 30	13	JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(2(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29 ± 0.28 OUR AVERAGE				

$3.1 \pm 1.5 \pm 0.1$	14k	¹ LEES	21	BABR	$10.6 e^+e^- \rightarrow$ $2(\pi^+ \pi^-) 3\pi^0 \gamma$
$2.26 \pm 0.08 \pm 0.27$	4.8k	ABLIKIM	05c	BES2	$e^+e^- \rightarrow 2(\pi^+ \pi^-) \eta$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best values $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(3(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}$ Γ_{179}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.24 \pm 0.96 \pm 1.11$	616	ABLIKIM	05c	BES2 $e^+e^- \rightarrow 3(\pi^+ \pi^-) \eta$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{183}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.06 ± 0.05 OUR AVERAGE				

$3.072 \pm 0.023 \pm 0.050$	1.8 k	¹ ABLIKIM	24AB	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$
$2.86 \pm 0.09 \pm 0.19$	1k	² METREVELI	12		$\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.39 \pm 0.24 \pm 0.22$	107	³ BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ± 0.9	6	³ BRANDELIK	79c	DASP e^+e^-

¹ Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.916 \pm 0.033)\%$.

² Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.


³ Interference with non-resonant $K^+ K^-$ production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{184}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.95 ± 0.11 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.

$1.93 \pm 0.01 \pm 0.05$	110k	ABLIKIM	17AH	BES3	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
$2.62 \pm 0.15 \pm 0.14$	0.3k	¹ METREVELI	12		$\psi(2S) \rightarrow \pi^+ \pi^- K_S^0 K_L^0$

1.82±0.04±0.13 2.1k ²BAI 04A BES2 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●
1.18±0.12±0.18 JOUSSET 90 DM2 $J/\psi \rightarrow$ hadrons
1.01±0.16±0.09 74 BALTRUSAIT..85D MRK3 $e^+ e^-$
¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$		Γ_{185}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-8}$	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$<1 \times 10^{-6}$	95	¹ BAI	04D BES	$e^+ e^-$
$<5.2 \times 10^{-6}$	90	¹ BALTRUSAIT..85C	MRK3	$e^+ e^-$

¹ Forbidden by *CP*.

$\Gamma(K \bar{K} \pi)/\Gamma_{\text{total}}$		Γ_{186}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ±10 OUR AVERAGE				
55.2±12.0	25	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$
78.0±21.0	126	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$		Γ_{187}/Γ		
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.01±0.12	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{187}/\Gamma_{170}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23k	LEES	17C	BABR $J/\psi \rightarrow h^+ h^- \pi^0$

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{188}/\Gamma_{170}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24k	LEES	17C	BABR $J/\psi \rightarrow h^0 h^+ h^-$

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.04±0.26±0.92	2671	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ±2.3	205	VANNUCCI	77	MRK1 $e^+ e^-$
----------	-----	----------	----	----------------

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{204}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
31±13	30	VANNUCCI	77	MRK1 $e^+ e^-$

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	¹ HUANG	03	BELL $B^+ \rightarrow 2(K^+ K^-) K^+$
0.7 ± 0.3		VANNUCCI	77	MRK1 $e^+ e^-$

¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$. $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.120±0.029 OUR AVERAGE

2.112±0.004±0.031	314k	ABLIKIM	12C	BES3 $e^+ e^-$
2.17 ±0.16 ±0.04	317	¹ WU	06	BELL $B^+ \rightarrow p\bar{p}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E	BES2 $e^+ e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98	FENI $e^+ e^-$
1.91 ±0.04 ±0.30		PALLIN	87	DM2 $e^+ e^-$
2.16 ±0.07 ±0.15	1420	EATON	84	MRK2 $e^+ e^-$
2.5 ±0.4	133	BRANDELIK	79C	DASP $e^+ e^-$
2.0 ±0.5		BESCH	78	BONA $e^+ e^-$
2.2 ±0.2	331	² PERUZZI	78	MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ±0.3	48	ANTONELLI	93	SPEC $e^+ e^-$
----------	----	-----------	----	----------------

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$


Γ_{215}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.1.		
$1.33 \pm 0.02 \pm 0.11$	11k	ABLIKIM	09B	BES2 e^+e^-
$1.13 \pm 0.09 \pm 0.09$	685	EATON	84	MRK2 e^+e^-
1.4 ± 0.4		BRANDELIK	79C	DASP e^+e^-
1.00 ± 0.15	109	PERUZZI	78	MRK1 e^+e^-

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{216}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$6.46 \pm 0.17 \pm 0.43$	1435	EATON	84	MRK2 e^+e^-
3.8 ± 1.6	48	BESCH	81	BONA e^+e^-
5.5 ± 0.6	533	PERUZZI	78	MRK1 e^+e^-



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{217}/Γ

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$

Γ_{218}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
$1.91 \pm 0.02 \pm 0.17$	13k	¹ ABLIKIM	09	BES2 e^+e^-

$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2	e^+e^-
2.5 ± 1.2		BRANDELIK	79C	DASP	e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1	e^+e^-

¹From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$


Γ_{219}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.31 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

Γ_{220}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$

Γ_{221}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.129 ± 0.014 OUR AVERAGE	Error includes scale factor of 2.0.			
$0.126 \pm 0.002 \pm 0.007$	16k	¹ ABLIKIM	19N	BES3 e^+e^-
$0.200 \pm 0.023 \pm 0.028$	265 ± 31	² ABLIKIM	09	BES2 e^+e^-
$0.68 \pm 0.23 \pm 0.17$	19	EATON	84	MRK2 e^+e^-
1.8 ± 0.6	19	PERUZZI	78	MRK1 e^+e^-

¹From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

²From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

$\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{222}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$6.8 \pm 1.2 \pm 1.3$	ABLIKIM	14N	BES3 $e^+e^- \rightarrow J/\psi$

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{223}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.519 ± 0.033 OUR AVERAGE				
$0.523 \pm 0.006 \pm 0.033$	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0$, $p\bar{p}K^+ K^-$
$0.45 \pm 0.13 \pm 0.07$		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{224}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.12 ± 0.09 OUR AVERAGE				
$2.36 \pm 0.02 \pm 0.21$	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow p\pi^-\bar{n}$
$2.47 \pm 0.02 \pm 0.24$	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{p}\pi^+n$
$2.02 \pm 0.07 \pm 0.16$	1288	EATON	84	MRK2 $e^+e^- \rightarrow p\pi^-$
$1.93 \pm 0.07 \pm 0.16$	1191	EATON	84	MRK2 $e^+e^- \rightarrow \bar{p}\pi^+$
1.7 ± 0.7	32	BESCH	81	BONA $e^+e^- \rightarrow p\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA $e^+e^- \rightarrow \bar{p}\pi^+$
2.16 ± 0.29	194	PERUZZI	78	MRK1 $e^+e^- \rightarrow p\pi^-$
2.04 ± 0.27	204	PERUZZI	78	MRK1 $e^+e^- \rightarrow \bar{p}\pi^+$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{225}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.09 ± 0.16 OUR AVERAGE				
$2.07 \pm 0.01 \pm 0.17$	36k	ABLIKIM	12C	BES3 e^+e^-
2.31 ± 0.49	79	BALDINI	98	FENI e^+e^-
1.8 ± 0.9		BESCH	78	BONA e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90 ± 0.55	40	ANTONELLI	93	SPEC e^+e^-


 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{226}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 3.6	5	BESCH	81	BONA e^+e^-

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{230}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.88 ± 0.08 OUR AVERAGE		Error includes scale factor of 2.6. See the ideogram below.		
$1.64 \pm 0.12 \pm 0.09$		ABLIKIM	23S	BES3 $e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$
$1.943 \pm 0.003 \pm 0.033$	441k	ABLIKIM	17L	BES3 e^+e^-
$2.03 \pm 0.03 \pm 0.15$	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
$1.96^{+0.47}_{-0.44} \pm 0.04$	46	¹ WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
$1.08 \pm 0.06 \pm 0.24$	631	BAI	98G	BES e^+e^-
$1.38 \pm 0.05 \pm 0.20$	1847	PALLIN	87	DM2 e^+e^-
$1.58 \pm 0.08 \pm 0.19$	365	EATON	84	MRK2 e^+e^-
2.6 ± 1.6	5	BESCH	81	BONA e^+e^-
1.1 ± 0.2	196	PERUZZI	78	MRK1 e^+e^-

¹WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{231}/Γ

VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G BES	e^+e^-
22 ±5 ±5	19	HENRARD	87 DM2	e^+e^-

¹Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

²Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{232}/Γ

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P BES2	J/ψ

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{233}/Γ

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$. $\Gamma(\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{234}/\Gamma$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.26 $\pm$ 0.05 OUR AVERAGE		Error includes scale factor of 1.2.		
1.244 $\pm$ 0.002 $\pm$ 0.045	2.6M	ABLIKIM	23BU BES3	$e^+ e^-$
1.52 $\pm$ 0.08 $\pm$ 0.16	589	¹ ABLIKIM	07H BES2	$e^+ e^-$
1.11 $\pm$ 0.06 $\pm$ 0.20	342 $\pm$ 18	HENRARD	87 DM2	$e^+ e^-$
1.38 $\pm$ 0.21 $\pm$ 0.35	118	EATON	84 MRK2	$e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$. $\Gamma(\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{235}/\Gamma$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 $\pm$ 0.07 OUR AVERAGE		Error includes scale factor of 1.8.		
1.221 $\pm$ 0.002 $\pm$ 0.038	2.7M	ABLIKIM	23BU BES3	$e^+ e^-$
0.90 $\pm$ 0.06 $\pm$ 0.16	225	HENRARD	87 DM2	$e^+ e^-$
1.53 $\pm$ 0.17 $\pm$ 0.38	135	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{236}/\Gamma$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 $\pm$ 0.11 OUR AVERAGE				
0.84 $^{+0.17}_{-0.15} \pm$ 0.02	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
0.89 $\pm$ 0.07 $\pm$ 0.14	307	EATON	84 MRK2	$e^+ e^-$

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p K^- \bar{\Sigma}^0)/\Gamma_{\text{total}} \quad \Gamma_{237}/\Gamma$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29 $\pm$ 0.06 $\pm$ 0.05	90	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(p K_S^0 \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{238}/\Gamma$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.725 $\pm$ 0.009 $\pm$ 0.050	120k	¹ ABLIKIM	24H BES3	$e^+ e^- \rightarrow J/\psi$

¹ The branching fractions for the charge-conjugate channels are measured separately as $(1.361 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $\bar{p} K_S^0 \Sigma^+$ and $(1.352 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $p K_S^0 \bar{\Sigma}^-$.

 $\Gamma(\bar{\Lambda} n K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{239}/\Gamma$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46 $\pm$ 0.20 $\pm$ 1.07	1058	¹ ABLIKIM	08C BES2	$e^+ e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+ \pi^-) = 69.2\%$.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{240}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

2.83±0.23 OUR AVERAGE2.74±0.24±0.22 234 ± 21 ¹ ABLIKIM 12B BES3 $J/\psi \rightarrow \Lambda\bar{\Sigma}^0$ 2.92±0.22±0.24 308 ± 24 ² ABLIKIM 12B BES3 $J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18 ² HENRARD 87 DM2 $J/\psi \rightarrow \bar{\Lambda}\Sigma^0$ <15 90 PERUZZI 78 MRK1 $e^+e^- \rightarrow \Lambda X$ ¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2. $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{241}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.07 ±0.04 OUR AVERAGE1.061±0.004±0.036 87k ABLIKIM 21AT BES3 $J/\psi \rightarrow p\pi^0\bar{p}\pi^0$ 1.50 ±0.10 ±0.22 399 ABLIKIM 08O BES2 $e^+e^- \rightarrow J/\psi$ $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{242}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.172±0.032 OUR AVERAGE Error includes scale factor of 1.4.1.164±0.004±0.023 111k ABLIKIM 17L BES3 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$ 1.33 ±0.04 ±0.11 1.7k ABLIKIM 06 BES2 $J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$ 1.06 ±0.04 ±0.23 884 PALLIN 87 DM2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$ 1.58 ±0.16 ±0.25 90 EATON 84 MRK2 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$ 1.3 ±0.4 52 PERUZZI 78 MRK1 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •


2.4 ±2.6 3 BESCH 81 BONA $e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$ $\Gamma(\Sigma^+\bar{\Sigma}^-\eta)/\Gamma_{\text{total}}$ Γ_{243}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.34±0.21±0.37 1821 ABLIKIM 22AY BES3 $J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-\eta$ $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{244}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.97 ±0.08 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.1.040±0.006±0.074 43k ABLIKIM 16L BES3 $J/\psi \rightarrow \Xi^-\bar{\Xi}^+$ 0.90 ±0.03 ±0.18 961 ABLIKIM 12P BES2 $J/\psi \rightarrow \Xi^-\bar{\Xi}^+$ 0.70 ±0.06 ±0.12 132 HENRARD 87 DM2 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ 1.14 ±0.08 ±0.20 194 EATON 84 MRK2 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ 1.4 ±0.5 51 PERUZZI 78 MRK1 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$



RADIATIVE DECAYS

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{245}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.41 ± 0.14 OUR FIT Error includes scale factor of 1.3.

1.7 ± 0.4 OUR AVERAGE Error includes scale factor of 1.5.

$2.00 \pm 0.31 \pm 0.02$	¹ MITCHELL	09	CLEO	$e^+e^- \rightarrow \gamma X$
1.27 ± 0.36	GAISER	86	CBAL	$J/\psi \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.40 ± 0.33	² ANASHIN	14	KEDR	$J/\psi \rightarrow \gamma\eta_c$
-----------------	----------------------	----	------	-----------------------------------

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Statistical uncertainty only.

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{248}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

11.6 ± 2.2 OUR AVERAGE

$11.3 \pm 1.8 \pm 2.0$	113 ± 18	ABLIKIM	13I	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
$12 \pm 3 \pm 2$	$24.2^{+7.2}_{-6.0}$	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	PARTRIDGE	80	CBAL	e^+e^-
-------	----	-----------	----	------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$					Γ_{249}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<9 \times 10^{-6}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$					Γ_{250}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<15 \times 10^{-6}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$					Γ_{251}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.39$\pm$0.08 OUR AVERAGE					
3.34 $\pm$ 0.02 $\pm$ 0.09	176k	ABLIKIM	23BD BES3	$J/\psi \rightarrow \pi^0 \gamma$	
3.59 $\pm$ 0.20 $\pm$ 0.03	1.6k	¹ ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$	
3.63 $\pm$ 0.36 $\pm$ 0.13		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0 \gamma$	
3.13 $^{+0.65}_{-0.47}$	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0 \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 $\pm$ 1.1 $\pm$ 0.7		BLOOM	83	CBAL	$e^+ e^-$
7.3 $\pm$ 4.7	10	BRANDELIK	79C	DASP	$e^+ e^-$


¹ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_{252}/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT	
1.15$\pm$0.05		¹ ABLIKIM	15AE BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$	

¹The uncertainty is systematic as statistical is negligible.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$					Γ_{253}/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT	
2.8 $\pm$0.5 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.					
4.32 $\pm$ 0.14 $\pm$ 0.73		¹ BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08 $\pm$ 0.13 $\pm$ 0.35		² BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05 $\pm$ 0.08 $\pm$ 0.45		² BALTRUSAIT..	86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 $\pm$ 0.45 $\pm$ 1.20		³ BURKE	82	MRK2	$e^+ e^-$
¹ 4 π mass less than 3.0 GeV.					
² 4 π mass less than 2.0 GeV.					
³ 4 π mass less than 2.5 GeV.					

WEIGHTED AVERAGE
 2.8 ± 0.5 (Error scaled by 1.9)



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$

Γ_{254}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.7 \pm 1.6$	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$

Γ_{255}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$8.2 \pm 0.8 \pm 1.7$	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$

Γ_{256}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.2 \pm 3.1$	¹ BALTRUSAIT...86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{257}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma (K \bar{K} \pi) [J^{PC} = 0^- +])/ \Gamma_{\text{total}}$

Γ_{258}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
$0.58 \pm 0.03 \pm 0.20$	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$2.1 \pm 0.1 \pm 0.7$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

¹ For a broad structure around 1800 MeV.² For a broad structure around 2040 MeV. $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{259}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.1 \pm 0.1 \pm 0.6$	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

 $\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{260}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges. $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{261}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.090 ± 0.013 OUR AVERAGE				
$1.096 \pm 0.001 \pm 0.019$	2.2M	ABLIKIM	23BD BES3	$J/\psi \rightarrow \eta\gamma$
$1.067 \pm 0.005 \pm 0.023$	87.9k	ABLIKIM	21AMBES3	$e^+e^- \rightarrow J/\psi$
$1.12 \pm 0.05 \pm 0.01$	18.6k	¹ ABLIKIM	18O BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
$1.101 \pm 0.029 \pm 0.022$		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta\gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.88 \pm 0.08 \pm 0.11$		BLOOM	83 CBAL	e^+e^-
0.82 ± 0.10		BRANDELIK	79C DASP	e^+e^-
1.3 ± 0.4	21	BARTEL	77 CNTR	e^+e^-

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{262}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$21.4 \pm 1.8 \pm 2.5$	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{263}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.5 ± 2.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{264}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5 ± 5	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{265}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4 ± 3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0) / \Gamma_{\text{total}}$ Γ_{266} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-6}$	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0) / \Gamma_{\text{total}}$ Γ_{267} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-6}$	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma \eta \pi \pi) / \Gamma_{\text{total}}$ Γ_{268} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 $\pm$ 1.0 OUR AVERAGE			
5.85 $\pm$ 0.3 $\pm$ 1.05	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta \pi^+ \pi^-$
7.8 $\pm$ 1.2 $\pm$ 2.4	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$

¹ Broad enhancement at 1700 MeV.

 $\Gamma(\gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{269} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.2 $\pm$ 2.2 $\pm$ 0.9	BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

 $\Gamma(\gamma \eta'(958)) / \Gamma_{\text{total}}$ Γ_{270} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.28 $\pm$ 0.06 OUR AVERAGE	Error includes scale factor of 1.3.			See the ideogram below.
5.40 $\pm$ 0.01 $\pm$ 0.11	638k	ABLIKIM	23BD	BES3 $J/\psi \rightarrow \gamma \eta'$
5.27 $\pm$ 0.03 $\pm$ 0.05	36k	ABLIKIM	19T	BES $J/\psi \rightarrow \gamma \eta'$
5.43 $\pm$ 0.23 $\pm$ 0.09	5.0k	¹ ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
4.76 $\pm$ 0.22 $\pm$ 0.06		² ABLIKIM	11	BES3 $J/\psi \rightarrow \eta' \gamma$
5.24 $\pm$ 0.12 $\pm$ 0.11		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta' \gamma$
5.55 $\pm$ 0.44	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta' \gamma$


• • • We do not use the following data for averages, fits, limits, etc. • • •

4.50 $\pm$ 0.14 $\pm$ 0.53		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \gamma \gamma$
4.30 $\pm$ 0.31 $\pm$ 0.71		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$
4.04 $\pm$ 0.16 $\pm$ 0.85	622	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
4.39 $\pm$ 0.09 $\pm$ 0.66	2420	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
4.1 $\pm$ 0.3 $\pm$ 0.6		BLOOM	83	CBAL $e^+ e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 $\pm$ 1.1	6	BRANDELIK	79C	DASP $e^+ e^- \rightarrow 3\gamma$
2.4 $\pm$ 0.7	57	BARTEL	76	CNTR $e^+ e^- \rightarrow 2\gamma \rho$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958)) / \Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma \gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958)) / \Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma \gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma \gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958)) / \Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+ \pi^- \eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we

rescale to our best values $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.



$\Gamma(\gamma f_0(980) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{271}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.3±0.2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{272}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8±0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma \rho \rho)/\Gamma_{\text{total}}$ Γ_{273}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
4.5 ±0.8 OUR AVERAGE				
4.7 ±0.3 ±0.9		¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75±1.05±1.20		² BURKE 82	MRK2	$J/\psi \rightarrow 4\pi\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.09	90	³ BISELLO 89B		$J/\psi \rightarrow 4\pi\gamma$
¹ 4π mass less than 2.0 GeV.				
² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .				
³ 4π mass in the range 2.0–25 GeV.				

$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$

Γ_{274}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-4}$	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$

Γ_{275}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.8 \times 10^{-5}$	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$

Γ_{276}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61 ± 0.33 OUR AVERAGE				
$6.0 \pm 4.8 \pm 1.8$		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
$1.41 \pm 0.2 \pm 0.42$	120 ± 17	BISELLO	87 SPEC	e^+e^- , hadrons γ
$1.76 \pm 0.09 \pm 0.45$		BALTRUSAIT..85C	MRK3	$e^+e^- \rightarrow$ hadrons γ

$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$

Γ_{277}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.2 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ $\phi\phi$ mass less than 2.9 GeV, η_C excluded.




$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$

Γ_{278} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1 ± 0.58	^{1,2} BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
3.8 ± 0.3 ± 0.6	³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
4.0 ± 0.7 ± 1.0	³ EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80 MRK2	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.78 ± 0.21 ± 0.33	^{3,5,6} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.83 ± 0.13 ± 0.18	^{3,7,8} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.66 ^{+0.17 +0.24} _{-0.16 -0.15}	^{3,6,9} BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$
1.03 ^{+0.21 +0.26} _{-0.18 -0.19}	^{3,8,10} BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

- ¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K \bar{K} \pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.
- ² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.
- ³ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.
- ⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.
- ⁵ From fit to the $a_0(980) \pi$ 0^-+ partial wave.
- ⁶ $a_0(980) \pi$ mode.
- ⁷ From fit to the $K^*(892) K$ 0^-+ partial wave.
- ⁸ $K^* K$ mode.
- ⁹ From $a_0(980) \pi$ final state.
- ¹⁰ From $K^*(890) K$ final state.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{279}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78±0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07±0.17±0.11	¹ BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64±0.12±0.07	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{280}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ±0.5 OUR AVERAGE				
2.6 ±0.7 ±0.4		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38±0.33±0.64		¹ BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ±0.6 ±1.1	261	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
---------------	-----	-----------------------	----	---

¹Via $a_0(980)\pi$.

²Includes unknown branching fraction to $\eta\pi^+\pi^-$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{281}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ±0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ±0.4	BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36±0.38	^{1,2} BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹Estimated by us from various fits.

²Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{282}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<82	95		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.03±0.92±0.91	1.3k		¹ ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
10.36±1.51±1.54	1.9k		² ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

²Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{283}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.11±0.11	743	ABLIKIM	12E	BES3 $J/\psi \rightarrow \gamma\eta(1405)$

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^0\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{284}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
7.10±0.82±0.72	198	ABLIKIM	12E	BES3 $J/\psi \rightarrow \gamma\eta(1405)$

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{285}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.63 × 10⁻⁶	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{286}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.86 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{287}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.09	1,2 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0\rho^0$. $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{288}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{289}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.80 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

 $\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{290}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.14^{+0.50}_{-0.19}$ OUR AVERAGE				

$2.40 \pm 0.10^{+2.47}_{-0.18}$		1,2 ABLIKIM	16N	BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I	BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$2.4^{+1.5}_{-1.0}$		3,4 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).² Includes unknown branching fraction to $\phi\phi$.³ Estimated by us from various fits.⁴ Includes unknown branching fraction to $\rho^0\rho^0$. $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{291}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.63 ± 0.12 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$2.07 \pm 0.16^{+0.02}_{-0.07}$	2.4k	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$1.63 \pm 0.26^{+0.02}_{-0.05}$		3 ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.42 \pm 0.21^{+0.02}_{-0.05}$		4 ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
$1.33 \pm 0.05 \pm 0.20$		5 AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.36 \pm 0.09 \pm 0.23$		5 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.48 \pm 0.25 \pm 0.30$	178	EDWARDS	82B	CBAL $e^+e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78	PLUT e^+e^-
1.2 ± 0.6	30	6 BRANDELIK	78B	DASP $e^+e^- \rightarrow \pi^+\pi^-\gamma$

¹Using CLEO-c data but not authored by the CLEO Collaboration.
²DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
³ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁴ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁵Estimated using $B(f_2(1270) \rightarrow \pi\pi)=0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.
⁶Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{292}/Γ
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$				Γ_{293}/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	
0.61 ± 0.08 OUR AVERAGE				
$0.69 \pm 0.16 \pm 0.20$	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \rho^0$	

$0.61 \pm 0.04 \pm 0.21$	² BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.45 \pm 0.09 \pm 0.17$	³ BAI	99	BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.625 \pm 0.063 \pm 0.103$	⁴ BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
$0.70 \pm 0.08 \pm 0.16$	⁵ BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta \pi \pi) / \Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.

⁵ Using $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta \pi$.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{294} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
38 ± 10	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{295} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.19 \pm 0.73 \pm 1.34$	478	¹ DOBBS 15		$J/\psi \rightarrow \gamma K \bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.3 ± 0.4		SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{296} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{297} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.5 ± 1.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{298} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.9 ± 0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{299} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.13 OUR AVERAGE			
$0.68 \pm 0.04 \pm 0.24$	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

$0.76 \pm 0.15 \pm 0.21$	^{1,2} AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.87 \pm 0.14^{+0.14}_{-0.11}$	¹ BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

²From fit to the $K^*(892) K 1^{++}$ partial wave.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{300} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.24 OUR AVERAGE				
$1.21 \pm 0.29 \pm 0.24$	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$1.00 \pm 0.03 \pm 0.45$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1.02 \pm 0.09 \pm 0.45$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ± 0.17	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
5.7 ± 0.8	^{3,4} BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$

¹Using CLEO-c data but not authored by the CLEO Collaboration.

²Including unknown branching fraction to $\pi \pi$.

³Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

⁴Assuming that $f_0(1500)$ decays only to two S -wave dipions.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{301} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.1 ± 0.4	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{302} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.3	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{303} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$18.1 \pm 1.1^{+1.9}_{-1.3}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

12 ± 5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
------------	-----------	----	------	--

¹From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{304} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f'_2(1525)) / \Gamma_{\text{total}}$ Γ_{305} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------


$5.7^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.0 \pm 0.9 \pm 0.2$	750	^{1,2} DOBBS	15		$J/\psi \rightarrow \gamma K \bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		³ BAI	03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		³ BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		³ BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.4	90	4	⁴ BRANDELIK	79C	DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
< 2.3	90	3	ALEXANDER	78	PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$

WEIGHTED AVERAGE
 $5.7^{+0.8}_{-0.5}$ (Error scaled by 1.5)



¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525)) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{306}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{307}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1565) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{308}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.32 \pm 0.05^{+0.12}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{309}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{310}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5 OUR AVERAGE				
$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.6 ± 0.2		³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K\bar{K}, \eta \eta, \omega \phi)$
$2.5 \pm 1.6 \pm 0.8$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{311}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.5 ± 1.0 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.
$8.00^{+0.12+1.24}_{-0.08-0.40}$			¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$11.76 \pm 0.54 \pm 0.94$		1.2k	² DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$9.62 \pm 0.29^{+3.51}_{-1.86}$			³ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$

$5.0 \pm 0.8 \begin{smallmatrix} +1.8 \\ -0.4 \end{smallmatrix}$		^{1,4} BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
$9.2 \pm 1.4 \pm 1.4$		¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$		¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$		¹ BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.3 ± 0.8		⁵ SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma$ ($\pi\pi, K\bar{K}, \eta\eta, \omega\phi$)
$1.6 \pm 0.2 \begin{smallmatrix} +0.6 \\ -0.2 \end{smallmatrix}$		^{1,6} BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90	⁷ BISELLO	89B		$J/\psi \rightarrow 4\pi\gamma$
$1.6 \pm 0.4 \pm 0.3$		⁸ BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 ± 1.6		⁹ EDWARDS	82D	CBAL	$e^+e^- \rightarrow \eta\eta\gamma$

¹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.


⁵ There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.

⁶ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁷ Includes unknown branching fraction to $\rho^0 \rho^0$.

⁸ Includes unknown branching fraction to $\pi^+ \pi^-$.

⁹ Includes unknown branching fraction to $\eta\eta$.



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{312}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{313}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.4 ² SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{314}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.5 ± 2.5	¹ SARANTSEV 21 RVUE		$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$ Γ_{315}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.6 OUR AVERAGE				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1 ± 0.1 ¹ SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{316}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{317}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma \eta_1(1855) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{318}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.70 \pm 0.41^{+0.16}_{-0.35}$	¹ ABLIKIM	22AI BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave. For analysis details see ABLIKIM 22AS.

$\Gamma(\gamma f_0(1770) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{319}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.11 \pm 0.01^{+0.04}_{-0.03}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{320}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--	-----	---------	---------	---

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{321}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$0.7 \pm 0.1 \pm 0.2$	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
---	-----	---------	---

$\Gamma(\gamma f_2(2010) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{322}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.06^{+0.10}_{-0.06}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{323}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

42 ± 10	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$\Gamma(\gamma f_0(2020) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{324}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

55 ± 25	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{325}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10 ± 10	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{326}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.63 \pm 0.06^{+0.31}_{-0.46}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
---	----------------------	----------	---

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{327} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.28 \pm 0.12^{+0.29}_{-0.20}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{328} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
---	-----------------------------	------	---

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

 $\Gamma(\gamma f_4(2050) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{329} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.01^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{330} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$
--	------	----------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 1.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{331} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

32 ± 20	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
-------------	-----------	----	------	--

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{332} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--	-----	--------------------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.8	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$ Γ_{333} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	----	-----	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{334}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.49 \pm 1.20$	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.5 ± 0.5		SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K \bar{K}, \eta\eta, \omega\phi)$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}$ Γ_{335}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 \bar{K}_S^0$

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{336}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5 ± 2	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K \bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{337}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.7 ± 0.4	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K \bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{338}/Γ

VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
> 300			¹ BAI	96B	BES $e^+e^- \rightarrow \gamma \bar{p} p, K \bar{K}$
> 250	99.9		² HASAN	96	SPEC $\bar{p} p \rightarrow \pi^+ \pi^-$
< 2.3	95		³ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		³ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 \bar{K}_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 \bar{K}_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 \bar{K}_S^0$.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{339}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$14 \pm 8 \pm 4$		BAI	98H	BES $J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$		BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{340}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6		³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.6 \pm 2.9 \pm 2.4$		BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{341}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{342}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.95 \pm 0.21^{+0.66}_{-0.72}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{343}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
4 ± 2	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{344}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
1.5 ± 0.4	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{345}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.09 \pm 0.64^{+4.00}_{-1.68}$	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{346}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
$0.10 \pm 0.02^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{347}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.60^{+0.62+2.37}_{-0.65-2.07}	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{348}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
5.54^{+0.34+3.82}_{-0.40-1.49}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{349}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
8.67$\pm$0.70^{+0.61}_{-1.67}	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_0(2470) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{350}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
8.18$\pm$1.77^{+3.73}_{-2.23}	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{351}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.7^{+0.6}_{-0.8}				OUR AVERAGE Error includes scale factor of 1.6.

3.93 $\pm$ 0.38^{+0.31}_{-0.84} ¹ ABLIKIM 16J BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

2.2 $\pm$ 0.4 $\pm$ 0.4 264 ABLIKIM 05R BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.87 $\pm$ 0.09^{+0.49}_{-0.52} 4265 ² ABLIKIM 11C BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma\eta(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{352}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.77^{+0.15}_{-0.09}$				OUR AVERAGE

$0.90^{+0.04+0.27}_{-0.11-0.55}$		¹ ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma p \bar{p}$
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10	CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F	BES2	$J/\psi \rightarrow \gamma p \bar{p}$

¹ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p \bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{353}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.31^{+0.33+1.96}_{-0.30-1.29}$	ABLIKIM	15T	BES3
			$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \phi(1020))/\Gamma_{\text{total}}$ Γ_{354}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.77 \pm 0.35 \pm 0.25$	305	¹ ABLIKIM	18i	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
$8.09 \pm 1.99 \pm 1.36$	1.3k	² ABLIKIM	18i	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{355}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.56 × 10⁻⁶	90	ABLIKIM	18o	BES3
				$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{356}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.44 \pm 0.36^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U	BES3
$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$				

 $\Gamma(\gamma \eta(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{357}/Γ

<i>VALUE</i> (units 10^{-5})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
$1.79 \pm 0.23 \pm 0.65$	ABLIKIM	20Q	BES3
			$J/\psi \rightarrow \gamma K^+ K^- \eta'$

 $\Gamma(\gamma \eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{358}/Γ

<i>VALUE</i> (units 10^{-5})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
$1.18 \pm 0.32 \pm 0.39$	ABLIKIM	20Q	BES3
			$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.31 \pm 0.22^{+2.85}_{-0.84}$	¹ ABLIKIM	24	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$
---------------------------------	----------------------	----	------	---

¹ Decaying via the intermediate $f_0(980) \eta'$, fitted together with $X(1835)$, a 600 MeV broad structure around 2.8 GeV, and the tail of the $\eta_c(1S)$.

$\Gamma(\gamma\eta(2370) \rightarrow \gamma\eta\eta\eta')/\Gamma_{\text{total}}$ Γ_{359}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.2 \times 10^{-6}$	90	ABLIKIM	21c	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$

 $\Gamma(\gamma D^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{360}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.1 \times 10^{-8}$	90	ABLIKIM	24BZ	BES3 $e^+e^- \rightarrow J/\psi(1S)$

 $\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{361}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.38 \pm 0.07 \pm 0.07$		49	EATON	84	MRK2 e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.11	90		PERUZZI	78	MRK1 e^+e^-

 $\Gamma(\gamma p \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{362}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2 e^+e^-

 $\Gamma(\gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{363}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<0.16 \times 10^{-3}$	90	BAI	98G	BES $e^+ e^-$

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{364}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.3 \times 10^{-6}$	90	3.7M	² INSLE	10	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

² The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$, reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

 $\Gamma(\gamma A^0 \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{365}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.9 \times 10^{-7}$	95	¹ ABLIKIM	24AD	BES3 $J/\psi \rightarrow \gamma\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.8 \times 10^{-6}$	95	² ABLIKIM	23E	BES3 $J/\psi \rightarrow \gamma\gamma\gamma$

¹ For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.18–2.85 GeV. The measured 95% CL limit as a function of m_{A^0} ranges from 3.7×10^{-8} to 4.85×10^{-7} .

² For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.165–2.84 GeV. The measured 95% CL limit as a function of m_{A^0} ranges from 8.3×10^{-8} to 1.8×10^{-6} .

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{366}/Γ
 (narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.8 \times 10^{-7}$	90	¹ ABLIKIM 22H	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.5 \times 10^{-5}$	90	² ABLIKIM 16E	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$< 2.1 \times 10^{-5}$	90	³ ABLIKIM 12	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

³ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

———— DALITZ DECAYS ————

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{367}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM 14i	BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{368}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.04 \pm 0.07$	2.47k	^{1,2} ABLIKIM 19A	BES3	$J/\psi \rightarrow \eta e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM 14i	BES3	$J/\psi \rightarrow \eta e^+ e^-$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14i.

$\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{369}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM 19H	BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.81 \pm 0.16 \pm 0.31$	1.4k	^{1,2} ABLIKIM 14i	BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

¹ Using both $\eta' \rightarrow \gamma \pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

² Superseded by ABLIKIM 19H.

$\Gamma(\eta(1405) e^+ e^- \rightarrow f_0(980) \pi^0 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{370}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.04 \pm 0.20 \pm 0.08$	203	¹ ABLIKIM 24i	BES3	$J/\psi \rightarrow e^+ e^- \eta(1405)$

¹ With a significance of 9.8 σ .

$\Gamma(X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{371}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.58 \pm 0.19 \pm 0.16$	1364	¹ ABLIKIM 22B	BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

¹ Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

$\Gamma(X(2120)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{372}/Γ

VALUE (units 10^{-6})	EVTs	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta' e^+e^-$

 $\Gamma(\eta(2370)e^+e^-, \eta \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{373}/Γ

VALUE (units 10^{-6})	EVTs	DOCUMENT ID	TECN	COMMENT
$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta' e^+e^-$

 $\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{374}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.11 \times 10^{-7}$	90	¹ ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+e^-$

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

 $\Gamma(\eta'(958)U \rightarrow \eta'(958)e^+e^-)/\Gamma_{\text{total}}$ Γ_{375}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-7}$	90	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+e^-$

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

 $\Gamma(\phi e^+e^-)/\Gamma_{\text{total}}$ Γ_{376}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	¹ ABLIKIM	19AB BES3	$J/\psi \rightarrow \phi e^+e^-$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.49 \pm 0.30)\%$.

WEAK DECAYS

 $\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{377}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.1 \times 10^{-8}$	90	ABLIKIM	21Q BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.2 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
------------------------	----	---------	----------	-----------------------------

 $\Gamma(D^- \mu^+ \nu_\mu + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{378}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-7}$	90	¹ ABLIKIM	24AMBES3	$e^+e^- \rightarrow J/\psi$

¹ Using $B(D^- \rightarrow K^+ \pi^- \pi^-) = 9.38 \pm 0.16 \%$.

 $\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{379}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.5 \times 10^{-8}$	90	¹ ABLIKIM	17AF BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.1 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
------------------------	----	---------	----------	-----------------------------

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{380}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R	BES3	$e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.6 \times 10^{-5}$	90	¹ ABLIKIM	06M	BES2	$e^+ e^- \rightarrow J/\psi$
¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$.					

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{381}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-6}$	90	ABLIKIM	14R	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{382}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.0 \times 10^{-8}$	90	ABLIKIM	24BI	BES3	$e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<7.5 \times 10^{-5}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{383}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.0 \times 10^{-7}$	90	ABLIKIM	24BI	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{384}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.7 \times 10^{-7}$	90	ABLIKIM	24BI	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{385}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-4}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{386}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	ABLIKIM	14K	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \eta + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{387}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.8 \times 10^{-7}$	90	ABLIKIM	24BI	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{388}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.2 \times 10^{-7}$	90	ABLIKIM	24BI	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{389}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-4}$	90	ABLIKIM	08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{390}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K	BES3	$e^+ e^- \rightarrow J/\psi$

CHARGE CONJUGATION (C), PARITY (P), LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{391}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-7}$	90	ABLIKIM	14Q	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 0.5 \times 10^{-5}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$< 1.6 \times 10^{-4}$	90	¹ WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma \gamma$
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J	BES2 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$< 50 \times 10^{-5}$	90	BARTEL	77	CNTR $e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{392}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-6}$	90	ABLIKIM	14Q	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{393}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-7}$	90	ABLIKIM	13L	BES3 $e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.1 \times 10^{-6}$	90	BAI	03D	BES $e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{394}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.5 \times 10^{-8}$	90	ABLIKIM	21M	BES3 $e^+ e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 8.3 \times 10^{-6}$	90	¹ ABLIKIM	04	BES $e^+ e^- \rightarrow J/\psi$

¹ Superseded by ABLIKIM 21M.

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{395}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-6}$	90	ABLIKIM	04	BES $e^+ e^- \rightarrow J/\psi$

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{396}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.9 \times 10^{-8}$	90	ABLIKIM	19AF	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow p K^- \pi^+ e^- (+ \text{c.c.})$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$ Γ_{397}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.6 \times 10^{-2}$	90	LEES	13I	BABR $B \rightarrow K^{(*)} J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$ Γ_{397}/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$$\Gamma(\mu^+\mu^-X^0 \rightarrow \mu^+\mu^- + \text{invisible})/\Gamma_{\text{total}} \quad \Gamma_{398}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-7}$	90	¹ ABLIKIM	24G BES3	$J/\psi \rightarrow \mu^+\mu^-X^0$ (scalar) $\rightarrow$ $\mu^+\mu^- + \text{invisible}$
$<9.6 \times 10^{-7}$	90	² ABLIKIM	24G BES3	$J/\psi \rightarrow \mu^+\mu^-X^0$ (vector) $\rightarrow$ $\mu^+\mu^- + \text{invisible}$

¹ For a light scalar, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 6.2×10^{-9} to 5.5×10^{-7} .

² For a light vector, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 4.5×10^{-9} to 9.6×10^{-7} .

$J/\psi(1S)$ REFERENCES

AAIJ	24AE	JHEP 2412 062	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	24	PRL 132 181901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AB	PR D110 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AD	PR D110 L031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AM	JHEP 2401 126	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BI	PR D110 032020	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BQ	PR D110 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BZ	PR D110 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CB	PR D110 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24G	PR D109 L031102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24H	PR D109 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24I	PR D109 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24L	PR D109 052006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAYRAPETY...	24A	PR D109 L111101	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
ABLIKIM	23BD	PR D108 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BU	PR D108 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23E	PL B838 137698	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23S	PR D107 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	23	JHEP 2306 196	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
GONG	23	PR D107 072008	G. Gong <i>et al.</i>	(BELLE Collab.)
LEES	23	PR D107 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LIAO	23	PR D107 112007	L. Liao <i>et al.</i>	
ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)

ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSION	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)

BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAITIS...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAITIS...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAITIS...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAITIS...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also				
EDWARDS	82E	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
LEMOIGNE	82	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BESCH	81	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
GIDAL	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)

SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI-...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)