$T_{b\overline{s}}(5568)^{+}$ 

$$I(J^P) = 1(??)$$

### OMITTED FROM SUMMARY TABLE was $X(5568)^{\pm}$

Seen as a peak in the  $B_s \pi^\pm$  mass spectrum with a significance of more than  $3\sigma$  by ABAZOV 16E and ABAZOV 18A in inclusive  $p\overline{p}$  collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

#### $T_{he}(5568)^{+}$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
$5566.9^{+3.2}_{-3.1}^{+0.6}_{-1.2}$	278	$^{ m 1}$ ABAZOV	18A	D0	$p\overline{p} \rightarrow B_S^0 \pi^{\pm} X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $5567.8 \pm 2.9 ^{+0.9}_{-1.0}$ 

133

<sup>2</sup> ABAZOV 16E D0 
$$p\overline{p} \rightarrow B_s^0 \pi^{\pm} X$$

### $T_{bs}(5568)^{+}$ WIDTH

VALUE (MeV)	<u>EVTS</u>	DOCUMENT ID		TECN	COMMENT
18.6 <sup>+7.9</sup> +3.5 -6.1-3.8	278	<sup>1</sup> ABAZOV	18A	D0	$ \rho \overline{\rho} \to B_{\mathcal{S}} \pi^{\pm} X $

• • • We do not use the following data for averages, fits, limits, etc. • • •

133

 $21.9 \pm 6.4 + 5.0$ 

ABAZOV 16E D0  $p\overline{p} \rightarrow B_s \pi^{\pm} X$ 

## $T_{bs}(5568)^+$ DECAY MODES

	Mode	Fraction $(\Gamma_i/\Gamma)$
Γ <sub>1</sub>	$B_s \pi^+$	seen

# $T_{b\overline{s}}(5568)^+$ BRANCHING RATIOS

$I\left(B_{s}\pi^{\top}\right)/I_{total}$						l <sub>1</sub> /l
VALUE	<b>EVTS</b>	DOCUMENT ID		TECN	COMMENT	
seen	145	$^{ m 1}$ ABAZOV	18A	D0	$ p\overline{p} \rightarrow B_{S}^{0} \pi^{\pm} X $ $ p\overline{p} \rightarrow B_{S}^{0} \pi^{\pm} X $	
seen	133	<sup>2</sup> ABAZOV	16E	D0	$p\overline{p} \rightarrow B_{S}^{0}\pi^{\pm}X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		<sup>3</sup> AABOUD	18L	ATLS	$pp \rightarrow B_s^0 \pi^{\pm} X$	
not seen		<sup>4</sup> AALTONEN	18A	CDF	$pp  ightarrow B_{S}^{0} \pi^{\pm} X$ $p\overline{p}  ightarrow B_{S}^{0} \pi^{\pm} X$	
					3	
https://pdg.lbl.gov		Page 1		Creat	ted: 5/30/2025	07:49

<sup>&</sup>lt;sup>1</sup> From the combined analysis of  $B_s^0 \to J/\psi \phi$  and  $B_s^0 \to D_s^\pm \mu^\mp X$  decays. <sup>2</sup> Assumes  $T_{b\overline{s}}(5568)^\pm \to B_s \pi^\pm$  decay. If  $T_{b\overline{s}}(5568)^\pm \to B_s^* \pi^\pm$  decay is assumed, the mass shifts upward by 49 MeV.

 $<sup>^1</sup>$  From the combined analysis of  $B^0_s \to J/\psi \phi$  and  $B^0_s \to D^\pm_s \mu^\mp X$  decays.

### $T_{b\bar{s}}(5568)^+$ REFERENCES

AABOUD	18L	PRL 120 202007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AALTONEN	18A	PRL 120 202006	T. Aaltonen et al.	` (CDF Collab.)
ABAZOV	18A	PR D97 092004	V.M. Abazov et al.	(D0 Collab.)
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	16AI	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Ábazov <i>et al.</i>	(D0 Collab.)

Created: 5/30/2025 07:49

<sup>&</sup>lt;sup>1</sup>With  $B_{s}$  mesons reconstructed in decays to  $D_{s}^{\pm} \mu^{\mp} X$ .

<sup>&</sup>lt;sup>2</sup> Seen in  $p\overline{p}$  collisions at 1.96 TeV at a rate of  $(8.6\pm1.9\pm1.4)\%$  relative to inclusive  $B_{\mathcal{S}}$  production in the kinematic region  $10 < p_{\mathcal{T}}(B_{\mathcal{S}}) < 30$  GeV/c, with  $B_{\mathcal{S}}$  mesons reconstructed in decays to  $J/\psi\phi$ . An alternative possibility,  $T_{b\overline{\mathcal{S}}}(5568)^{\pm} \to B_{\mathcal{S}}^*\pi^{\pm}$  with a missing  $\gamma$ , could not be ruled out.

 $<sup>^3</sup>$  Not seen in 24.4 fb $^{-1}$  of pp collision data at  $\sqrt{s}=7$  and 8 TeV with  $B_{\rm S}$  mesons reconstructed in decays to  $J/\psi\,\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^\pm\to B_{\rm S}\,\pi^\pm$  relative to inclusive  $B_{\rm S}$  production is less than 1.5% at  $p_T(B_{\rm S})>10$  GeV/c and less than 1.6% at  $p_T(B_{\rm S})>15$  GeV/c at 95% CL.

<sup>&</sup>lt;sup>4</sup> Not seen in 9.6 fb<sup>-1</sup> of  $p\overline{p}$  collision data at  $\sqrt{s}=1.96$  TeV with  $B_{\rm S}$  mesons reconstructed in decays to  $J/\psi\,\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^{\pm}\to B_{\rm S}\,\pi^{\pm}$  relative to inclusive  $B_{\rm S}$  production is less than 6.7% at 95% CL.

 $<sup>^5</sup>$  Not seen in  $19.7~{\rm fb}^{-1}$  of  $p\,p$  collisions data at  $\sqrt{s}=8$  TeV with  $B_{\rm S}$  mesons reconstructed in decays to  $J/\psi\,\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^\pm\to~B_{\rm S}\,\pi^\pm$  relative to inclusive  $B_{\rm S}$  production is less than 1.1% at  $p_T(B_{\rm S})>10~{\rm GeV/c}$  and less than 1.0% at  $p_T(B_{\rm S})>15~{\rm GeV/c}$  at  $95\%{\rm CL}$ .

<sup>&</sup>lt;sup>6</sup> Not seen in 3 fb<sup>-1</sup> of  $p\,p$  collision data at  $\sqrt{s}=7$  and 8 TeV in a scan over the  $T_{b\overline{s}}(5568)$  mass and width, with  $B_s$  mesons reconstructed in decays to  $D_s^-\pi^+$  or  $J/\psi\,\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^\pm\to B_s\,\pi^\pm$  relative to inclusive  $B_s$  production is less than 2.1% at  $p_T(B_s)>10$  GeV/c at 90% CL.