

$T_{c\bar{c}}(4250)^+$

$I^G(J^{PC}) = 1^-(?^?+)$
 I, G, C need confirmation.

OMITTED FROM SUMMARY TABLE
was $X(4250)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by MIZUK 08 in the $\pi^+\chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

$T_{c\bar{c}}(4250)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4248^{+44+180}_{-29-35}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

¹From a Dalitz plot analysis with two Breit-Wigner amplitudes.

$T_{c\bar{c}}(4250)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$177^{+54+316}_{-39-61}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

¹From a Dalitz plot analysis with two Breit-Wigner amplitudes.

$T_{c\bar{c}}(4250)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^+\chi_{c1}(1P)$	seen

$T_{c\bar{c}}(4250)^+$ BRANCHING RATIOS

$\Gamma(\pi^+\chi_{c1}(1P))/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID		TECN	COMMENT	
seen	¹	MIZUK	08	BELL	$\overline{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
not seen	²	LEES	12B	BABR	$B \rightarrow K\pi\chi_{c1}(1P)$
¹ With a product branching fraction measurement of $B(\overline{B}^0 \rightarrow K^-T_{c\overline{c}}(4250)^+) \times B(T_{c\overline{c}}(4250)^+ \rightarrow \pi^+\chi_{c1}(1P)) = (4.0^{+2.3+19.7}_{-0.9-0.5}) \times 10^{-5}$.					
² With a product branching fraction limit of $B(\overline{B}^0 \rightarrow T_{c\overline{c}}(4250)^+K^-) \times B(T_{c\overline{c}}(4250)^+ \rightarrow \chi_{c1}\pi^+) < 4.0 \times 10^{-5}$ at 90% CL.					

$T_{c\bar{c}}(4250)^+$ REFERENCES

LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)