

$\chi_{c1}(1P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on “Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ ”
before the $\chi_{c0}(1P)$ Listings.

 $\chi_{c1}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3510.67 $\pm$ 0.05 OUR AVERAGE		Error includes scale factor of 1.2.		
3509.84 $\pm$ 0.69 $\pm$ 0.64	2.8k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+ (K_S^0 K \pi)$
3508.4 $\pm$ 1.9 $\pm$ 0.7	460	¹ AAIJ	17BB LHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+ K^-)X$
3510.71 $\pm$ 0.04 $\pm$ 0.09	4.8k	² AAIJ	17BI LHCb	$\chi_{c1} \rightarrow J/\psi \mu^+ \mu^-$
3510.30 $\pm$ 0.14 $\pm$ 0.16		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3510.719 $\pm$ 0.051 $\pm$ 0.019		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
3509.4 $\pm$ 0.9		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.60 $\pm$ 0.087 $\pm$ 0.019	513	³ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
3511.3 $\pm$ 0.4 $\pm$ 0.4	30	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
3512.3 $\pm$ 0.3 $\pm$ 4.0		⁴ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 $\pm$ 1.7	91	⁵ LEMOIGNE	82 GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3510.4 $\pm$ 0.6		OREGLIA	82 CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
3510.1 $\pm$ 1.1	254	⁶ HIMEL	80 MRK2	$e^+ e^- \rightarrow J/\psi 2\gamma$
3509 $\pm$ 11	21	BRANDELIK	79B DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
3507 $\pm$ 3		⁶ BARTEL	78B CNTR	$e^+ e^- \rightarrow J/\psi 2\gamma$
3505.0 $\pm$ 4 $\pm$ 4		^{6,7} TANENBAUM	78 MRK1	$e^+ e^-$
3513 $\pm$ 7	367	⁶ BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$
3500 $\pm$ 10	40	TANENBAUM	75 MRK1	Hadrons γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c1}(1P)$ fixed to the PDG 16 value.

² AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

³ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

⁴ Using mass of $\psi(2S) = 3686.0$ MeV.

⁵ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁶ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁷ From a simultaneous fit to radiative and hadronic decay channels.

 $\chi_{c1}(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 $\pm$ 0.04 OUR FIT			Error includes scale factor of 1.1.		
0.88 $\pm$ 0.05 OUR AVERAGE					
1.39 $^{+0.40}_{-0.38}$ $^{+0.26}_{-0.77}$			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
0.876 $\pm$ 0.045 $\pm$ 0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
0.87 $\pm$ 0.11 $\pm$ 0.08	513	¹ ARMSTRONG	92 E760		$\bar{p}p \rightarrow e^+ e^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	95	BAGLIN	86B	SPEC	$\bar{p}p \rightarrow e^+e^-X$
<3.8	90	GAISER	86	CBAL	$\psi(2S) \rightarrow \gamma X$

¹ Recalculated by ANDREOTTI 05A.

$\chi_{c1}(1P)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1	hadrons		
Γ_2	e^+e^-	$(1.4^{+1.5}_{-1.0}) \times 10^{-7}$	
Hadronic decays			
Γ_3	$3(\pi^+\pi^-)$	$(1.04 \pm 0.16) \%$	S=4.6
Γ_4	$2(\pi^+\pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$	
Γ_5	$\pi^+\pi^-\pi^0\pi^0$	$(1.19 \pm 0.15) \%$	
Γ_6	$\rho^+\pi^-\pi^0 + \text{c.c.}$	$(1.45 \pm 0.24) \%$	
Γ_7	$\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$	
Γ_8	$4\pi^0$	$(5.4 \pm 0.8) \times 10^{-4}$	
Γ_9	$\pi^+\pi^-K^+K^-$	$(4.5 \pm 1.0) \times 10^{-3}$	
Γ_{10}	$K^+K^-\pi^0\pi^0$	$(1.12 \pm 0.27) \times 10^{-3}$	
Γ_{11}	$K^+K^-\pi^+\pi^-\pi^0$	$(1.15 \pm 0.13) \%$	
Γ_{12}	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$	
Γ_{13}	$K^\pm \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(8.6 \pm 1.4) \times 10^{-3}$	
Γ_{14}	$\rho^- K^+ \bar{K}^0 + \text{c.c.}$	$(5.0 \pm 1.2) \times 10^{-3}$	
Γ_{15}	$K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{16}	$K^+K^-\eta\pi^0$	$(1.12 \pm 0.34) \times 10^{-3}$	
Γ_{17}	$\pi^+\pi^-K_S^0 K_S^0$	$(6.9 \pm 2.9) \times 10^{-4}$	
Γ_{18}	$K^+K^-\eta$	$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{19}	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$(7.0 \pm 0.6) \times 10^{-3}$	S=1.1
Γ_{20}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.03 \pm 0.15) \times 10^{-3}$	
Γ_{21}	$K^*(892)^+ K^- + \text{c.c.}$	$(1.21 \pm 0.23) \times 10^{-3}$	
Γ_{22}	$K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K^+ \pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%
Γ_{23}	$K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K_S^0 K^+ \pi^- + \text{c.c.}$	$< 2.1 \times 10^{-3}$	CL=90%
Γ_{24}	$K^+K^-\pi^0$	$(1.81 \pm 0.24) \times 10^{-3}$	
Γ_{25}	$\eta\pi^+\pi^-$	$(4.62 \pm 0.24) \times 10^{-3}$	
Γ_{26}	$a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(3.2 \pm 0.4) \times 10^{-3}$	S=2.1
Γ_{27}	$a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(1.76 \pm 0.24) \times 10^{-4}$	
Γ_{28}	$a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(4.6 \pm 0.7) \times 10^{-5}$	
Γ_{29}	$f_2(1270)\eta \rightarrow \eta\pi^+\pi^-$	$(3.5 \pm 0.6) \times 10^{-4}$	

Γ_{30}	$f_4(2050)\eta \rightarrow \eta\pi^+\pi^-$	$(2.5 \pm 0.9) \times 10^{-5}$	
Γ_{31}	$\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 5 \times 10^{-5}$	CL=90%
Γ_{32}	$\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{33}	$\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 8 \times 10^{-6}$	CL=90%
Γ_{34}	$f_2(1270)\eta$	$(6.7 \pm 1.1) \times 10^{-4}$	
Γ_{35}	$\pi^+\pi^-\eta'$	$(2.2 \pm 0.4) \times 10^{-3}$	
Γ_{36}	$K^+K^-\eta'(958)$	$(8.8 \pm 0.9) \times 10^{-4}$	
Γ_{37}	$K_0^*(1430)^+K^- + \text{c.c.}$	$(6.4^{+2.2}_{-2.8}) \times 10^{-4}$	
Γ_{38}	$f_0(980)\eta'(958)$	$(1.6^{+1.4}_{-0.7}) \times 10^{-4}$	
Γ_{39}	$f_0(1710)\eta'(958)$	$(7^{+7}_{-5}) \times 10^{-5}$	
Γ_{40}	$f'_2(1525)\eta'(958)$	$(9 \pm 6) \times 10^{-5}$	
Γ_{41}	$K_2^*(1430)^+K^- + \text{c.c.}$	$(1.61 \pm 0.31) \times 10^{-3}$	
Γ_{42}	$K_2^*(1430)\bar{K}^0 + \text{c.c.}$	$(1.17 \pm 0.20) \times 10^{-3}$	
Γ_{43}	$\pi^0 f_0(980) \rightarrow \pi^0\pi^+\pi^-$	$(3.5 \pm 0.9) \times 10^{-7}$	
Γ_{44}	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$	
Γ_{45}	$K^*(892)^0\bar{K}^*(892)^0$	$(1.4 \pm 0.4) \times 10^{-3}$	
Γ_{46}	$K^+K^-K_S^0K_S^0$	$< 4 \times 10^{-4}$	CL=90%
Γ_{47}	$K_S^0K_S^0K_S^0K_S^0$	$(3.5 \pm 1.0) \times 10^{-5}$	
Γ_{48}	$K^+K^-K^+K^-$	$(5.4 \pm 1.1) \times 10^{-4}$	
Γ_{49}	$K^+K^-\phi$	$(4.1 \pm 1.5) \times 10^{-4}$	
Γ_{50}	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	$(3.3 \pm 0.5) \times 10^{-3}$	
Γ_{51}	$K^+K^-\pi^0\phi$	$(1.62 \pm 0.30) \times 10^{-3}$	
Γ_{52}	$3(K^+K^-)$	$(4.2 \pm 1.1) \times 10^{-6}$	
Γ_{53}	$\phi\pi^+\pi^-\pi^0$	$(7.5 \pm 1.0) \times 10^{-4}$	
Γ_{54}	$\omega\omega$	$(5.7 \pm 0.7) \times 10^{-4}$	
Γ_{55}	ωK^+K^-	$(7.8 \pm 0.9) \times 10^{-4}$	
Γ_{56}	$\omega\phi$	$(2.7 \pm 0.4) \times 10^{-5}$	
Γ_{57}	$\phi\phi$	$(4.26 \pm 0.21) \times 10^{-4}$	
Γ_{58}	$\phi\phi\eta$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{59}	$p\bar{p}$	$(7.6 \pm 0.4) \times 10^{-5}$	S=1.2
Γ_{60}	$p\bar{p}\pi^0$	$(1.55 \pm 0.18) \times 10^{-4}$	
Γ_{61}	$p\bar{p}\eta$	$(1.45 \pm 0.25) \times 10^{-4}$	
Γ_{62}	$p\bar{p}\omega$	$(2.12 \pm 0.31) \times 10^{-4}$	
Γ_{63}	$p\bar{p}\phi$	$< 1.7 \times 10^{-5}$	CL=90%
Γ_{64}	$p\bar{p}\pi^+\pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$	
Γ_{65}	$p\bar{p}\pi^0\pi^0$	$< 5 \times 10^{-4}$	CL=90%
Γ_{66}	$p\bar{p}K^+K^- (\text{non-resonant})$	$(1.27 \pm 0.22) \times 10^{-4}$	
Γ_{67}	$p\bar{p}K_S^0K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%
Γ_{68}	$p\bar{p}K_S^0K^- \pi^+ + \text{c.c.}$	$(4.2 \pm 0.5) \times 10^{-5}$	
Γ_{69}	$p\bar{n}\pi^-$	$(3.8 \pm 0.5) \times 10^{-4}$	
Γ_{70}	$\bar{p}n\pi^+$	$(3.9 \pm 0.5) \times 10^{-4}$	
Γ_{71}	$p\bar{n}\pi^-\pi^0$	$(1.03 \pm 0.12) \times 10^{-3}$	

Γ_{72}	$\bar{p}n\pi^+\pi^0$	$(1.01 \pm 0.12) \times 10^{-3}$	
Γ_{73}	$\Lambda\bar{\Lambda}$	$(1.27 \pm 0.09) \times 10^{-4}$	S=1.1
Γ_{74}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.9 \pm 0.5) \times 10^{-4}$	
Γ_{75}	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$(2.5 \pm 0.6) \times 10^{-4}$	
Γ_{76}	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{77}	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{78}	$\Lambda\bar{\Lambda}\eta$	$(5.9 \pm 1.5) \times 10^{-5}$	
Γ_{79}	$\Lambda\bar{\Lambda}\omega$	$(1.01 \pm 0.16) \times 10^{-4}$	
Γ_{80}	$\Lambda\bar{\Lambda}\phi$	$(6.0 \pm 1.0) \times 10^{-5}$	
Γ_{81}	$K^+\bar{p}\Lambda + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-4}$	S=1.2
Γ_{82}	$nK_S^0\bar{\Lambda} + \text{c.c.}$	$(1.66 \pm 0.17) \times 10^{-4}$	
Γ_{83}	$\bar{p}\Lambda(1520)K_S^0\pi^+ + \text{c.c.}$	$(4.1 \pm 0.9) \times 10^{-5}$	
Γ_{84}	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	$(4.9 \pm 0.7) \times 10^{-4}$	
Γ_{85}	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(1.7 \pm 0.4) \times 10^{-4}$	
Γ_{86}	$\Lambda(1520)\bar{\Lambda}(1520)$	$< 9 \times 10^{-5}$	CL=90%
Γ_{87}	$\Sigma^0\bar{\Sigma}^0$	$(4.2 \pm 0.6) \times 10^{-5}$	
Γ_{88}	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	$(1.53 \pm 0.12) \times 10^{-4}$	
Γ_{89}	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	$(1.46 \pm 0.10) \times 10^{-4}$	
Γ_{90}	$\Sigma^+\bar{\Sigma}^-$	$(3.6 \pm 0.7) \times 10^{-5}$	
Γ_{91}	$\Sigma^+\bar{\Sigma}^-\eta$	$(5.1 \pm 1.4) \times 10^{-5}$	
Γ_{92}	$\Sigma^-\bar{\Sigma}^+$	$(5.7 \pm 1.5) \times 10^{-5}$	
Γ_{93}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 9 \times 10^{-5}$	CL=90%
Γ_{94}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90%
Γ_{95}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.35 \pm 0.24) \times 10^{-4}$	
Γ_{96}	$\Xi^0\bar{\Xi}^0$	$(7.5 \pm 1.3) \times 10^{-5}$	
Γ_{97}	$\Xi^-\bar{\Xi}^+$	$(6.0 \pm 0.6) \times 10^{-5}$	
Γ_{98}	$\Omega^-\bar{\Omega}^+$	$(1.49 \pm 0.25) \times 10^{-5}$	
Γ_{99}	$\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$	
Γ_{100}	$K_S^0K_S^0$	$< 6 \times 10^{-5}$	CL=90%
Γ_{101}	$\eta_c\pi^+\pi^-$	$< 3.2 \times 10^{-3}$	CL=90%

Radiative decays

Γ_{102}	$\gamma J/\psi(1S)$	$(34.3 \pm 1.3) \%$	S=1.3
Γ_{103}	$\gamma\rho^0$	$(2.16 \pm 0.17) \times 10^{-4}$	
Γ_{104}	$\gamma\omega$	$(6.8 \pm 0.8) \times 10^{-5}$	
Γ_{105}	$\gamma\phi$	$(2.4 \pm 0.5) \times 10^{-5}$	
Γ_{106}	$\gamma\gamma$	$< 6.3 \times 10^{-6}$	CL=90%
Γ_{107}	$e^+e^- J/\psi(1S)$	$(3.46 \pm 0.24) \times 10^{-3}$	
Γ_{108}	$\mu^+\mu^- J/\psi(1S)$	$(2.33 \pm 0.29) \times 10^{-4}$	

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 88 branching ratios uses 255 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 393.1$ for 206 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{48}	4				
x_{59}	−1	0			
x_{73}	12	5	−1		
x_{102}	20	9	−25	23	
Γ	−9	−4	−60	−11	−31
	x_{19}	x_{48}	x_{59}	x_{73}	x_{102}

$\chi_{c1}(1P)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					Γ_2
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.12^{+0.13}_{-0.08}$	250	¹ ABLIKIM	22AF BES3	$e^+e^- \rightarrow \chi_{c1} \rightarrow \gamma J/\psi$	
¹ Assuming $\Gamma(\chi_{c1} \rightarrow \gamma J/\psi) = 0.28$ MeV.					

$\chi_{c1}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$					$\Gamma_{59} \Gamma_{102} / \Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
21.9 ± 0.8 OUR FIT					
21.4 ± 0.9 OUR AVERAGE					
21.5 ± 0.5 ± 0.8		¹ ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+e^-\gamma$	
21.4 ± 1.5 ± 2.2		^{1,2} ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+e^-\gamma$	
19.9 ^{+4.4} _{−4.0}		¹ BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+e^-X$	
¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.					
² Recalculated by ANDREOTTI 05A.					

$\chi_{c1}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-)) / \Gamma_{\text{total}}$					Γ_3 / Γ
VALUE (units 10 ^{−3})	EVTS	DOCUMENT ID	TECN	COMMENT	
10.4 ± 1.6 OUR AVERAGE		Error includes scale factor of 4.6.			
10.92 ± 0.23 ± 0.30	84K	¹ ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$	
5.4 ± 0.7 ± 0.9		² BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$	

$16.0 \pm 5.9 \pm 0.8$ 2 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

1 ABLIKIM 22Q reports $(1.092 \pm 0.004 \pm 0.035) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.6 $\pm$ 2.6	OUR EVALUATION	Treating systematic error as correlated.		
8 $\pm$ 4	OUR AVERAGE	Error includes scale factor of 1.5.		
4.6 $\pm$ 2.1 $\pm$ 2.6	1 BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
12.5 $\pm$ 4.2 $\pm$ 0.6	1 TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.85 $\pm$ 0.26 $\pm$ 0.19 670 2 ABLIKIM 24BT BES3 $\psi(2S) \rightarrow \gamma \chi_{c1}$

1 Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

2 ABLIKIM 24BT reports $(0.685 \pm 0.001 \pm 0.031) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 $\pm$ 0.15 $\pm$ 0.03	604.7	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

1 HE 08B reports $1.28 \pm 0.06 \pm 0.15 \pm 0.08\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.45 $\pm$ 0.24 $\pm$ 0.04	712.3	1,2 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

1 HE 08B reports $1.56 \pm 0.13 \pm 0.22 \pm 0.10\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.9 $\pm$ 3.5	1 TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ **Γ_8/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.4 \pm 0.8 \pm 0.1$	608	¹ ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 11A reports $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ **Γ_9/Γ**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.5 ± 1.0 OUR EVALUATION	Treating systematic error as correlated.		
4.5 ± 0.9 OUR AVERAGE			

$4.2 \pm 0.4 \pm 0.9$ ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c1}$

$7.3 \pm 3.0 \pm 0.4$ ¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.12 \pm 0.27 \pm 0.03$	45.1	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $(0.12 \pm 0.02 \pm 0.02 \pm 0.01) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$11.46 \pm 0.12 \pm 1.29$	12k	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.52 \pm 0.11 \pm 0.79$	5.1k	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.86 \pm 0.13 \pm 0.02$	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $0.92 \pm 0.09 \pm 0.11 \pm 0.06\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.50±0.12±0.01	141.3	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.54 \pm 0.11 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.23±0.06±0.01	141.3	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.25 \pm 0.06 \pm 0.03 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.112±0.034±0.003	141.3	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.12 \pm 0.03 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.9±2.9±0.2	19.8 ± 7.7	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$ which we divide by our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 1.0 ± 0.1		¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.30 ± 0.05	8690	² ABLIKIM	24BWBES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-------------	------	----------------------	----------	---

¹ ATHAR 07 reports $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² No systematic error reported.

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03 ± 0.15 OUR AVERAGE				
$1.04 \pm 0.13 \pm 0.10$	262	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$
$0.98 \pm 0.37 \pm 0.03$	22	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ AAIJ 23AH reports $(1.04 \pm 0.13 \pm 0.04 \pm 0.09) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

² ABLIKIM 06R reports $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ± 0.23 OUR AVERAGE				
$1.18 \pm 0.17 \pm 0.17$	288	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$
$1.43 \pm 0.65 \pm 0.04$	27	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ AAIJ 23AH reports $(1.18 \pm 0.17 \pm 0.14 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

² ABLIKIM 06R reports $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-4}$	90	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$$\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{23}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-3}$	90	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $< 2.4 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{24}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.81 \pm 0.24 \pm 0.05$	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{25}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.62±0.24 OUR AVERAGE					
4.58±0.23±0.13		^{1,2} ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$
4.7 ±0.5 ±0.1		³ ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.3 ±0.9 ±0.1	222	⁴ ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(4.67 \pm 0.03 \pm 0.23 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ATHAR 07 reports $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06R reports $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{26}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.2 $\pm$0.4 OUR AVERAGE Error includes scale factor of 2.1.					
3.33 $\pm$ 0.19 $\pm$ 0.09		^{1,2} ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1.79 $\pm$ 0.63 $\pm$ 0.05	58	³ ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(3.40 \pm 0.03 \pm 0.19 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06R reports $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{27}/Γ
VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
0.176$\pm$0.023$\pm$0.005	1,2	ABLIKIM	17K	BES3
				$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(0.18 \pm 0.01 \pm 0.02 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 0.7 \pm 0.1$	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(4.7 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f_2(1270) \eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 0.6 \pm 0.1$	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(0.36 \pm 0.01 \pm 0.06 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270) \eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f_4(2050) \eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.9 \pm 0.1$	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(2.6 \pm 0.4 \pm 0.8 \pm 0.1) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_4(2050) \eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5 \times 10^{-5}$	90	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 4.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-5}$	90	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 1.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-6}$	90	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.67 ± 0.11 OUR AVERAGE				
$0.63 \pm 0.11 \pm 0.02$		1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
$2.7 \pm 0.8 \pm 0.1$	53	³ ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 17K reports $(6.4 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From an amplitude analysis using an isobar model.

³ ABLIKIM 06R reports $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.2 \pm 0.4 \pm 0.1$	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.75 ± 0.87	310	¹ ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

$$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{41}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.61 \pm 0.19 \pm 0.24$	351	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ AAIJ 23AH reports $(1.61 \pm 0.19 \pm 0.19 \pm 0.14) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $\text{B}(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

$$\Gamma(K_2^*(1430) \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.17 \pm 0.16 \pm 0.11$	278	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ AAIJ 23AH reports $(1.17 \pm 0.16 \pm 0.05 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_2^*(1430) \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $\text{B}(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

$$\Gamma(K_0^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$6.41 \pm 0.57 + 2.09 - 2.71$	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Normalized to $\text{B}(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$$\Gamma(f_0(980) \eta'(958))/\Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.65 \pm 0.47 + 1.32 - 0.56$	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Normalized to $\text{B}(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$$\Gamma(f_0(1710) \eta'(958))/\Gamma_{\text{total}} \quad \Gamma_{39}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.71 \pm 0.22 + 0.68 - 0.48$	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Normalized to $\text{B}(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$$\Gamma(f_2'(1525) \eta'(958))/\Gamma_{\text{total}} \quad \Gamma_{40}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.92 \pm 0.23 + 0.55 - 0.51$	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Normalized to $\text{B}(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$$\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{43}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.09		ABLIKIM	18D BES3	$\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 ¹ ABLIKIM 11D BES3 $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

¹ ABLIKIM 11D reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ < 6.0×10^{-7} which we divide by our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
32 ± 21	¹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.44 ± 0.36 ± 0.04	28.4 ± 5.5	^{1,2} ABLIKIM	04H BES	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

¹ ABLIKIM 04H reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 4 × 10⁻⁴	90	3.2 ± 2.4	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.10 ± 0.01	22	¹ ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4 K_S^0$

¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (3.4 \pm 0.9 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.15 ± 0.01	17	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2 K^+ 2 K^-$

¹ ABLIKIM 06T reports $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.27 ± 0.28 ± 0.46	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.62 ± 0.12 ± 0.28	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.75 \pm 0.06 \pm 0.08$	373	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 0.7 \pm 0.2$	597	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons

¹ ABLIKIM 11K reports $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.78 \pm 0.04 \pm 0.08$	628	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.27 \pm 0.04 \pm 0.01$	105	¹ ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.21 \pm 0.06 \pm 0.01$ 15 ^{2,3} ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ = $(2.67 \pm 0.31 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 11K reports $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by ABLIKIM 19J.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.26 \pm 0.17 \pm 0.12$	1529	^{1,2} ABLIKIM	23N BES3	$\psi(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.2 \pm 0.5 \pm 0.1$ 366 ³ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

² ABLIKIM 23N reports $(4.26 \pm 0.13 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 11K reports $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$ **Γ_{58}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.0 \pm 0.5 \pm 0.1$	83.6	¹ ABLIKIM	20B	BES3 $\psi(2S) \rightarrow \gamma\phi\phi\eta$

¹ ABLIKIM 20B reports $(2.96 \pm 0.43 \pm 0.22) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ **Γ_{60}/Γ**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.155 ± 0.018 OUR AVERAGE			
$0.163 \pm 0.019 \pm 0.004$	¹ ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$
$0.112 \pm 0.047 \pm 0.003$	² ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ **Γ_{61}/Γ**

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$0.145 \pm 0.024 \pm 0.004$		¹ ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$
< 0.15	90	² ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ONYISI 10 reports $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $< 0.16 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ **Γ_{62}/Γ**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.212 \pm 0.030 \pm 0.006$	¹ ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

¹ ONYISI 10 reports $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 × 10⁻⁵	90	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $< 1.82 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
0.50±0.19 OUR EVALUATION	Treating systematic error as correlated.		
0.50±0.19 OUR AVERAGE			

0.46±0.12±0.15	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
1.08±0.77±0.05	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5 × 10⁻⁴	90	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $< 0.05 \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
1.27±0.22±0.04	82 ± 9	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$.

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±0.5±0.1	1412	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$

Γ_{70}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 0.5 \pm 0.1$	1625	¹ ABLIKIM 12J	BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{71}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$10.3 \pm 1.1 \pm 0.3$	1082	¹ ABLIKIM 12J	BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$

Γ_{72}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$10.1 \pm 1.1 \pm 0.3$	1261	¹ ABLIKIM 12J	BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{74}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$29 \pm 5 \pm 1$		105	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150	90	² ABLIKIM 06D	BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
------	----	--------------------------	------	--

¹ ABLIKIM 12I reports $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$

Γ_{75}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$25 \pm 6 \pm 1$	13	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

¹ ABLIKIM 12I reports $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow$

$\gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$
¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.				

$\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<13	90	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$
¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.				

$\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.2.			
$9.2^{+2.8}_{-2.4} \pm 0.4$	24	¹ LU	19	BELL $B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
$4.2 \pm 0.4 \pm 0.1$	3k	^{2,3} ABLIKIM	13D	BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
$3.1 \pm 0.9 \pm 0.1$		⁴ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ LU 19 reports $(9.15^{+2.63}_{-2.25} \pm 0.86) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(1P)K^+)]$ assuming $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.79 \pm 0.23) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.74 \pm 0.22) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

⁴ ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.66 \pm 0.16 \pm 0.05$	399	¹ ABLIKIM 21AV	BES3	$\psi(2S) \rightarrow \gamma nK_S^0\bar{\Lambda} + \text{c.c.}$

¹ ABLIKIM 21AV reports $(1.66 \pm 0.12 \pm 0.12) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow nK_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0975 \pm 0.0024$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$.

$\Gamma(K^*(892)^+ \bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{84}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.9 \pm 0.7 \pm 0.1$	328	¹ ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p}\Lambda$

¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ \bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (4.8 \pm 0.5 \pm 0.4) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{85}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.71 \pm 0.44 \pm 0.05$	48 ± 10	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}$

Γ_{86}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-5}$	90	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $< 1.00 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

Γ_{87}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 0.6 \pm 0.1$		103	¹ ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
< 4	90	3.8 ± 2.5	³ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (0.41 \pm 0.05 \pm 0.03) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13H reports $< 0.62 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

³ NAIK 08 reports $< 0.44 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 0.6 \pm 0.1$		59	¹ ABLIKIM	18v BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
< 6	90	4.3 ± 2.3	³ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

¹ ABLIKIM 18v reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ = $(0.35 \pm 0.06 \pm 0.02) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13H reports $< 0.87 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

³ NAIK 08 reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.1 \pm 1.4 \pm 0.1$	36	¹ ABLIKIM	24CA BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}(1P)$

¹ ABLIKIM 24CA reports $(5.10 \pm 1.21 \pm 0.67) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 1.5 \pm 0.2$	214	¹ ABLIKIM	20i BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

¹ ABLIKIM 20i reports $(5.7 \pm 1.4 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-5}$	90	¹ ABLIKIM	12i BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

¹ ABLIKIM 12i reports $< 10 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5 \times 10^{-5}$	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
¹ ABLIKIM 12I reports $< 5.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.				

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.35 \pm 0.24 \pm 0.04$	49	¹ ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$
¹ ABLIKIM 15I reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.5 \pm 1.2 \pm 0.2$		325	¹ ABLIKIM	22O BES3	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6	90	1.7 ± 2.4	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
¹ ABLIKIM 22O reports $(0.75 \pm 0.11 \pm 0.06) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.60 ± 0.06 OUR AVERAGE					
$0.58 \pm 0.06 \pm 0.02$		692	¹ ABLIKIM	22O BES3	$\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$
$0.80 \pm 0.21 \pm 0.02$	16.4 ± 4.3		² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^+ \Xi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.4	90		³ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
¹ ABLIKIM 22O reports $(0.58 \pm 0.04 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
³ Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma) (9.1 \pm 0.6)\%$.					

$\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.49 \pm 0.23 \pm 0.10$	277	ABLIKIM	23T BES3	$\chi_{cJ} \rightarrow \Omega^- \bar{\Omega}^+$

$[\Gamma(\pi^+ \pi^-) + \Gamma(K^+ K^-)]/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 21 \times 10^{-4}$		¹ FELDMAN	77 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 38 \times 10^{-4}$	90	¹ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	----	------------------------	----------	---

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6 \times 10^{-5}$	90	¹ ABLIKIM	05O BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

¹ ABLIKIM 05O reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$
 $< 0.6 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.2 \times 10^{-3}$	90	^{1,2} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 4.4 \times 10^{-3}$	90	^{1,3} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
------------------------	----	------------------------	----------	---

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

² Using the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

³ Using the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

————— RADIATIVE DECAYS —————

$\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
216 ± 17 OUR AVERAGE				
$215 \pm 22 \pm 6$	432 ± 25	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
$217 \pm 24 \pm 6$	186 ± 15	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$

¹ ABLIKIM 11E reports $(228 \pm 13 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $(243 \pm 19 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
68 ± 8 OUR AVERAGE				
$66 \pm 9 \pm 2$	136 ± 14	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
$74 \pm 17 \pm 2$	39 ± 7	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

¹ ABLIKIM 11E reports $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $(83 \pm 15 \pm 12) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$24 \pm 5 \pm 1$		43 ± 9	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 23	90	5.2 ± 3.1	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$

¹ ABLIKIM 11E reports $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $< 26 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-6}$	90	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 3.5 \times 10^{-5}$	90	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$
$< 150 \times 10^{-5}$	90	¹ YAMADA	77 DASP	$e^+e^- \rightarrow 3\gamma$

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

$\Gamma(e^+e^-J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.65 \pm 0.23 \pm 0.10$	1.9k	^{1,2} ABLIKIM	17I BES3	$\psi(2S) \rightarrow \gamma e^+e^-J/\psi$

¹ ABLIKIM 17I reports $(3.73 \pm 0.09 \pm 0.25) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow e^+e^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent from other measurements reported by ABLIKIM 17I

$$\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S)) \quad \Gamma_{107}/\Gamma_{102}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.1±0.3±0.5	1.9k	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^- \gamma J/\psi$
¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.				

$$\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S)) \quad \Gamma_{108}/\Gamma_{107}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
6.73±0.51±0.50	222	ABLIKIM	19Z BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$

$\chi_{c1}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$$\Gamma(\chi_{c1}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.68±0.01±0.25	670	¹ ABLIKIM	24BT BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
¹ Calculated by us. The value given here is derived from the value of $B(\chi_{c1} \rightarrow 2(\pi^+\pi^-))$ reported in ABLIKIM 24BT using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.75 \pm 0.24)\%$ [PDG 22].				

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.8±0.6 OUR FIT	Error includes scale factor of 1.1.		
7.2±0.6 OUR AVERAGE			
7.3±0.5±0.5	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
7.0±0.5±0.9	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K^0 K^+ \pi^- + \text{c.c.})$ reported by ATHAR 07 was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$.			
² Calculated by us. ABLIKIM 06R reports $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$. We use $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$.			

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
19.6±1.6 OUR FIT	Error includes scale factor of 1.1.		
13.2±2.4±3.2	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].			

$$\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \quad \Gamma_{48}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.53±0.11 OUR FIT				
0.61±0.11±0.08	54	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{48} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}{\Gamma_{182}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.52±0.31 OUR FIT			
1.13±0.40±0.29	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow 3(K^+ K^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{52} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$


VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
4.1±0.9±0.5	24.9 ± 5.1	¹ ABLIKIM	24P BES3	$e^+ e^- \rightarrow \psi(2S)$

¹ Systematic error derived by us, based on the text.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{59} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
7.4±0.4 OUR FIT	Error includes scale factor of 1.3.			
7.8±0.6 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
7.9±0.4±0.3	453	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p \bar{p}$
8.2±0.7±0.4	141 ± 13	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
4.8 ^{+1.4} _{-1.3} ± 0.6	18.2 ^{+5.5} _{-4.9}	BAI	04F BES	$\psi(2S) \rightarrow \gamma \chi_{c1}(1P) \rightarrow \gamma p \bar{p}$

WEIGHTED AVERAGE
7.8±0.6 (Error scaled by 1.4)



$$\Gamma(\chi_{c1}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} \text{ (units)}$$

10^{-6})

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{59}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.13±0.13 OUR FIT	Error includes scale factor of 1.3.		
1.1 ±1.0	¹ BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\bar{p}p$

¹ Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{68}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.05±0.24±0.39	396	ABLIKIM	24BX BES3	$\psi(2S) \rightarrow \gamma\chi_{c1}$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{73}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±0.9 OUR FIT	Error includes scale factor of 1.1.			
12.3±0.9 OUR AVERAGE	Error includes scale factor of 1.2.			

12.8±0.6±0.6	528	ABLIKIM	21L BES3	$\psi(2S) \rightarrow \gamma p\pi^- \bar{p}\pi^+$
10.5±1.6±0.6	46	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.2±1.0±0.9	136	^{2,3} ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
--------------	-----	------------------------	----------	---

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

² Superseded by ABLIKIM 21L

³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ from a measurement of $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c1})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{73}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.58±0.25 OUR FIT	Error includes scale factor of 1.1.			
7.1 $\begin{smallmatrix} +2.8 \\ -2.4 \end{smallmatrix} \pm 1.3$	$9.0 \begin{smallmatrix} +3.5 \\ -3.1 \end{smallmatrix}$	¹ BAI	03E BES	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ BAI 03E reports $[B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33 \begin{smallmatrix} +0.52 \\ -0.46 \end{smallmatrix} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda \bar{\Lambda} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{78} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.72 \pm 1.34 \pm 0.65$	21	ABLIKIM	22AO BES3	$\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+ \gamma \gamma$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda \bar{\Lambda} \omega) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{79} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.8 \pm 1.0 \pm 1.1$	202 ± 20	¹ ABLIKIM	24BE BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Calculated by us. The authors report $B(\chi_{c1} \rightarrow \Lambda \bar{\Lambda} \omega)$ obtained from a product using PDG 22 value of $B(\psi(2S) \rightarrow \gamma \chi_{c1})$.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda \bar{\Lambda} \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{80} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.87 \pm 0.39$	51.6	ABLIKIM	24AC BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{p} \Lambda(1520) K_S^0 \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{83} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.96^{+0.77}_{-0.74} \pm 0.50$	88	ABLIKIM	24BX BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{88} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.49 \pm 0.09 \pm 0.07$	258	¹ ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c1} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (1.53 \pm 0.10 \pm 0.08) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.75 \pm 0.24)\%$ and other branching fractions from PDG 18.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{89} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma \psi(2S)}$$


VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.07 \pm 0.06$	493	¹ ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_{c1} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (1.46 \pm 0.07 \pm 0.07) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}^0) = (9.75 \pm 0.24)\%$ and other branching fractions from PDG 20.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{102}}{\Gamma} \times \frac{\Gamma_{182}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.34 ± 0.10 OUR FIT	Error includes scale factor of 1.7.			
3.24 ± 0.16 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.			
3.518 ± 0.010 ± 0.120	143k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
3.442 ± 0.010 ± 0.132	1.9M	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
2.81 ± 0.05 ± 0.23	13k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
2.56 ± 0.12 ± 0.20		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
2.78 ± 0.30		² OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.2 ± 0.5		³ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.9 ± 0.5		³ BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c1}$
5.0 ± 1.5		⁴ BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$
2.8 ± 0.9		² WHITAKER	76 MRK1	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.377 ± 0.009 ± 0.183	142k	⁵ ABLIKIM	12O BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.56 ± 0.03 ± 0.12	24.9k	⁶ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.44 ± 0.06 ± 0.13	3.7k	⁷ ADAM	05A CLEO	Repl. by MENDEZ 08

WEIGHTED AVERAGE
3.24 ± 0.16 (Error scaled by 2.1)



$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))}{\Gamma_{\text{total}}} \quad (\text{units } 10^{-2})$$

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{102}}{\Gamma} \times \frac{\Gamma_{182}^{\psi(2S)}}{\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.62±0.29 OUR FIT	Error includes scale factor of 1.7.			
10.15±0.28 OUR AVERAGE				
10.17±0.07±0.27	24.9k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma \chi_{c1}$
12.6 ±0.3 ±3.8	3k	¹ ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
8.5 ±2.1		² HIMEL	80	MRK2 $\psi(2S) \rightarrow \gamma \chi_{c1}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.24±0.17±0.23	3.7k	³ ADAM	05A	CLEO Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.


³ Not independent from other values reported by ADAM 05A.

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

$a_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
−6.7 ±0.9 OUR AVERAGE	Error includes scale factor of 2.6. See the ideogram below.			
−7.40±0.33±0.34	164k	¹ ABLIKIM	17N	BES3 $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
−6.26±0.63±0.24	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.2 ±3.2 ±0.4	2090	AMBROGIANI	02	E835 $p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi \gamma$
−0.2 ^{+0.8} _{−2.0}	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

¹ Correlated with b_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.



MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma \chi_{c1}(1S)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.4 OUR AVERAGE				
$2.29 \pm 0.39 \pm 0.27$	164k	¹ ABLIKIM	17N	BES3 $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$2.76 \pm 0.73 \pm 0.23$	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$7.7^{+5.0}_{-4.5}$	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Correlated with a_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS
 $\psi(2S) \rightarrow \gamma \chi_{c1}(1S)$ and $\chi_{c1} \rightarrow \gamma J/\psi(1S)$

a_2/b_2 Magnetic quadrupole transition amplitude ratio

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-2.27^{+0.57}_{-0.99}$	39k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Statistical and systematic errors combined. Not independent of $a_2(\chi_{c1})$ and $b_2(\chi_{c1})$ values from ARTUSO 09.

$\chi_{c1}(1P)$ REFERENCES

ABLIKIM	24AC	PR D110 032016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BE	PR D110 032022	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BT	PR D110 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BW	PR D110 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BX	PR D110 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CA	PR D110 112013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24P	PR D109 072016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23N	JHEP 2305 069	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23T	PR D107 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AF	PRL 129 122001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BI	PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)

ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)
