

$\eta_c(2S)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

 $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637.8 ± 0.6	OUR AVERAGE	Error includes scale factor of 1.1.		
3637.8 ± 0.8 ± 0.2	1.6k	ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$
3637.90 ± 0.54 ± 1.40	3.7k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+ (K_S^0 K \pi)$
3643.4 ± 2.3 ± 4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1 ± 3.7 ± 2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6 ± 1.7 ± 0.6	106	¹ AAIJ	17AD LHCb	$pp \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4 ± 4.1 ± 0.7	365	² AAIJ	17BB LHCb	$pp \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-) X$
3637.0 ± 5.7 ± 3.4	178	^{3,4} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
3635.1 ± 5.8 ± 2.1	47	^{3,5} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
3646.9 ± 1.6 ± 3.6	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127	⁶ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
3638.5 ± 1.5 ± 0.8	624	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 ^{+3.9} _{-4.2} ^{+0.7} _{-2.0}	128	⁷ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁸ ABE	07 BELL	$e^+ e^- \rightarrow J/\psi (c \bar{c})$
3645.0 ± 5.5 ^{+4.9} _{-7.8}	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3639 ± 7	98	⁹ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c \bar{c}}$
3630.8 ± 3.4 ± 1.0	112	¹⁰ AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39	¹¹ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		¹² EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$ MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

² From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³ Ignoring possible interference with continuum.

⁴ With a width fixed to 11.3 MeV.

⁵ With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁷ Accounts for interference with non-resonant continuum.

⁸ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰Superseded by DEL-AMO-SANCHEZ 11M.
¹¹Superseded by VINOKUROVA 11.
¹²Assuming mass of $\psi(2S) = 3686$ MeV.

$\eta_c(2S)$ WIDTH						
VALUE (MeV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT	
11.6 $\pm$ 1.4 OUR AVERAGE						
10.5 $\pm$ 1.7 $\pm$ 3.5	1.6k		ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$	
10.77 $\pm$ 1.62 $\pm$ 1.08	3.7k		AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$	
19.8 $\pm$ 3.9 $\pm$ 3.1	569		ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$	
9.9 $\pm$ 4.8 $\pm$ 2.9	57		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	
16.9 $\pm$ 6.4 $\pm$ 4.8	127	1	ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$	
13.4 $\pm$ 4.6 $\pm$ 3.2	624	2	DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
6.6 $\begin{smallmatrix} + & 8.4 \\ - & 5.1 \end{smallmatrix} \begin{smallmatrix} +2.6 \\ -0.9 \end{smallmatrix}$	128	3	VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$	
6.3 $\pm$ 12.4 $\pm$ 4.0	61		ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta_c' \rightarrow K_S^0 K^\pm \pi^\mp$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 23	90	98	4 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
22 $\pm$ 14		121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$	
17.0 $\pm$ 8.3 $\pm$ 2.5		112	5 AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$	
<55	90	39	6 CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$	
<8.0	95		7 EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$	
1 From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.						
2 Ignoring possible interference with continuum.						
3 Accounts for interference with non-resonant continuum.						
4 From the fit of the kaon momentum spectrum. Systematic errors not evaluated.						
5 Superseded by DEL-AMO-SANCHEZ 11M.						
6 For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.						
7 For a mass value of 3594 ± 5 MeV						

$\eta_c(2S)$ DECAY MODES			
Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 hadrons	seen		
Γ_2 $K \bar{K} \pi$	(1.9 $^{+1.2}_{-1.0}$) %		
Γ_3 $K \bar{K} \eta$	(7 $^{+5}_{-4}$) $\times 10^{-3}$		
Γ_4 $2\pi^+ 2\pi^-$	< 2.6 $\times 10^{-3}$	90%	
Γ_5 $a_0(1450)\pi$	seen		
Γ_6 $a_2(1700)\pi$	seen		
Γ_7 $a_0(1710)\pi$	seen		

Γ_8	$\rho^0 \rho^0$	$< 2.3 \times 10^{-3}$	90%
Γ_9	$3\pi^+ 3\pi^-$	$(1.7^{+0.9}_{-1.1}) \%$	
Γ_{10}	$K^+ K^- \pi^+ \pi^-$	$< 2.5 \times 10^{-3}$	90%
Γ_{11}	$K^{*0} \bar{K}^{*0}$	$< 4 \times 10^{-3}$	90%
Γ_{12}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.5^{+1.0}_{-0.9}) \%$	
Γ_{13}	$K^+ K^- 2\pi^+ 2\pi^-$	$< 1.8 \%$	90%
Γ_{14}	$K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	$(1.4^{+0.8}_{-1.0}) \%$	
Γ_{15}	$2K^+ 2K^-$	$< 1.4 \times 10^{-3}$	90%
Γ_{16}	$K_2^*(1430) \bar{K} + \text{c.c.}$	seen	
Γ_{17}	$K_0^*(1950) \bar{K} + \text{c.c.}$	seen	
Γ_{18}	$K_0^*(2600) \bar{K} + \text{c.c.}$	seen	
Γ_{19}	$\phi \phi$	$< 1.4 \times 10^{-3}$	90%
Γ_{20}	$p \bar{p}$	$< 3.2 \times 10^{-4}$	90%
Γ_{21}	$p \bar{p} \pi^+ \pi^-$	seen	
Γ_{22}	$\gamma \gamma$	$(1.8^{+1.0}_{-1.1}) \times 10^{-4}$	
Γ_{23}	$\gamma J/\psi(1S)$	$< 1.8 \%$	90%
Γ_{24}	$\pi^+ \pi^- \eta$	$(5.5^{+3.3}_{-4.0}) \times 10^{-3}$	
Γ_{25}	$\pi^+ \pi^- \eta'$	$(2.7^{+2.0}_{-1.8}) \times 10^{-3}$	
Γ_{26}	$\pi^+ \pi^- \eta_c(1S)$	$< 4 \%$	90%

FIT INFORMATION

A multiparticle fit to $\eta_c(2S)$ and $\psi(2S)$ with 4 branching ratios uses 5 measurements to determine 3 parameters. The overall fit has a $\chi^2 = 2.6$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_3	$\begin{array}{ c } \hline 92 \\ \hline \end{array}$
x_2	

$\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$					Γ_{22}
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.44 ± 0.14	106	¹ XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
1.3 ± 0.6		² ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Assuming that the branching fraction into $\eta' \pi^+ \pi^-$ is the same as for $\eta_c(1S)$.

² They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.

$\eta_c(2S) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_2 \Gamma_{22} / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
39 ± 6 OUR AVERAGE				
41 ± 4 ± 6	624	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
33.6 ± 7.2 ± 8.1		¹ NAKAZAWA 08	BELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

¹ NAKAZAWA 08 reports $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) \times \Gamma(\gamma\gamma) = 11.2 \pm 2.4 \pm 2.7$ eV which we multiplied by 3 to account for isospin symmetry.

$\Gamma(2\pi^+ 2\pi^-) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_4 \Gamma_{22} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.5	90	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+ \pi^-)$

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{10} \Gamma_{22} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{12} \Gamma_{22} / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
30 ± 6 ± 5	1201	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(2K^+ 2K^-) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{15} \Gamma_{22} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$

$\Gamma(\pi^+ \pi^- \eta') \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{25} \Gamma_{22} / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.6^{+1.2}_{-1.1} ± 1.1	106	XU	18	BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{26} \Gamma_{22} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<133	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

$\eta_c(2S) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma^2(\text{total})$

$\Gamma(p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

$\Gamma_{20} / \Gamma \times \Gamma_{22} / \Gamma$

VALUE (units 10 ⁻⁸)	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	^{1,2,3} AMBROGIANI 01	E835	$\bar{p} p \rightarrow \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.0	90	^{1,2,4} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	^{2,4} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

¹ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

² For a total width $\Gamma=5$ MeV.

³ For the resonance mass region 3589–3599 MeV/ c^2 .

⁴ For the resonance mass region 3575–3660 MeV/ c^2 .

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABREU	98O	DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
seen	¹ EDWARDS	82C	CBAL	$e^+e^- \rightarrow \gamma X$

¹ For a mass value of 3594 ± 5 MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.9^{+1.2}_{-1.0}$ OUR FIT

$1.9 \pm 0.4 \pm 1.1$	59 ± 12	¹ AUBERT	08AB	BABR	$B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
-----------------------	-------------	---------------------	------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	127 ± 18	ABLIKIM	12G	BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	39 ± 11	² CHOI	02	BELL	$B \rightarrow K K_S K^- \pi^+$

¹ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.

² For a mass value of 3654 ± 6 MeV

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$

Γ_3/Γ_2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

39 $\pm$ 10 OUR FIT Error includes scale factor of 1.3.

$27.3 \pm 7.0 \pm 9.0$	225	¹ LEES	14E	BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$
------------------------	-----	-------------------	-----	------	---

¹ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	------	---------------------------------------

$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	¹ AAIJ	23AH	LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
------	-------------------	------	------	------------------------------------

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$

$\Gamma(a_2(1700)\pi)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$. $\Gamma(a_0(1710)\pi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$. $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

 $\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$ Γ_{12}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.73 \pm 0.17 \pm 0.17$	1201	¹ DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M. We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$. $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

 $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

 $\Gamma(K_2^*(1430) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(K_0^*(1950)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(K_0^*(2600)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ABLIKIM 11H BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 106 ¹ AAIJ 17AD LHCb $pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$

¹ AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 110 ¹ CHILIKIN 19 BELL $e^+e^- \rightarrow \gamma(4S)$

¹ CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S)K^+$ and $B^0 \rightarrow \eta_c(2S)K_S^0$ with 12.3 and 5.9 standard deviations, respectively.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4 \times 10^{-4}$ 90 ¹ WICHT 08 BELL $B^\pm \rightarrow K^\pm \gamma\gamma$

not seen AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$

<0.01 90 LEE 85 CBAL $\psi' \rightarrow \text{photons}$

¹ WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$.

$\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$ Γ_{26}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.33 90 ¹ LEES 12AE BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

¹ We divided the reported limit by 3 to take into account isospin relations.

$\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \\ \Gamma_2 / \Gamma \times \Gamma_{185}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03 ± 0.12 OUR FIT	Error includes scale factor of 1.2.			
1.00 ± 0.10 OUR AVERAGE				
$0.97 \pm 0.06 \pm 0.09$	1.6k	ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$
$1.30 \pm 0.20 \pm 0.30$	127	ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \\ \Gamma_3 / \Gamma \times \Gamma_{185}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.0 OUR FIT	Error includes scale factor of 1.3.				
$4.78 \pm 0.64 \pm 0.68$	362	¹ ABLIKIM	24BW BES3		$\psi(2S) \rightarrow \gamma K^+ K^- \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 11.8	90	² CRONIN-HEN..10	CLEO		$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
¹ ABLIKIM 24BW reports a value of $(2.39 \pm 0.32 \pm 0.34) \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 to account for isospin symmetry.					
² CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.					

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \\ \Gamma_4 / \Gamma \times \Gamma_{185}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.43 \times 10^{-6}$	90	ABLIKIM	24BT BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 14.6 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO		$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.					

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \\ \Gamma_8 / \Gamma \times \Gamma_{185}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 12.7 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \\ \Gamma_9 / \Gamma \times \Gamma_{185}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$9.2 \pm 1.0 \pm 1.2$		569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 13.2	90	¹ CRONIN-HEN..10	CLEO		$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.					

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{10}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 9.6 \times 10^{-6}$ 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{11}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 19.6 \times 10^{-7}$ 90 ABLIKIM 11H BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{12}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 43.0 \times 10^{-6}$ 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{13}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 9.7 \times 10^{-6}$ 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{14}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE (units 10^{-6})	CL% EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	----------	-------------	------	---------

7.9 $\pm$ 1.8 OUR AVERAGE

9.31 $\pm$ 0.72 $\pm$ 2.77 3140 ABLIKIM 24Q BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

7.03 $\pm$ 2.10 $\pm$ 0.7 60 ABLIKIM 13K BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.2 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow \phi\phi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))}{\Gamma_{\text{total}}} \frac{\Gamma_{19}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 7.8 \times 10^{-7}$ 90 ABLIKIM 11H BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{23}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{24}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10^{-6})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$2.97 \pm 0.81 \pm 0.26$		106	ABLIKIM	23Q BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.3	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$
--------	----	-----------------------------	------	---

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{25}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<14.2 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$
------------------------	----	-----------------------------	------	--

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{26}/\Gamma \times \Gamma_{185}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.21 \times 10^{-5}$	90	¹ ABLIKIM	24Q BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.7 \times 10^{-4}$	90	² CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$
-----------------------	----	-----------------------------	------	---

¹ $\eta_c(1S)$ reconstructed in the final states $K^+K^-\pi^0$ and $K_S^0 K^\pm \pi^\mp$.

² Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\eta_c(2S)$ REFERENCES

ABLIKIM	24BT	PR D110 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BW	PR D110 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24J	PR D109 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24Q	PR D109 072017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23Q	PR D107 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
NAKAZAWA	08	NPBPS 184 220	H. Nakazawa	(BELLE Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)