

$\omega(782)$


$$I^G(J^{PC}) = 0^-(1^--)$$

 $\omega(782)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
782.66±0.13 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
777.9 ±2.2 ^{+4.3} _{-2.2}	900k	ANDREEV 20	H1	$e p \rightarrow e \pi^+ \pi^- p$
783.20±0.13±0.16	18680	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
782.68±0.09±0.04	11200	¹ AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.79±0.08±0.09	1.2M	² ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.96±0.17±0.80	11k	³ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
782.08±0.36±0.82	3463	⁴ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
781.96±0.13±0.17	15k	AMSLER 93B	CBAR	0.0 $\bar{p} p \rightarrow \omega \pi^0 \pi^0$
782.4 ±0.2	270k	WEIDENAUER 93	ASTE	$\bar{p} p \rightarrow 2\pi^+ 2\pi^- \pi^0$
782.4 ±0.5	7000	⁵ KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
782.43±0.03	34M	⁶ IGNATOV 24	CMD3	$e^+ e^- \rightarrow \pi^+ \pi^-$
782.58±0.03±0.01		⁷ HOID 20	RVUE	$e^+ e^- \rightarrow \pi^0 \gamma$
781.68±0.09±0.03		⁸ COLANGELO 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
782.63±0.03±0.01		⁹ HOFERICH... 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.91±0.24		¹⁰ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
782.7 ±0.1 ±1.5	19500	¹¹ WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He} \omega$
781.78±0.10		¹¹ BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.2 ±0.4	1488	¹² KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
783.3 ±0.4	433	CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.5 ±0.8	33260	ROOS 80	RVUE	0.0-3.6 $\bar{p} p$
782.6 ±0.8	3000	BENKHEIRI 79	OMEG	9-12 $\pi^\pm p$
781.8 ±0.6	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p} p \rightarrow 5\pi$
782.7 ±0.9	535	VANAPEL... 78	HBC	7.2 $\bar{p} p \rightarrow \bar{p} p \omega$
783.5 ±0.8	2100	GESSAROLI 77	HBC	11 $\pi^- p \rightarrow \omega n$
782.5 ±0.8	418	AGUILAR... 72B	HBC	3.9,4.6 $K^- p$
783.4 ±1.0	248	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K^+ K^- \omega$
781.0 ±0.6	510	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K_1^+ K_1^- \omega$
783.7 ±1.0	3583	¹³ COYNE 71	HBC	3.7 $\pi^+ p \rightarrow p \pi^+ \pi^+ \pi^- \pi^0$
784.1 ±1.2	750	ABRAMOVI... 70	HBC	3.9 $\pi^- p$
783.2 ±1.6		¹⁴ BIGGS 70B	CNTR	<4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
782.4 ±0.5	2400	BIZZARRI 69	HBC	0.0 $\bar{p} p$

¹ Update of AKHMETSHIN 00C.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ From the $\eta \rightarrow \gamma \gamma$ decay.⁴ From the $\eta \rightarrow 3\pi^0$ decay.⁵ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

- 6 From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. Systematic errors not estimated.
- 7 The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 782.736 ± 0.024 MeV.
- 8 The ω mass was extracted from a dispersively improved Breit-Wigner parameterization, the ω width fixed at 8.49 ± 0.08 MeV. The value does not include vacuum polarization which would shift the mass to $781.81 \pm 0.09 \pm 0.03$ MeV. The mixing parameter is assumed real valued.
- 9 The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.
- 10 From the $\rho - \omega$ interference in the $\pi^+\pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.
- 11 Systematic uncertainties underestimated.
- 12 Systematic uncertainties not estimated.
- 13 From best-resolution sample of COYNE 71.
- 14 From ω - ρ interference in the $\pi^+\pi^-$ mass spectrum assuming ω width 12.6 MeV.



$\omega(782)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
8.68±0.13 OUR AVERAGE				
8.68±0.23±0.10	11200	1 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.68±0.04±0.15	1.2M	2 ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.57 ± 0.06	34M	³ IGNATOV	24	CMD3	$e^+e^- \rightarrow \pi^+\pi^-$
$8.65 \pm 0.06 \pm 0.01$		⁴ HOID	20	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
$8.71 \pm 0.04 \pm 0.04$		⁵ HOFERICH...	19	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.13 ± 0.45		⁶ LEES	12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
8.2 ± 0.3	19500	⁷ WURZINGER	95	SPEC	$1.33 \text{ } p d \rightarrow {}^3\text{He} \omega$
8.4 ± 0.1		⁸ AULCHENKO	87	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.30 ± 0.40		⁷ BARKOV	87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.8 ± 0.9	1488	⁹ KURDADZE	83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 ± 0.8	433	⁷ CORDIER	80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
12 ± 2	1430	COOPER	78B	HBC	$0.7\text{--}0.8 \text{ } \bar{p} p \rightarrow 5\pi$
9.4 ± 2.5	2100	GESSAROLI	77	HBC	$11 \text{ } \pi^- p \rightarrow \omega n$
10.22 ± 0.43	20000	¹⁰ KEYNE	76	CNTR	$\pi^- p \rightarrow \omega n$
13.3 ± 2	418	AGUILAR-...	72B	HBC	$3.9, 4.6 \text{ } K^- p$
9.1 ± 0.8	451	⁷ BENAKSAS	72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
10.5 ± 1.5		BORENSTEIN	72	HBC	$2.18 \text{ } K^- p$
$7.70 \pm 0.9 \pm 1.15$	940	BROWN	72	MMS	$2.5 \text{ } \pi^- p \rightarrow n \text{ } \omega$
10.3 ± 1.4	510	BIZZARRI	71	HBC	$0.0 \text{ } p \bar{p} \rightarrow K_1^0 K_1^0 \omega$
12.8 ± 3.0	248	BIZZARRI	71	HBC	$0.0 \text{ } p \bar{p} \rightarrow K^+ K^- \omega$
9.5 ± 1.0	3583	COYNE	71	HBC	$3.7 \text{ } \pi^+ p \rightarrow$ $p \pi^+ \pi^+ \pi^- \pi^0$

¹ Update of AKHMETSHIN 00C.

² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

³ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. Systematic errors not estimated.

⁴ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 8.63 ± 0.05 MeV.

⁵ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

⁶ From the $\rho - \omega$ interference in the $\pi^+\pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.

⁷ Systematic uncertainties underestimated.

⁸ Relativistic Breit-Wigner includes radiative corrections. Systematic uncertainties not estimated.

⁹ Systematic uncertainties not estimated.

¹⁰ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

$\omega(782)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \quad \pi^+\pi^-\pi^0$	$(89.2 \pm 0.7) \%$	
$\Gamma_2 \quad \pi^0\gamma$	$(8.33 \pm 0.25) \%$	S=2.1
$\Gamma_3 \quad \pi^+\pi^-$	$(1.53 \pm 0.12) \%$	S=1.2
$\Gamma_4 \quad \text{neutrals (excluding } \pi^0\gamma)$	$(7^{+7}_{-5}) \times 10^{-3}$	S=1.1

Γ_5	$\eta\gamma$	$(4.5 \pm 0.4) \times 10^{-4}$	S=1.1
Γ_6	$\pi^0 e^+ e^-$	$(7.7 \pm 0.6) \times 10^{-4}$	
Γ_7	$\pi^0 \mu^+ \mu^-$	$(1.34 \pm 0.18) \times 10^{-4}$	S=1.5
Γ_8	$\eta e^+ e^-$		
Γ_9	$e^+ e^-$	$(7.41 \pm 0.19) \times 10^{-5}$	S=1.8
Γ_{10}	$\pi^+ \pi^- \pi^0 \pi^0$	$< 2 \times 10^{-4}$	CL=90%
Γ_{11}	$\pi^+ \pi^- \gamma$	$< 3.6 \times 10^{-3}$	CL=95%
Γ_{12}	$\pi^+ \pi^- \pi^+ \pi^-$	$< 1 \times 10^{-3}$	CL=90%
Γ_{13}	$\pi^0 \pi^0 \gamma$	$(6.7 \pm 1.1) \times 10^{-5}$	
Γ_{14}	$\eta \pi^0 \gamma$	$< 3.3 \times 10^{-5}$	CL=90%
Γ_{15}	$\mu^+ \mu^-$	$(7.4 \pm 1.8) \times 10^{-5}$	
Γ_{16}	3γ	$< 1.9 \times 10^{-4}$	CL=95%

Charge conjugation (C) violating modes

Γ_{17}	$\eta \pi^0$	C	$< 2.1 \times 10^{-4}$	CL=90%
Γ_{18}	$2\pi^0$	C	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{19}	$3\pi^0$	C	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{20}	invisible		$< 7 \times 10^{-5}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to 15 branching ratios uses 49 measurements and one constraint to determine 10 parameters. The overall fit has a $\chi^2 = 48.9$ for 40 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	24								
x_3	-18	-4							
x_4	-93	-55	1						
x_5	7	16	-1	-12					
x_6	-1	0	0	0	0				
x_7	0	0	0	0	0	0			
x_9	-28	-59	5	44	-28	0	0		
x_{13}	1	4	0	-2	1	0	0	-2	
x_{15}	0	0	0	0	0	0	0	0	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_9	x_{13}

$\omega(782)$ PARTIAL WIDTHS **$\Gamma(\pi^0\gamma)$** **Γ_2**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
880 ± 50	7815	¹ ACHASOV	13	SND $1.05\text{--}2.00$ $e^+e^- \rightarrow \pi^0\pi^0\gamma$
$788 \pm 12 \pm 27$	36500	² ACHASOV	03	SND $0.60\text{--}0.97$ $e^+e^- \rightarrow \pi^0\gamma$
764 ± 51	10625	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

¹ Systematic uncertainty not estimated.² Using $\Gamma_\omega = 8.44 \pm 0.09$ MeV and $B(\omega \rightarrow \pi^0\gamma)$ from ACHASOV 03. **$\Gamma(\eta\gamma)$** **Γ_5**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
6.1 ± 2.5	¹ DOLINSKY	89	ND $e^+e^- \rightarrow \eta\gamma$

¹ Using $\Gamma_\omega = 8.4 \pm 0.1$ MeV and $B(\omega \rightarrow \eta\gamma)$ from DOLINSKY 89. **$\Gamma(e^+e^-)$** **Γ_9**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.60 $\pm$ 0.02 OUR EVALUATION				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.591 ± 0.015	11200	^{1,2} AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$0.653 \pm 0.003 \pm 0.021$	1.2M	³ ACHASOV	03D	RVUE $0.44\text{--}2.00$ $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.600 ± 0.031	10625	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.891 \pm 0.007$ and $\Gamma_{\text{total}} = 8.44 \pm 0.09$ MeV.² Update of AKHMETSHIN 00C.³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$. **$\omega(782) \Gamma(i) \Gamma(e^+e^-) / \Gamma(\text{total})$** **$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-) / \Gamma_{\text{total}}$** **$\Gamma_1\Gamma_9/\Gamma$**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
569.8 $\pm$ 3.1 $\pm$ 8.2	¹ LEES	21B	BABR 10.5 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

¹ From the cross section for $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. **$\omega(782) \Gamma(e^+e^-) \Gamma(i) / \Gamma^2(\text{total})$** **$\Gamma(e^+e^-) / \Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0) / \Gamma_{\text{total}}$** **$\Gamma_9/\Gamma \times \Gamma_1/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.61 $\pm$ 0.16 OUR FIT Error includes scale factor of 2.0.				
6.45 $\pm$ 0.15 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
$6.82 \pm 0.04 \pm 0.23$	123k	¹ IGNATOV	24	CMD3 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$6.24 \pm 0.11 \pm 0.08$	11.2k	² AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$6.74 \pm 0.04 \pm 0.24$	1.2M	^{3,4} ACHASOV	03D	RVUE $0.44\text{--}2.00$ $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
6.37 ± 0.35		³ DOLINSKY	89	ND $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

6.20±0.13		⁵	BENAYOUN	10	RVUE	0.4–1.05	e^+e^-
6.70±0.06±0.27		⁶	AUBERT,B	04N	BABR	10.6	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
6.45±0.24		⁷	BARKOV	87	CMD		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.79±0.42	1488	⁸	KURDADZE	83B	OLYA		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.89±0.54	433	⁷	CORDIER	80	DM1		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
7.54±0.84	451	⁷	BENAKSAS	72B	OSPK		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

- ¹ From a fit of the cross section using a VMD model which includes the interference of the ω and ϕ and non-resonant background, with some parameters, including ϕ mass and width, fixed at their PDG 22 values.

² Update of AKHMETSHIN 00C.

³ Recalculated by us from the cross section in the peak.


⁴ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁵ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

⁶ Superseded by LEES 21B.

⁷ Recalculated by us from the cross section in the peak. Systematic uncertainties underestimated.

⁸ Recalculated by us from the cross section in the peak. Systematic uncertainties not estimated.



$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$				$\Gamma_9/\Gamma \times \Gamma_2/\Gamma$	
<i>VALUE (units 10^{-6})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	
6.17 ±0.16	OUR FIT	Error includes scale factor of 2.0.			
6.34 ±0.10	OUR AVERAGE				
6.336±0.056±0.089		¹ ACHASOV	16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$

6.47 ±0.14 ±0.39	18k	AKHMETSHIN 05	CMD2	0.60-1.38	$e^+e^- \rightarrow \pi^0\gamma$
6.34 ±0.21 ±0.21	10k	² DOLINSKY 89	ND		$e^+e^- \rightarrow \pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.80 ±0.13		³ BENAYOUN 10	RVUE	0.4-1.05	e^+e^-
6.50 ±0.11 ±0.20	36k	⁴ ACHASOV 03	SND	0.60-0.97	$e^+e^- \rightarrow \pi^0\gamma$

¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 03.

² Recalculated by us from the cross section in the peak.

³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.


⁴ Using $\sigma(\phi \rightarrow \pi^0\gamma)$ from ACHASOV 00 and $m_\omega = 782.57$ MeV in the model with the energy-independent phase of ρ - ω interference equal to $(-10.2 \pm 7.0)^\circ$.

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_3/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.225±0.030 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
1.204±0.013±0.023	34M	¹ IGNATOV 24	CMD3	$e^+e^- \rightarrow \pi^+\pi^-$
1.318±0.051±0.021		² ACHASOV 21	SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.225±0.058±0.041	800k	³ ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.166±0.036		⁴ BENAYOUN 13	RVUE	0.4-1.05 e^+e^-
1.05 ±0.08		⁵ DAVIER 13	RVUE	$e^+e^- \rightarrow \pi^+\pi^-(\gamma)$

WEIGHTED AVERAGE

1.225±0.030 (Error scaled by 1.3)



¹ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference with ω and ϕ masses and widths constrained by the values and their errors from PDG 22, and leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit.

²From a fit of the cross section in the energy range $0.525 < \sqrt{s} < 0.883$ GeV parameterized by the sum of the Breit-Wigner amplitudes for the $\rho(770)$, ω and $\rho(1450)$ resonances. The measured phase of the $\rho(770)$ — ω interference is $(110.7 \pm 1.5 \pm 1.0)^\circ$.

³Supersedes ACHASOV 05A.

⁴A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data. Supersedes BENAYOUN 10.

⁵From $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data of LEES 12G.

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.32±0.28 OUR FIT Error includes scale factor of 1.1.

3.18±0.28 OUR AVERAGE

3.10±0.31±0.11	33k	¹ ACHASOV	07B	SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
----------------	-----	----------------------	-----	-----	--

3.17 ^{+1.85} _{-1.31} ±0.21	17.4k	² AKHMETSHIN 05	CMD2		0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
--	-------	----------------------------	------	--	---

3.41±0.52±0.21	23k	^{3,4} AKHMETSHIN 01B	CMD2		$e^+e^- \rightarrow \eta\gamma$
----------------	-----	-------------------------------	------	--	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.50±0.10		⁵ BENAYOUN 10	RVUE		0.4–1.05 e^+e^-
-----------	--	--------------------------	------	--	-------------------

¹From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

²From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

³From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

⁴The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

⁵A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_{15}/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.3±1.8±2.2	4.5M	¹ ANASTASI	17	KLOE	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
--------------------	------	-----------------------	----	------	---------------------------------------

¹From a fit of the real part of the vacuum polarization by a sum of the leptonic and hadronic contributions, where the hadronic contribution is parametrized as a sum of Breit-Wigner resonances $\omega(782)$, $\phi(1020)$ and using a GOUNARIS 68 parametrization for the $\rho(770)$, and a non-resonant term.

$\omega(782)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_1/Γ

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9024±0.0019		¹ AMBROSINO 08G	KLOE		1.0–1.03 $e^+e^- \rightarrow \pi^+\pi^-\pi^0, 2\pi^0\gamma$
---------------	--	----------------------------	------	--	---

0.8965±0.0016±0.0048	1.2M	^{2,3} ACHASOV	03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
----------------------	------	------------------------	-----	------	--

0.880 ±0.020 ±0.032	11200	^{3,4} AKHMETSHIN 00C	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
---------------------	-------	-------------------------------	------	--	--------------------------------------

0.8942±0.0062		³ DOLINSKY 89	ND		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
---------------	--	--------------------------	----	--	--------------------------------------

¹Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

²Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

³Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.

⁴Using $\Gamma(e^+e^-) = 0.60 \pm 0.02$ keV.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.88 ± 0.18		¹ ACHASOV 16A	SND	$0.60\text{--}1.38 e^+e^- \rightarrow \pi^0\gamma$
8.09 ± 0.14		² AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
$9.06 \pm 0.20 \pm 0.57$	18k	^{3,4} AKHMETSHIN 05	CMD2	$0.60\text{--}1.38 e^+e^- \rightarrow \pi^0\gamma$
$9.34 \pm 0.15 \pm 0.31$	36k	⁴ ACHASOV 03	SND	$0.60\text{--}0.97 e^+e^- \rightarrow \pi^0\gamma$
$8.65 \pm 0.16 \pm 0.42$	1.2M	^{5,6} ACHASOV 03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.39 ± 0.24	9k	⁷ BENAYOUN 96	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.88 ± 0.62	10k	⁴ DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

¹ Using $B(\omega \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 03.² Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.³ Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$.⁴ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.⁵ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.⁶ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.⁷ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions. $\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_2/Γ_1


VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
9.33 ± 0.28 OUR FIT	Error includes scale factor of 2.3.		
9.05 ± 0.27 OUR AVERAGE	Error includes scale factor of 1.8.		
8.97 ± 0.16	AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
$9.94 \pm 0.36 \pm 0.38$	¹ AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
8.4 ± 1.3	KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
10.9 ± 2.5	BENAKSAS 72C	OSPK	$e^+e^- \rightarrow \pi^0\gamma$
8.1 ± 2.0	BALDIN 71	HLBC	$2.9 \pi^+p$
13 ± 4	JACQUET 69B	HLBC	$2.05 \pi^+p \rightarrow \pi^+p\omega$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$9.7 \pm 0.2 \pm 0.5$	^{2,3} ACHASOV 03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.9 ± 0.7	² DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

¹ From $\sigma_0^{\omega\pi^0 \rightarrow \pi^0\pi^0\gamma}(m_\phi)/\sigma_0^{\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0\pi^0}(m_\phi)$ with a phase-space correction factor of 1/1.023.² Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.³ Using ACHASOV 03. Based on 1.2M events. $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_3/Γ See also $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.53 ± 0.12 OUR FIT	Error includes scale factor of 1.2.			
1.49 ± 0.13 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$1.46 \pm 0.12 \pm 0.02$	900k	¹ AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
$1.30 \pm 0.24 \pm 0.05$	11.2k	² AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
$2.38^{+1.77}_{-0.90} \pm 0.18$	5.4k	³ ACHASOV 02E	SND	$1.1\text{--}1.38 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

2.3 ± 0.5		BARKOV	85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
1.6 $^{+0.9}_{-0.7}$		QUENZER	78	DM1	$e^+e^- \rightarrow \pi^+\pi^-$
3.6 ± 1.9		BENAKSAS	72	OSPK	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.29 $\pm 0.22 \pm 0.03$	970k	4,5 ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.28 $\pm 0.22 \pm 0.03$	970k	6,7 ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.52 ± 0.08		8 HANHART	18	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1.75 ± 0.11	4.5M	9 ACHASOV	05A	SND	$e^+e^- \rightarrow \pi^+\pi^-$
2.01 ± 0.29		10 BENAYOUN	03	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1.9 ± 0.3		11 GARDNER	99	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
2.3 ± 0.4		12 BENAYOUN	98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
1.0 ± 0.11		13 WICKLUND	78	ASPK	3,4,6 $\pi^\pm N$
1.22 ± 0.30		ALVENSLEB...	71C	CNTR	Photoproduction
1.3 $^{+1.2}_{-0.9}$		MOFFEIT	71	HBC	2,8,4.7 γp
0.80 $^{+0.28}_{-0.20}$		14 BIGGS	70B	CNTR	4.2 $\gamma C \rightarrow \pi^+\pi^- C$

WEIGHTED AVERAGE
1.49 $\pm$ 0.13 (Error scaled by 1.3)



- ¹ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.
² Update of AKHMETSHIN 02.
³ From the $m_{\pi^+\pi^-}$ spectrum taking into account the interference of the $\rho\pi$ and $\omega\pi$ amplitudes.
⁴ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.
⁵ ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.25 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm$

$0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.

⁷ ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.22 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ Dispersive analysis. Value extracted from average of data from AUBERT 09AS, AKHMETSHIN 07, ACHASOV 06, AMBROSINO 11A, BABUSCI 13D, ABLIKIM 16B normalised by PDG 16 evaluation for $\Gamma(\omega \rightarrow e^+e^-)$.

⁹ Using $\Gamma(\omega \rightarrow e^+e^-)$ from the 2004 Edition of this Review (PDG 04).

¹⁰ Using the data of AKHMETSHIN 02 in the hidden local symmetry model.

¹¹ Using the data of BARKOV 85.

¹² Using the data of BARKOV 85 in the hidden local symmetry model.

¹³ From a model-dependent analysis assuming complete coherence.

¹⁴ Re-evaluated under $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ by BEHREND 71 using more accurate $\omega \rightarrow \rho$ photoproduction cross-section ratio.

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_3/Γ_1

See also $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0172±0.0014 OUR FIT	Error includes scale factor of 1.2.		
0.026 ±0.005 OUR AVERAGE			
0.021 ^{+0.028} _{-0.009}	1,2 RATCLIFF	72 ASPK	15 $\pi^- p \rightarrow n 2\pi$
0.028 ±0.006	1 BEHREND	71 ASPK	Photoproduction
0.022 ^{+0.009} _{-0.01}	3 ROOS	70 RVUE	

¹ The fitted width of these data is 160 MeV in agreement with present average, thus the ω contribution is overestimated. Assuming ρ width 145 MeV.

² Significant interference effect observed. NB of $\omega \rightarrow 3\pi$ comes from an extrapolation.

³ ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\gamma)$

Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.20±0.04	1.98M	1 ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ Using the data of ALOISIO 02D.

$\Gamma(\text{neutrals})/\Gamma_{\text{total}}$

$(\Gamma_2+\Gamma_4)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.091±0.006 OUR FIT				
0.081±0.011 OUR AVERAGE				
0.075±0.025		BIZZARRI	71 HBC	0.0 $p\bar{p}$
0.079±0.019		DEINET	69B OSPK	1.5 $\pi^- p$
0.084±0.015		BOLLINI	68C CNTR	2.1 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.073±0.018	42	BASILE	72B CNTR	1.67 $\pi^- p$

$\Gamma(\text{neutrals})/\Gamma(\pi^+\pi^-\pi^0)$ $(\Gamma_2+\Gamma_4)/\Gamma_1$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.102±0.008 OUR FIT**0.103^{+0.011}_{-0.010} OUR AVERAGE**

0.15 ±0.04	46	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
0.10 ±0.03	19	BARASH	67B	HBC	0.0 $\bar{p} p$
0.134±0.026	850	DIGIUGNO	66B	CNTR	1.4 $\pi^- p$
0.097±0.016	348	FLATTE	66	HBC	1.4 – 1.7 $K^- p \rightarrow \Lambda MM$
0.06 ^{+0.05} _{-0.02}		JAMES	66	HBC	2.1 $\pi^+ p$
0.08 ±0.03	35	KRAEMER	64	DBC	1.2 $\pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ±0.02	20	BUSCHBECK	63	HBC	1.5 $K^- p$

 $\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$ $\Gamma_2/(\Gamma_2+\Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.78±0.07		¹ DAKIN	72	OSPK	1.4 $\pi^- p \rightarrow n MM$
>0.81	90	DEINET	69B	OSPK	

¹ Error statistical only. Authors obtain good fit also assuming $\pi^0\gamma$ as the only neutral decay. $\Gamma(\text{neutrals})/\Gamma(\text{charged particles})$ $(\Gamma_2+\Gamma_4)/(\Gamma_1+\Gamma_3)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.100±0.008 OUR FIT**0.124±0.021** FELDMAN 67C OSPK 1.2 $\pi^- p$ $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.5 ±0.4 OUR FIT Error includes scale factor of 1.1.**6.3 ±1.3 OUR AVERAGE** Error includes scale factor of 1.2.

6.6 ±1.7		¹ ABELE	97E	CBAR	0.0 $\bar{p} p \rightarrow 5\gamma$
8.3 ±2.1		ALDE	93	GAM2	38 $\pi^- p \rightarrow \omega n$
3.0 ^{+2.5} _{-1.8}		² ANDREWS	77	CNTR	6.7–10 γCu

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 ±0.4 ±0.1	33k	³ ACHASOV	07B	SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
4.44 ^{+2.59} _{-1.83} ±0.28	17.4k	^{4,5} AKHMETSHIN	05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
5.10±0.72±0.34	23k	⁶ AKHMETSHIN	01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
0.7 to 5.5		⁷ CASE	00	CBAR	0.0 $p\bar{p} \rightarrow \eta\eta\gamma$
6.56 ^{+2.41} _{-2.55}	3525	^{2,8} BENAYOUN	96	RVUE	$e^+e^- \rightarrow \eta\gamma$
7.3 ±2.9		^{2,4} DOLINSKY	89	ND	$e^+e^- \rightarrow \eta\gamma$

¹ No flat $\eta\eta\gamma$ background assumed.² Solution corresponding to constructive ω - ρ interference.³ ACHASOV 07B reports $[\Gamma(\omega(782) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow e^+e^-)] = (3.10 \pm 0.31 \pm 0.11) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow e^+e^-) = (7.41 \pm 0.19) \times 10^{-5}$. Our first error is their experiment's error and our second

error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁴ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁵ Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁶ Using $B(\omega \rightarrow e^+e^-) = (7.07 \pm 0.19) \times 10^{-5}$ and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$. Solution corresponding to constructive ω - ρ interference. The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively). Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁷ Depending on the degree of coherence with the flat $\eta\eta\gamma$ background and using $B(\omega \rightarrow \pi^0\gamma) = (8.5 \pm 0.5) \times 10^{-2}$.

⁸ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$

Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0098 ± 0.0024	¹ ALDE	93	GAM2 $38\pi^- p \rightarrow \omega n$
0.0082 ± 0.0033	² DOLINSKY	89	ND $e^+e^- \rightarrow \eta\gamma$
0.010 ± 0.045	APEL	72B	OSPK $4-8 \pi^- p \rightarrow n 3\gamma$

¹ Model independent determination.

² Solution corresponding to constructive ω - ρ interference.

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$

Γ_6/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.7 ± 0.6	OUR FIT			
7.7 ± 0.6	OUR AVERAGE			
$7.61 \pm 0.53 \pm 0.64$		ACHASOV	08	SND $0.36-0.97 e^+e^- \rightarrow \pi^0 e^+e^-$
$8.19 \pm 0.71 \pm 0.62$		AKHMETSHIN	05A	CMD2 $0.72-0.84 e^+e^-$
5.9 ± 1.9	43	DOLINSKY	88	ND $e^+e^- \rightarrow \pi^0 e^+e^-$

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.34 ± 0.18	OUR FIT Error includes scale factor of 1.5.			
1.34 ± 0.19	OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.			
$1.41 \pm 0.09 \pm 0.15$		ARNALDI	16	NA60 400 GeV (p -A) collisions
$1.72 \pm 0.25 \pm 0.14$	3k	ARNALDI	09	NA60 158A In-In collisions
0.96 ± 0.23		DZHELYADIN	81B	CNTR $25-33 \pi^- p \rightarrow \omega n$

WEIGHTED AVERAGE
1.34±0.19 (Error scaled by 1.5)



$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$

Γ_8/Γ

VALUE (units 10 ⁻⁵)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<1.1	AKHMETSHIN 05A	CMD2	0.72-0.84 $e^+ e^-$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

Γ_9/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.741±0.019 OUR FIT	Error includes scale factor of 1.8.			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.700±0.016	11200	1,2 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.752±0.004±0.024	1.2M	2,3 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.714±0.036		2 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.72 ±0.03		2 BARKOV	87 CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.64 ±0.04	1488	2 KURDADZE	83B OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.675±0.069	433	2 CORDIER	80 DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.83 ±0.10	451	2 BENAKSAS	72B OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.77 ±0.06		4 AUGUSTIN	69D OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.65 ±0.13	33	5 ASTVACAT...	68 OSPK	Assume SU(3)+mixing

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = 0.891 \pm 0.007$. Update of AKHMETSHIN 00C.

² Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}^2$.

³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+ \pi^-) = (1.70 \pm 0.28)\%$.

⁴ Rescaled by us to correspond to ω width 8.4 MeV. Systematic errors underestimated.

⁵ Not resolved from ρ decay. Error statistical only.

$$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	ACHASOV	09A SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<200	90	KURDADZE	86 OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0036	95	WEIDENAUER	90 ASTE	$\rho\bar{p} \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.004	95	BITYUKOV	88B SPEC	$32\pi^-\rho \rightarrow \pi^+\pi^-\gamma X$

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{11}/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.066	90	KALBFLEISCH	75 HBC	$2.18 K^-\rho \rightarrow \Lambda\pi^+\pi^-\gamma$
<0.05	90	FLATTE	66 HBC	$1.2 - 1.7 K^-\rho \rightarrow \Lambda\pi^+\pi^-\gamma$

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 × 10⁻³	90	KURDADZE	88 OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 1.1 OUR FIT				
6.5 ± 1.2 OUR AVERAGE				
$6.4^{+2.4}_{-2.0} \pm 0.8$	190	¹ AKHMETSHIN	04B CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
$6.6^{+1.4}_{-1.3} \pm 0.6$	295	ACHASOV	02F SND	$0.36-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$11.8^{+2.1}_{-1.9} \pm 1.4$	190	² AKHMETSHIN	04B CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
$7.8 \pm 2.7 \pm 2.0$	63	^{1,3} ACHASOV	00G SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$12.7 \pm 2.3 \pm 2.5$	63	^{2,3} ACHASOV	00G SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ and $f_0(500)\gamma$ mechanisms.

² In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ mechanism only.

³ Superseded by ACHASOV 02F.

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{13}/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00045	90	DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.08	95	JACQUET	69B HLBC	$2.05 \pi^+\rho \rightarrow \pi^+\rho\omega$

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^0\gamma)$ Γ_{13}/Γ_2

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
8.0 ± 1.3 OUR FIT					
8.5 ± 2.9		40 ± 14	ALDE	94B	GAM2 $38\pi^- p \rightarrow \pi^0\pi^0\gamma n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 50	90		DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
< 1800	95		KEYNE	76	CNTR $\pi^- p \rightarrow \omega n$
< 1500	90		BENAKSAS	72C	OSPK e^+e^-
< 1400			BALDIN	71	HLBC $2.9\pi^+ p$
< 1000	90		BARMIN	64	HLBC $1.3\text{--}2.8\pi^- p$

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\text{neutrals})$ $\Gamma_{13}/(\Gamma_2+\Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.22 ± 0.07		¹ DAKIN	72	OSPK $1.4\pi^- p \rightarrow n\pi\pi$
< 0.19	90	DEINET	69B	OSPK
¹ See $\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$.				

 $\Gamma(\eta\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	AKHMETSHIN 04B	CMD2	$0.6\text{--}0.97 e^+e^- \rightarrow \eta\pi^0\gamma$

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
7.4 ± 1.8 OUR FIT				
7.4 ± 1.8 OUR AVERAGE				
$6.6 \pm 1.4 \pm 1.7$	4.5M	¹ ANASTASI	17	KLOE $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$9.0 \pm 2.9 \pm 1.1$	18	HEISTER	02C	ALEP $Z \rightarrow \mu^+\mu^- + X$

¹ Assuming lepton universality in the decay $\omega \rightarrow \ell^+\ell^-$ and correcting for different phase space between electron and muon final states.

 $\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{15}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.2	90	WILSON	69	OSPK $12\pi^- C \rightarrow \text{Fe}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.7	74	FLATTE	66	HBC $1.2\text{--}1.7 K^- p \rightarrow \Lambda\mu^+\mu^-$
< 1.2		BARBARO-...	65	HBC $2.7 K^- p$

 $\Gamma(\pi^0\mu^+\mu^-)/\Gamma(\mu^+\mu^-)$ Γ_7/Γ_{15}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.2 ± 0.6	30	¹ DZHELYADIN	79	CNTR $25\text{--}33\pi^- p$
¹ Superseded by DZHELYADIN 81B result above.				

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	95	¹ ABELE 97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<2	90	¹ PROKOSHKIN 95	GAM2	$38 \pi^- p \rightarrow 3\gamma n$

¹ From direct 3γ decay search.

$\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

Violates C conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.001	90	ALDE 94B	GAM2	$38\pi^- p \rightarrow \eta\pi^0 n$

$[\Gamma(\eta\gamma) + \Gamma(\eta\pi^0)]/\Gamma(\pi^+\pi^-\pi^0)$ $(\Gamma_5 + \Gamma_{17})/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.016	90	¹ FLATTE 66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda\pi^+\pi^- \text{MM}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.045	95	JACQUET 69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p\omega$

¹ Restated by us using $B(\eta \rightarrow \text{charged modes}) = 29.2\%$.

$\Gamma(\eta\pi^0)/\Gamma(\pi^0\gamma)$ Γ_{17}/Γ_2

Violates C conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ STAROSTIN 09	CRYM	$\gamma p \rightarrow \eta\pi^0 p$
¹ STAROSTIN 09 reports $[\Gamma(\omega(782) \rightarrow \eta\pi^0)/\Gamma(\omega(782) \rightarrow \pi^0\gamma)] \times [B(\eta \rightarrow 2\gamma)] < 1.01 \times 10^{-3}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.				

$\Gamma(2\pi^0)/\Gamma(\pi^0\gamma)$ Γ_{18}/Γ_2

Violates C conservation and Bose-Einstein statistics.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.59	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 2\pi^0 p$

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

Violates C conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<3 \times 10^{-4}$	90	PROKOSHKIN 95	GAM2	$38 \pi^- p \rightarrow 3\pi^0 n$

$\Gamma(3\pi^0)/\Gamma(\pi^0\gamma)$ Γ_{19}/Γ_2

Violates C conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.72	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 3\pi^0 p$

$\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{19}/Γ_1

Violates C conservation.

VALUE	CL%	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.009	90	BARBERIS 01	$450 p p \rightarrow p_f 3\pi^0 p_s$

$\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{20}/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.1 \times 10^{-5}$	90	ABLIKIM	18S BES3	$J/\psi \rightarrow \omega\eta \rightarrow \omega\pi^+\pi^-\pi^0$

PARAMETER Λ IN $\omega \rightarrow \pi^0 \ell^+ \ell^-$ DECAY

In the pole approximation the electromagnetic transition form factor for a resonance of mass M is given by the expression:

$$|F|^2 = (1 - M^2/\Lambda^2)^{-2},$$

where for the parameter Λ vector dominance predicts $\Lambda = M_\rho \approx 0.770$ GeV. The ARNALDI 09 measurement is in obvious conflict with this expectation. Note that for $\eta \rightarrow \gamma\mu^+\mu^-$ decay ARNALDI 09 and DZHELYADIN 80 obtain the value of Λ consistent with vector dominance.

PARAMETER Λ IN $\omega \rightarrow \pi^0 \mu^+ \mu^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.670 ± 0.006 OUR AVERAGE				
0.6707 ± 0.0039 ± 0.0056		¹ ARNALDI	16 NA60	400 GeV (p -A) collisions
0.668 ± 0.009 ± 0.003	3k	² ARNALDI	09 NA60	158A In-In collisions
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.65 ± 0.03		DZHELYADIN	81B CNTR	25–33 $\pi^- p \rightarrow \omega n$
¹ ARNALDI 16 reports $\Lambda^{-2}(\omega) = 2.223 \pm 0.026 \pm 0.037 \text{ GeV}^{-2}$ which we converted to the quoted Λ value.				
² ARNALDI 09 reports $\Lambda^{-2}(\omega) = 2.24 \pm 0.06 \pm 0.02 \text{ GeV}^{-2}$ which we converted to the quoted Λ value.				

PARAMETER Λ IN $\omega \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.709 ± 0.037	1.1k	¹ ADLARSON	17B A2MM	$\gamma p \rightarrow \omega p$
¹ ADLARSON 17B reports $\Lambda^{-2}(\omega\pi^0) = 1.99 \pm 0.21 \text{ GeV}^{-2}$ that we converted to the quoted Λ value.				

ENERGY DEPENDENCE OF $\omega \rightarrow \pi^+\pi^-\pi^0$ DALITZ PLOT

The following experiments fit to one or more of the coefficients α, β, γ for $|\text{matrix element}|^2 \propto P(1 + 2\alpha Z + 2\beta Z^{3/2} \sin(3\phi) + 2\gamma Z^2 + O(Z^{5/2}))$ where P is the P -wave phase-space factor and Z, ϕ are kinematical variables as defined in ADLARSON 17.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.133 ± 0.008 OUR AVERAGE				
0.1321 ± 0.0067 ± 0.0046	260k	¹ ABLIKIM	18AD BES3	$J/\psi \rightarrow \omega\eta$
0.147 ± 0.036	44k	ADLARSON	17 WASA	α in $pd \rightarrow {}^3\text{He } \omega$, $pp \rightarrow pp\omega$
¹ Keeping a term linear in Z only. A fit with the terms proportional to Z and $Z^{3/2}$ gives $\alpha = 0.133 \pm 0.041$ and $\beta = 0.037 \pm 0.054$.				

$\omega(782)$ REFERENCES

IGNATOV	24	PR D109 112002	F.V. Ignatov <i>et al.</i>	(CMD-3 Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ACHASOV	21	JHEP 2101 113	M.N. Achasov <i>et al.</i>	(SND Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ANDREEV	20	EPJ C80 1189	V. Andreev <i>et al.</i>	(H1 Collab.)
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)
COLANGELO	19	JHEP 1902 006	G. Colangelo, M. Hoferichter, P. Stoffer	(BERN+)
HOFRICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)
ABLIKIM	18AD	PR D98 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HANHART	18	EPJ C78 450	C. Hanhart <i>et al.</i>	
ADLARSON	17	PL B770 418	P. Adlarson <i>et al.</i>	(WASA-at-COSY Collab.)
ADLARSON	17B	PR C95 035208	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)
ANASTASI	17	PL B767 485	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
ARNALDI	16	PL B757 437	R. Arnaldi <i>et al.</i>	(NA60 Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)
BABUSCI	13D	PL B720 336	D. Babusci <i>et al.</i>	(CATA, CALB, BARI)
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)
DAVIER	13	EPJ C73 2597	M. Davier <i>et al.</i>	
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 136 442.		
ARNALDI	09	PL B677 260	R. Arnaldi <i>et al.</i>	(NA60 Collab.)
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)
STAROSTIN	09	PR C79 065201	A. Starostin <i>et al.</i>	(Crystal Ball Collab. at MAMI)
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 134 80.		
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 130 437.		
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 84 491.		
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 128 1201.		
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 82 841.		
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)
BENAYOUN	03	EPJ C29 397	M. Benayoun <i>et al.</i>	
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 72 411.		
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 71 519.		

AKHMETSHIN	00C	PL B476 33	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 117 1067.		
CASE	00	PR D61 032002	T. Case <i>et al.</i>	(Crystal Barrel Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
GARDNER	99	PR D59 076002	S. Gardner, H.B. O'Connell	
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)
ABELE	97E	PL B411 361	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)
PROKOSHKIN	95	PD 40 273	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)
		Translated from DANS 342 610.		
WURZINGER	95	PR C51 443	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)
ALDE	94B	PL B340 122	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
AMSLER	94C	PL B327 425	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)
ALDE	93	PAN 56 1229	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)
		Translated from YAF 56 137.		
Also		ZPHY C61 35	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)
AMSLER	93B	PL B311 362	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
WEIDENAUER	90	ZPHY C47 353	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)
BITYUKOV	88B	SJNP 47 800	S.I. Bityukov <i>et al.</i>	(SERP)
		Translated from YAF 47 1258.		
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)
		Translated from YAF 48 442.		
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)
		Translated from ZETFP 47 432.		
AULCHENKO	87	PL B186 432	V.M. Aulchenko <i>et al.</i>	(NOVO)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)
		Translated from ZETFP 43 497.		
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)
KURDADZE	83B	JETPL 36 274	A.M. Kurdadze <i>et al.</i>	(NOVO)
		Translated from ZETFP 36 221.		
DZHELYADIN	81B	PL 102B 296	R.I. Dzheilyadin <i>et al.</i>	(SERP)
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)
DZHELYADIN	80	PL 94B 548	R.I. Dzheilyadin <i>et al.</i>	(SERP)
ROOS	80	LNC 27 321	M. Roos, A. Pellinen	(HELS)
BENKHEIRI	79	NP B150 268	P. Benkheiri <i>et al.</i>	(EPOL, CERN, CDEF+)
DZHELYADIN	79	PL 84B 143	R.I. Dzheilyadin <i>et al.</i>	(SERP)
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)
QUENZER	78	PL 76B 512	A. Quenzer <i>et al.</i>	(LALO)
VANAPEL...	78	NP B133 245	G.W. van Apeldoorn <i>et al.</i>	(ZEEM)
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)
KEYNE	76	PR D14 28	J. Keyne <i>et al.</i>	(LOIC, SHMP)
Also		PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
APEL	72B	PL 41B 234	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)
BASILE	72B	Phil. Conf. 153	M. Basile <i>et al.</i>	(CERN)
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)
BENAKSAS	72B	PL 42B 507	D. Benaksas <i>et al.</i>	(ORSAY)
BENAKSAS	72C	PL 42B 511	D. Benaksas <i>et al.</i>	(ORSAY)
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)
BROWN	72	PL 42B 117	R.M. Brown <i>et al.</i>	(ILL, ILLC)
DAKIN	72	PR D6 2321	J.T. Dakin <i>et al.</i>	(PRIN)
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)
ALVENSLEB...	71C	PRL 27 888	H. Alvensleben <i>et al.</i>	(DESY)
BALDIN	71	SJNP 13 758	A.B. Baldin <i>et al.</i>	(ITEP)
		Translated from YAF 13 1318.		
BEHREND	71	PRL 27 61	H.J. Behrend <i>et al.</i>	(ROCH, CORN, FNAL)
BIZZARRI	71	NP B27 140	R. Bizzarri <i>et al.</i>	(CERN, CDEF)
COYNE	71	NP B32 333	D.G. Coyne <i>et al.</i>	(LRL)
MOFFEIT	71	NP B29 349	K.C. Moffeit <i>et al.</i>	(LRL, UCB, SLAC+)
ABRAMOVI...	70	NP B20 209	M. Abramovich <i>et al.</i>	(CERN)
BIGGS	70B	PRL 24 1201	P.J. Biggs <i>et al.</i>	(DARE)
BIZZARRI	70	PRL 25 1385	R. Bizzarri <i>et al.</i>	(ROMA, SYRA)

ROOS	70	DNPL/R7 173	M. Roos	(CERN)
		Proc. Daresbury Study Weekend No. 1.		
AUGUSTIN	69D	PL 28B 513	J.E. Augustin <i>et al.</i>	(ORSAY)
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)
DEINET	69B	PL 30B 426	W. Deinet <i>et al.</i>	(KARL, CERN)
JACQUET	69B	NC 63A 743	F. Jacquet <i>et al.</i>	(EPOL, BERG)
WILSON	69	Private Comm.	R. Wilson	(HARV)
		Also		
		PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)
ASTVACAT...	68	PL 27B 45	R.G. Astvatsaturov <i>et al.</i>	(JINR, MOSU)
BOLLINI	68C	NC 56A 531	D. Bollini <i>et al.</i>	(CERN, BGNA, STRB)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)
FELDMAN	67C	PR 159 1219	M. Feldman <i>et al.</i>	(PENN)
DIGIUGNO	66B	NC 44A 1272	G. Di Giugno <i>et al.</i>	(NAPL, FRAS, TRST)
FLATTE	66	PR 145 1050	S.M. Flatte <i>et al.</i>	(LRL)
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)
BARBARO-...	65	PRL 14 279	A. Barbaro-Galtieri, R.D. Tripp	(LRL)
BARMIN	64	JETP 18 1289	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from ZETF 45		
KRAEMER	64	PR 136 B496	R.W. Kraemer <i>et al.</i>	(JHU, NWES, WOOD)
BUSCHBECK	63	Siena Conf. 1 166	B. Buschbeck <i>et al.</i>	(VIEN, CERN, ANIK)
