

Other Particle Searches

OMITTED FROM SUMMARY TABLE  
OTHER PARTICLE SEARCHES

Revised February 2018 by K. Hikasa (Tohoku University).

We collect here those searches which do not appear in any other search categories. These are listed in the following order:

- Concentration of stable particles in matter
- General new physics searches
- Limits on jet-jet resonance in hadron collisions
- Limits on neutral particle production at accelerators
- Limits on charged particles in  $e^+e^-$  collisions
- Limits on charged particles in hadron reactions
- Limits on charged particles in cosmic rays
- Searches for quantum black hole production

Note that searches appear in separate sections elsewhere for Higgs bosons (and technipions), other heavy bosons (including  $W_R$ ,  $W'$ ,  $Z'$ , leptoquarks, axigluons), axions (including pseudo-Goldstone bosons, Majorons, familons), WIMPs, heavy leptons, heavy neutrinos, free quarks, monopoles, supersymmetric particles, and compositeness.

We no longer list for limits on tachyons and centauros. See our 1994 edition for these limits.

CONCENTRATION OF STABLE PARTICLES IN MATTER

Concentration of Heavy (Charge +1) Stable Particles in Matter

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4 \times 10^{-17}$	95	<sup>1</sup> YAMAGATA	93 SPEC	Deep sea water, $M=5\text{--}1600m_p$
$<6 \times 10^{-15}$	95	<sup>2</sup> VERKERK	92 SPEC	Water, $M=10^5$ to $3 \times 10^7$ GeV
$<7 \times 10^{-15}$	95	<sup>2</sup> VERKERK	92 SPEC	Water, $M=10^4$ , $6 \times 10^7$ GeV
$<9 \times 10^{-15}$	95	<sup>2</sup> VERKERK	92 SPEC	Water, $M=10^8$ GeV
$<3 \times 10^{-23}$	90	<sup>3</sup> HEMMICK	90 SPEC	Water, $M=1000m_p$

$<2 \times 10^{-21}$	90	<sup>3</sup> HEMMICK	90	SPEC	Water, $M = 5000m_p$
$<3 \times 10^{-20}$	90	<sup>3</sup> HEMMICK	90	SPEC	Water, $M = 10000m_p$
$<1. \times 10^{-29}$		SMITH	82B	SPEC	Water, $M=30\text{--}400m_p$
$<2. \times 10^{-28}$		SMITH	82B	SPEC	Water, $M=12\text{--}1000m_p$
$<1. \times 10^{-14}$		SMITH	82B	SPEC	Water, $M > 1000 m_p$
$<(0.2\text{--}1.) \times 10^{-21}$		SMITH	79	SPEC	Water, $M=6\text{--}350 m_p$

<sup>1</sup>YAMAGATA 93 used deep sea water at 4000 m since the concentration is enhanced in deep sea due to gravity.

<sup>2</sup>VERKERK 92 looked for heavy isotopes in sea water and put a bound on concentration of stable charged massive particle in sea water. The above bound can be translated into into a bound on charged dark matter particle ( $5 \times 10^6$  GeV), assuming the local density,  $\rho=0.3$  GeV/cm<sup>3</sup>, and the mean velocity  $\langle v \rangle=300$  km/s.

<sup>3</sup>See HEMMICK 90 Fig. 7 for other masses 100–10000  $m_p$ .

### Concentration of Heavy Stable Particles Bound to Nuclei

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 2 \times 10^{-17}$ /nucleon	95	<sup>1</sup> AFEK	21	millicharged particle search
$<1.2 \times 10^{-11}$	95	<sup>2</sup> JAVORSEK	01	SPEC Au, $M= 3$ GeV
$<6.9 \times 10^{-10}$	95	<sup>2</sup> JAVORSEK	01	SPEC Au, $M= 144$ GeV
$<1 \times 10^{-11}$	95	<sup>3</sup> JAVORSEK	01B	SPEC Au, $M= 188$ GeV
$<1 \times 10^{-8}$	95	<sup>3</sup> JAVORSEK	01B	SPEC Au, $M= 1669$ GeV
$<6 \times 10^{-9}$	95	<sup>3</sup> JAVORSEK	01B	SPEC Fe, $M= 188$ GeV
$<1 \times 10^{-8}$	95	<sup>3</sup> JAVORSEK	01B	SPEC Fe, $M= 647$ GeV
$<4 \times 10^{-20}$	90	<sup>4</sup> HEMMICK	90	SPEC C, $M = 100m_p$
$<8 \times 10^{-20}$	90	<sup>4</sup> HEMMICK	90	SPEC C, $M = 1000m_p$
$<2 \times 10^{-16}$	90	<sup>4</sup> HEMMICK	90	SPEC C, $M = 10000m_p$
$<6 \times 10^{-13}$	90	<sup>4</sup> HEMMICK	90	SPEC Li, $M = 1000m_p$
$<1 \times 10^{-11}$	90	<sup>4</sup> HEMMICK	90	SPEC Be, $M = 1000m_p$
$<6 \times 10^{-14}$	90	<sup>4</sup> HEMMICK	90	SPEC B, $M = 1000m_p$
$<4 \times 10^{-17}$	90	<sup>4</sup> HEMMICK	90	SPEC O, $M = 1000m_p$
$<4 \times 10^{-15}$	90	<sup>4</sup> HEMMICK	90	SPEC F, $M = 1000m_p$
$< 1.5 \times 10^{-13}$ /nucleon	68	<sup>5</sup> NORMAN	89	SPEC $^{206}\text{Pb}X^-$
$< 1.2 \times 10^{-12}$ /nucleon	68	<sup>5</sup> NORMAN	87	SPEC $^{56,58}\text{Fe}X^-$

<sup>1</sup>AFEK 21 search for millicharged particles bound to matter using an optomechanical device. No signal was observed. Limits placed in the abundance vs. charge plane (Fig. 3). This is translated to the mass versus charge plane by requiring bound states to be stable.

<sup>2</sup>JAVORSEK 01 search for (neutral) SIMPs (strongly interacting massive particles) bound to Au nuclei. Here  $M$  is the effective SIMP mass.

<sup>3</sup>JAVORSEK 01B search for (neutral) SIMPs (strongly interacting massive particles) bound to Au and Fe nuclei from various origins with exposures on the earth's surface, in a satellite, heavy ion collisions, etc. Here  $M$  is the mass of the anomalous nucleus. See also JAVORSEK 02.

<sup>4</sup>See HEMMICK 90 Fig. 7 for other masses 100–10000  $m_p$ .

<sup>5</sup>Bound valid up to  $m_{X^-} \sim 100$  TeV.

## GENERAL NEW PHYSICS SEARCHES

This subsection lists some of the search experiments which look for general signatures characteristic of new physics, independent of the framework of a specific model.

The observed events are compatible with Standard Model expectation, unless noted otherwise.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1	HAYRAPETY...24AO	CMS	soft unclustered energy search
2	ALKHATIB 21A	SCDM	fractionally charged relics
3	AGUILAR-AR...20B	CONN	$\nu$ elastic scatter on nuclei
4	FEDDERKE 20		CHAMPs from white dwarfs
5	SIRUNYAN 20A	CMS	SUSY/LQ search with mT2 or long-lived charged particles
6	ALCANTARA 19		Auger, superheavy DM
7	PORAYKO 18	PPTA	pulsar timing fuzzy DM search
8	AAD 15AT	ATLS	$t + \cancel{E}_T$
9	KHACHATRY...15F	CMS	$t + \cancel{E}_T$
10	AALTONEN 14J	CDF	$W + 2$ jets
11	AAD 13A	ATLS	$W W \rightarrow \ell \nu \ell' \nu$
12	AAD 13C	ATLS	$\gamma + \cancel{E}_T$
13	AALTONEN 13I	CDF	Delayed $\gamma + \cancel{E}_T$
14	CHATRCHYAN 13	CMS	$\ell^+ \ell^- + \text{jets} + \cancel{E}_T$
15	AAD 12C	ATLS	$t \bar{t} + \cancel{E}_T$
16	AALTONEN 12M	CDF	jet + $\cancel{E}_T$
17	CHATRCHYAN 12AP	CMS	jet + $\cancel{E}_T$
18	CHATRCHYAN 12Q	CMS	$Z + \text{jets} + \cancel{E}_T$
19	CHATRCHYAN 12T	CMS	$\gamma + \cancel{E}_T$
20	AAD 11S	ATLS	jet + $\cancel{E}_T$
21	AALTONEN 11AF	CDF	$\ell^\pm \ell^\pm$
22	CHATRCHYAN 11C	CMS	$\ell^+ \ell^- + \text{jets} + \cancel{E}_T$
23	CHATRCHYAN 11U	CMS	jet + $\cancel{E}_T$
24	AALTONEN 10AF	CDF	$\gamma \gamma + \ell, \cancel{E}_T$
25	AALTONEN 09AF	CDF	$\ell \gamma b \cancel{E}_T$
26	AALTONEN 09G	CDF	$\ell \ell \ell \cancel{E}_T$

<sup>1</sup> HAYRAPETYAN 24AO report on search for soft unclustered energy deposits. No signal observed. Limits placed in mediator mass vs. decay temperature plane.

<sup>2</sup> ALKHATIB 21A search for lightly ionizing fractionally charged relics scattering from Ge. No signal observed. Limits plotted in fractional charge  $f$  vs. vertical intensity plane for  $m \sim 5$  MeV to 100 TeV.

<sup>3</sup> AGUILAR-AREVALO 20B search for light BSM mediator effect on  $\nu$  elastic scatter on nuclei; no signal; limits placed in  $m(\text{mediator})$  vs. coupling plane for two models of MeV-scale mediators.

<sup>4</sup> FEDDERKE 20 place limits on cosmic relic charged massive particles (CHAMPs) due to their capture and subsequent disruption of old white dwarf stars; limits placed in the  $m(\text{CHAMP})$  vs. relic density parameter plane.

<sup>5</sup> SIRUNYAN 20A search for SUSY and LQ production using mT2 or presence of long-lived charged particle; no signal, limits placed in various mass planes for different BSM scenarios and various assumed lifetimes.

- <sup>6</sup> ALCANTARA 19 place limits on  $m(\text{WIMPzilla}=X)$  vs lifetime from upper bound on ultra high energy cosmic rays at Auger experiment: e.g.  $\tau(X) < 4 \times 10^{22}$  yr for  $m(X) = 10^{16}$  GeV.
  - <sup>7</sup> PORAYKO 18 search for deviations in the residuals of pulsar timing data using PPTA. No signal observed. Limits set on fuzzy DM with  $3 \times 10^{-24} < m(\text{DM}) < 2 \times 10^{-22}$  eV.
  - <sup>8</sup> AAD 15AT search for events with a top quark and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV with  $L = 20.3 \text{ fb}^{-1}$ .
  - <sup>9</sup> KHACHATRYAN 15F search for events with a top quark and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV with  $L = 19.7 \text{ fb}^{-1}$ .
  - <sup>10</sup> AALTONEN 14J examine events with a  $W$  and two jets in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 8.9 \text{ fb}^{-1}$ . Invariant mass distributions of the two jets are consistent with the Standard Model expectation.
  - <sup>11</sup> AAD 13A search for resonant  $WW$  production in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.7 \text{ fb}^{-1}$ .
  - <sup>12</sup> AAD 13C search for events with a photon and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.6 \text{ fb}^{-1}$ .
  - <sup>13</sup> AALTONEN 13I search for events with a photon and missing  $E_T$ , where the photon is detected after the expected timing, in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 6.3 \text{ fb}^{-1}$ . The data are consistent with the Standard Model expectation.
  - <sup>14</sup> CHATRCHYAN 13 search for events with an opposite-sign lepton pair, jets, and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.98 \text{ fb}^{-1}$ .
  - <sup>15</sup> AAD 12C search for events with a  $t\bar{t}$  pair and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 1.04 \text{ fb}^{-1}$ .
  - <sup>16</sup> AALTONEN 12M search for events with a jet and missing  $E_T$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 6.7 \text{ fb}^{-1}$ .
  - <sup>17</sup> CHATRCHYAN 12AP search for events with a jet and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 5.0 \text{ fb}^{-1}$ .
  - <sup>18</sup> CHATRCHYAN 12Q search for events with a  $Z$ , jets, and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.98 \text{ fb}^{-1}$ .
  - <sup>19</sup> CHATRCHYAN 12T search for events with a photon and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 5.0 \text{ fb}^{-1}$ .
  - <sup>20</sup> AAD 11S search for events with one jet and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 33 \text{ pb}^{-1}$ .
  - <sup>21</sup> AALTONEN 11AF search for high- $p_T$  like-sign dileptons in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 6.1 \text{ fb}^{-1}$ .
  - <sup>22</sup> CHATRCHYAN 11C search for events with an opposite-sign lepton pair, jets, and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 34 \text{ pb}^{-1}$ .
  - <sup>23</sup> CHATRCHYAN 11U search for events with one jet and missing  $E_T$  in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 36 \text{ pb}^{-1}$ .
  - <sup>24</sup> AALTONEN 10AF search for  $\gamma\gamma$  events with  $e, \mu, \tau$ , or missing  $E_T$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 1.1\text{--}2.0 \text{ fb}^{-1}$ .
  - <sup>25</sup> AALTONEN 09AF search for  $\ell\gamma b$  events with missing  $E_T$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 1.9 \text{ fb}^{-1}$ . The observed events are compatible with Standard Model expectation including  $t\bar{t}\gamma$  production.
  - <sup>26</sup> AALTONEN 09G search for  $\mu\mu\mu$  and  $\mu\mu e$  events with missing  $E_T$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 976 \text{ pb}^{-1}$ .
-

## LIMITS ON JET-JET RESONANCES

### Heavy Particle Production Cross Section

Limits are for a particle decaying to two hadronic jets.

Units(pb)	CL%	Mass(GeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
			<sup>1</sup> HAYRAPETY...24G	CMS	trijet resonance search
			<sup>2</sup> TUMASYAN 23L	CMS	dijet resonance in 4-jet events
			<sup>3</sup> AAD 20AD	ATLS	$pp$ at 13 TeV, dijet resonance
			<sup>4</sup> AAD 20T	ATLS	dijet resonance search
			<sup>5</sup> AAD 20W	ATLS	dijet resonance plus lepton
			<sup>6</sup> SIRUNYAN 20AI	CMS	dijet resonance search
			<sup>7</sup> AABOUD 19AJ	ATLS	$pp \rightarrow \gamma X, X \rightarrow jj$
			<sup>8</sup> SIRUNYAN 19B	CMS	$pp \rightarrow jA, A \rightarrow b\bar{b}$
			<sup>9</sup> SIRUNYAN 19CD	CMS	$pp \rightarrow Z'\gamma, Z' \rightarrow jj$
			<sup>10</sup> AABOUD 18AD	ATLS	$pp \rightarrow Y \rightarrow HX \rightarrow (bb) + (qq)$
			<sup>11</sup> AABOUD 18CK	ATLS	$pp \rightarrow bbb + \cancel{E}_T$
			<sup>12</sup> AABOUD 18CL	ATLS	$pp \rightarrow$ vector-like quarks
			<sup>13</sup> AABOUD 18N	ATLS	$pp \rightarrow jj$ resonance
			<sup>14</sup> SIRUNYAN 18DJ	CMS	$pp \rightarrow ZZ$ or $WZ \rightarrow \ell\bar{\ell}jj$
			<sup>15</sup> SIRUNYAN 18DY	CMS	$pp \rightarrow RR; R \rightarrow jj$
			<sup>16</sup> KHACHATRY...17W	CMS	$pp \rightarrow jj$ resonance
			<sup>17</sup> KHACHATRY...17Y	CMS	$pp \rightarrow (8-10) j + \cancel{E}_T$
			<sup>18</sup> SIRUNYAN 17F	CMS	$pp \rightarrow jj$ angular distribution
			<sup>19</sup> AABOUD 16	ATLS	$pp \rightarrow b + \text{jet}$
			<sup>20</sup> AAD 16N	ATLS	$pp \rightarrow 3$ high $E_T$ jets
			<sup>21</sup> AAD 16S	ATLS	$pp \rightarrow jj$ resonance
			<sup>22</sup> KHACHATRY...16K	CMS	$pp \rightarrow jj$ resonance
			<sup>23</sup> KHACHATRY...16L	CMS	$pp \rightarrow jj$ resonance
			<sup>24</sup> AAD 13D	ATLS	7 TeV $pp \rightarrow 2$ jets
			<sup>25</sup> AALTONEN 13R	CDF	1.96 TeV $p\bar{p} \rightarrow 4$ jets
			<sup>26</sup> CHATRCHYAN 13A	CMS	7 TeV $pp \rightarrow 2$ jets
			<sup>27</sup> CHATRCHYAN 13A	CMS	7 TeV $pp \rightarrow b\bar{b}X$
			<sup>28</sup> AAD 12S	ATLS	7 TeV $pp \rightarrow 2$ jets
			<sup>29</sup> CHATRCHYAN 12BL	CMS	7 TeV $pp \rightarrow t\bar{t}X$
			<sup>30</sup> AAD 11AG	ATLS	7 TeV $pp \rightarrow 2$ jets
			<sup>31</sup> AALTONEN 11M	CDF	1.96 TeV $p\bar{p} \rightarrow W + 2$ jets
			<sup>32</sup> ABAZOV 11I	D0	1.96 TeV $p\bar{p} \rightarrow W + 2$ jets
			<sup>33</sup> AAD 10	ATLS	7 TeV $pp \rightarrow 2$ jets
			<sup>34</sup> KHACHATRY...10	CMS	7 TeV $pp \rightarrow 2$ jets
			<sup>35</sup> ABE 99F	CDF	1.8 TeV $p\bar{p} \rightarrow b\bar{b} + \text{anything}$
			<sup>36</sup> ABE 97G	CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
<2603	95	200	<sup>37</sup> ABE 93G	CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 44	95	400	<sup>37</sup> ABE 93G	CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 7	95	600	<sup>37</sup> ABE 93G	CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets

<sup>1</sup> HAYRAPETYAN 24G search for trijet resonance at CMS with  $138 \text{ fb}^{-1}$  of data at 13 TeV. No signal observed. Limits placed on various models vs mass(resonance).

<sup>2</sup> TUMASYAN 23L search for dijet resonance in 4-jet events with  $138 \text{ fb}^{-1}$  fb of data. There are two events in the tails of the distributions, each with a four-jet mass of 8 TeV and an average dijet mass of 2 TeV, resulting in local and global significances of 3.9 and

- 1.6 standard deviations, respectively, if interpreted as a signal. Limits set for simplified diquark model.
- <sup>3</sup> AAD 20AD search for weakly supervised dijet resonance in ATLAS with  $139 \text{ fb}^{-1}$  at 13 TeV; no signal; various limits placed depending on kinematics and production cross section.
  - <sup>4</sup> AAD 20T search for dijet resonance with or without  $b$ -jets at 13 TeV and  $139 \text{ fb}^{-1}$ ; no signal; limits placed in  $\sigma \cdot \text{BF}$  vs mass plane for various BSM models.
  - <sup>5</sup> AAD 20W search for dijet resonance plus lepton with ATLAS at 13 TeV and  $139 \text{ fb}^{-1}$ ; no signal; limits placed in  $\sigma \cdot \text{BF}$  vs. mass plane for various BSM models.
  - <sup>6</sup> SIRUNYAN 20AI search for dijet resonance in CMS at 13 TeV with  $137 \text{ fb}^{-1}$ ; no signal; limits set in  $\sigma$  vs. mass plane for various BSM models.
  - <sup>7</sup> AABOUD 19AJ search for low mass dijet resonance in  $pp \rightarrow \gamma X, X \rightarrow jj$  at 13 TeV with  $79.8 \text{ fb}^{-1}$  of data; no signal found; limits placed on  $Z'$  model in coupling vs.  $m(Z')$  plane.
  - <sup>8</sup> SIRUNYAN 19B search for low mass resonance  $pp \rightarrow jA, A \rightarrow b\bar{b}$  at 13 TeV using  $35.9 \text{ fb}^{-1}$ ; no signal; exclude resonances 50–350 GeV depending on production and decay.
  - <sup>9</sup> SIRUNYAN 19CD search for  $pp \rightarrow Z'\gamma, Z' \rightarrow jj$  with fat jet ( $jj$ ); no signal, limits placed in  $m(Z')$  vs. coupling plane for  $Z'$  masses from 10 to 125 GeV.
  - <sup>10</sup> AABOUD 18AD search for new heavy particle  $Y \rightarrow HX \rightarrow (bb) + (qq)$ . No signal observed. Limits set on  $m(Y)$  vs.  $m(X)$  in the ranges of  $m(Y)$  in 1–4 TeV and  $m(X)$  in 50–1000 GeV.
  - <sup>11</sup> AABOUD 18CK search for SUSY Higgsinos in gauge-mediation via  $pp \rightarrow bbb + \cancel{E}_T$  at 13 TeV using two complementary analyses with  $24.3/36.1 \text{ fb}^{-1}$ ; no signal is found and Higgsinos with masses between 130 and 230 GeV and between 290 and 880 GeV are excluded at the 95% confidence level.
  - <sup>12</sup> AABOUD 18CL search for  $pp \rightarrow \text{vector-like quarks} \rightarrow \text{jets}$  at 13 TeV with  $36 \text{ fb}^{-1}$ ; no signal seen; limits set on various VLQ scenarios. For pure  $B \rightarrow Hb$  or  $T \rightarrow Ht$ , set the mass limit  $m > 1010 \text{ GeV}$ .
  - <sup>13</sup> AABOUD 18N search for dijet resonance at Atlas with 13 TeV and  $29.3 \text{ fb}^{-1}$ ; limits set on  $m(Z')$  in the mass range of 450–1800 GeV.
  - <sup>14</sup> SIRUNYAN 18DJ search for  $pp \rightarrow ZZ$  or  $WZ \rightarrow \ell\bar{\ell}jj$  resonance at 13 TeV,  $35.9 \text{ fb}^{-1}$ ; no signal; limits set in the 400–4500 GeV mass range, exclusion of  $W'$  up to 2270 GeV in the HVT model A, and up to 2330 GeV for HVT model B. WED bulk graviton exclusion up to 925 GeV.
  - <sup>15</sup> SIRUNYAN 18DY search for  $pp \rightarrow RR; R \rightarrow jj$  two dijet resonances at 13 TeV  $35.9 \text{ fb}^{-1}$ ; no signal; limits placed on RPV top-squark pair production.
  - <sup>16</sup> KHACHATRYAN 17W search for dijet resonance in  $12.9 \text{ fb}^{-1}$  data at 13 TeV; see Fig. 2 for limits on axigluons, diquarks, dark matter mediators etc.
  - <sup>17</sup> KHACHATRYAN 17Y search for  $pp \rightarrow (8-10)j$  in  $19.7 \text{ fb}^{-1}$  at 8 TeV. No signal seen. Limits set on colorons, axigluons, RPV, and SUSY.
  - <sup>18</sup> SIRUNYAN 17F measure  $pp \rightarrow jj$  angular distribution in  $2.6 \text{ fb}^{-1}$  at 13 TeV; limits set on LEDs and quantum black holes.
  - <sup>19</sup> AABOUD 16 search for resonant dijets including one or two  $b$ -jets with  $3.2 \text{ fb}^{-1}$  at 13 TeV; exclude excited  $b^*$  quark from 1.1–2.1 TeV; exclude leptophilic  $Z'$  with SM couplings from 1.1–1.5 TeV.
  - <sup>20</sup> AAD 16N search for  $\geq 3$  jets with  $3.6 \text{ fb}^{-1}$  at 13 TeV; limits placed on micro black holes (Fig. 10) and string balls (Fig. 11).
  - <sup>21</sup> AAD 16S search for high mass jet-jet resonance with  $3.6 \text{ fb}^{-1}$  at 13 TeV; exclude portions of excited quarks,  $W', Z'$  and contact interaction parameter space.
  - <sup>22</sup> KHACHATRYAN 16K search for dijet resonance in  $2.4 \text{ fb}^{-1}$  data at 13 TeV; see Fig. 3 for limits on axigluons, diquarks etc.
  - <sup>23</sup> KHACHATRYAN 16L use data scouting technique to search for  $jj$  resonance on  $18.8 \text{ fb}^{-1}$  of data at 8 TeV. Limits on the coupling of a leptophobic  $Z'$  to quarks are set, improving on the results by other experiments in the mass range between 500–800 GeV.

- <sup>24</sup> AAD 13D search for dijet resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.8 \text{ fb}^{-1}$ . The observed events are compatible with Standard Model expectation. See their Fig. 6 and Table 2 for limits on resonance cross section in the range  $m = 1.0\text{--}4.0$  TeV.
- <sup>25</sup> AALTONEN 13R search for production of a pair of jet-jet resonances in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 6.6 \text{ fb}^{-1}$ . See their Fig. 5 and Tables I, II for cross section limits.
- <sup>26</sup> CHATRCHYAN 13A search for  $qq$ ,  $qg$ , and  $gg$  resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.8 \text{ fb}^{-1}$ . See their Fig. 3 and Table 1 for limits on resonance cross section in the range  $m = 1.0\text{--}4.3$  TeV.
- <sup>27</sup> CHATRCHYAN 13A search for  $b\bar{b}$  resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.8 \text{ fb}^{-1}$ . See their Fig. 8 and Table 4 for limits on resonance cross section in the range  $m = 1.0\text{--}4.0$  TeV.
- <sup>28</sup> AAD 12S search for dijet resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 1.0 \text{ fb}^{-1}$ . See their Fig. 3 and Table 2 for limits on resonance cross section in the range  $m = 0.9\text{--}4.0$  TeV.
- <sup>29</sup> CHATRCHYAN 12BL search for  $t\bar{t}$  resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 4.4 \text{ fb}^{-1}$ . See their Fig. 4 for limits on resonance cross section in the range  $m = 0.5\text{--}3.0$  TeV.
- <sup>30</sup> AAD 11AG search for dijet resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 36 \text{ pb}^{-1}$ . Limits on number of events for  $m = 0.6\text{--}4$  TeV are given in their Table 3.
- <sup>31</sup> AALTONEN 11M find a peak in two jet invariant mass distribution around 140 GeV in  $W + 2$  jet events in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 4.3 \text{ fb}^{-1}$ .
- <sup>32</sup> ABAZOV 11I search for two-jet resonances in  $W + 2$  jet events in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 4.3 \text{ fb}^{-1}$  and give limits  $\sigma < (2.6\text{--}1.3) \text{ pb}$  (95% CL) for  $m = 110\text{--}170$  GeV. The result is incompatible with AALTONEN 11M.
- <sup>33</sup> AAD 10 search for narrow dijet resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 315 \text{ nb}^{-1}$ . Limits on the cross section in the range  $10\text{--}10^3 \text{ pb}$  is given for  $m = 0.3\text{--}1.7$  TeV.
- <sup>34</sup> KHACHATRYAN 10 search for narrow dijet resonances in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 2.9 \text{ pb}^{-1}$ . Limits on the cross section in the range  $1\text{--}300 \text{ pb}$  is given for  $m = 0.5\text{--}2.6$  TeV separately in the final states  $qq$ ,  $qg$ , and  $gg$ .
- <sup>35</sup> ABE 99F search for narrow  $b\bar{b}$  resonances in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8$  TeV. Limits on  $\sigma(p\bar{p} \rightarrow X + \text{anything}) \times \text{B}(X \rightarrow b\bar{b})$  in the range  $3\text{--}10^3 \text{ pb}$  (95%CL) are given for  $m_X = 200\text{--}750$  GeV. See their Table I.
- <sup>36</sup> ABE 97G search for narrow dijet resonances in  $p\bar{p}$  collisions with  $106 \text{ pb}^{-1}$  of data at  $E_{\text{cm}} = 1.8$  TeV. Limits on  $\sigma(p\bar{p} \rightarrow X + \text{anything}) \cdot \text{B}(X \rightarrow jj)$  in the range  $10^4\text{--}10^{-1} \text{ pb}$  (95%CL) are given for dijet mass  $m = 200\text{--}1150$  GeV with both jets having  $|\eta| < 2.0$  and the dijet system having  $|\cos\theta^*| < 0.67$ . See their Table I for the list of limits. Supersedes ABE 93G.
- <sup>37</sup> ABE 93G give cross section times branching ratio into light ( $d$ ,  $u$ ,  $s$ ,  $c$ ,  $b$ ) quarks for  $\Gamma = 0.02 M$ . Their Table II gives limits for  $M = 200\text{--}900$  GeV and  $\Gamma = (0.02\text{--}0.2) M$ .

## LIMITS ON NEUTRAL PARTICLE PRODUCTION

### Production Cross Section of Radiatively-Decaying Neutral Particle

VALUE (pb)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		1 AAD	24AT ATLS	$h \rightarrow \text{ALPs}$ search
		2 ALBERT	18C HAWC	$\gamma$ from Sun
		3 KHACHATRY...17D	CMS	$Z\gamma$ resonance
<0.0008	95	4 AAD	16AI ATLS	$pp \rightarrow \gamma + \text{jet}$

		<sup>5</sup> KHACHATRYAN...16M	CMS	$pp \rightarrow \gamma\gamma$ resonance
<(0.043–0.17)	95	<sup>6</sup> ABBIENDI 00D	OPAL	$e^+e^- \rightarrow X^0 Y^0$ , $X^0 \rightarrow Y^0 \gamma$
<(0.05–0.8)	95	<sup>7</sup> ABBIENDI 00D	OPAL	$e^+e^- \rightarrow X^0 X^0$ , $X^0 \rightarrow Y^0 \gamma$
<(2.5–0.5)	95	<sup>8</sup> ACKERSTAFF 97B	OPAL	$e^+e^- \rightarrow X^0 Y^0$ , $X^0 \rightarrow Y^0 \gamma$
<(1.6–0.9)	95	<sup>9</sup> ACKERSTAFF 97B	OPAL	$e^+e^- \rightarrow X^0 X^0$ , $X^0 \rightarrow Y^0 \gamma$

<sup>1</sup> AAD 24AT search for  $h \rightarrow$  ALPs with  $ALP \rightarrow \gamma\gamma$ . No signal observed. Limits placed in  $BF(h)$  vs.  $m(ALP)$  plane.

<sup>2</sup> ALBERT 18C search for WIMP annihilation in Sun to long-lived, radiatively decaying mediator; no signal; limits set on  $\sigma^{SD}(\chi p)$  assuming long-lived mediator.

<sup>3</sup> KHACHATRYAN 17D search for new scalar resonance decaying to  $Z\gamma$  with  $Z \rightarrow e^+e^-$ ,  $\mu^+\mu^-$  in  $pp$  collisions at 8 and 13 TeV; no signal seen.

<sup>4</sup> AAD 16AI search for excited quarks (EQ) and quantum black holes (QBH) in 3.2 fb<sup>-1</sup> at 13 TeV of data; exclude EQ below 4.4 TeV and QBH below 3.8 (6.2) TeV for RS1 (ADD) models. The visible cross section limit was obtained for 5 TeV resonance with  $\sigma_G/M_G = 2\%$ .

<sup>5</sup> KHACHATRYAN 16M search for  $\gamma\gamma$  resonance using 19.7 fb<sup>-1</sup> at 8 TeV and 3.3 fb<sup>-1</sup> at 13 TeV; slight excess at 750 GeV noted; limit set on RS graviton.

<sup>6</sup> ABBIENDI 00D associated production limit is for  $m_{X^0} = 90$ –188 GeV,  $m_{Y^0}=0$  at  $E_{cm}=189$  GeV. See also their Fig. 9.

<sup>7</sup> ABBIENDI 00D pair production limit is for  $m_{X^0} = 45$ –94 GeV,  $m_{Y^0}=0$  at  $E_{cm}=189$  GeV. See also their Fig. 12.

<sup>8</sup> ACKERSTAFF 97B associated production limit is for  $m_{X^0} = 80$ –160 GeV,  $m_{Y^0}=0$  from 10.0 pb<sup>-1</sup> at  $E_{cm} = 161$  GeV. See their Fig. 3(a).

<sup>9</sup> ACKERSTAFF 97B pair production limit is for  $m_{X^0} = 40$ –80 GeV,  $m_{Y^0}=0$  from 10.0 pb<sup>-1</sup> at  $E_{cm} = 161$  GeV. See their Fig. 3(b).

## Heavy Particle Production Cross Section

VALUE (cm <sup>2</sup> /N)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<sup>1</sup> AAD		24AE ATLS		heavy vector triplet search
<sup>2</sup> AAD		24BV ATLS		VBF di-Higgs to 4 <i>b</i> events
<sup>3</sup> AAD		24CD ATLS		hadronic $W/Z + MET$
<sup>4</sup> AAD		24E ATLS		resonance search
<sup>5</sup> AAD		24S ATLS		QBH via lepton + jet
<sup>6</sup> HAYRAPETY...24AB		CMS		heavy neutral lepton search
<sup>7</sup> HAYRAPETY...24AC		CMS		heavy neutrino from $B$ -decay
<sup>8</sup> HAYRAPETY...24AZ		CMS		four-muon events search
<sup>9</sup> AAD		23P ATLS		exotica search in association with $h \rightarrow \gamma\gamma$
<sup>10</sup> TUMASYAN		23BC CMS		$\gamma$ -jet resonance search
<sup>11</sup> TUMASYAN		23BF CMS		$pp + \gamma/Z + X$ search
<sup>12</sup> TUMASYAN		22AG CMS		SIMP search
<sup>13</sup> AAD		21F ATLS		monojet search
<sup>14</sup> AAIJ		20AL LHCB		$pp$ at 13 TeV, dimuon resonance
<sup>15</sup> SIRUNYAN		20AY CMS		$\Upsilon(1S)\mu^+\mu^-$ decay states

		16	SIRUNYAN	20Z	CMS	multilepton BSM search, 13 TeV
		17	AABOUD	19H	ATLS	di-photon-jet resonance
		18	AABOUD	19V	ATLS	review, mediator-based DM
		19	SIRUNYAN	19O	CMS	$pp \rightarrow \gamma \cancel{E}_T$
		20	AABOUD	18CJ	ATLS	$pp \rightarrow VV/\ell\ell/\ell\nu$ , $V =$ $W, Z, h$
		21	AABOUD	18CM	ATLS	$pp \rightarrow e\mu/e\tau/\mu\tau$
		22	AAIJ	18AJ	LHCB	$pp \rightarrow A' \rightarrow \mu^+\mu^-$ ; dark photon
		23	BANERJEE	18	NA64	$eZ \rightarrow eZ X(A')$
		24	BANERJEE	18A	NA64	$eZ \rightarrow eZ A'$ , $A' \rightarrow \chi\chi$
		25	MARSICANO	18	E137	$e^+e^- \rightarrow A'(\gamma)$ visible decay
		26	SIRUNYAN	18BB	CMS	$pp \rightarrow Z' \rightarrow \ell^+\ell^-$ at 13 TeV
		27	SIRUNYAN	18DA	CMS	$pp \rightarrow$ Black Hole, string ball, sphaleron
		28	SIRUNYAN	18DD	CMS	$pp \rightarrow jj$
		29	SIRUNYAN	18DR	CMS	$pp \rightarrow b\mu\bar{\mu}$
		30	SIRUNYAN	18DU	CMS	$pp \rightarrow \gamma\gamma$
		31	SIRUNYAN	18ED	CMS	$pp \rightarrow V \rightarrow Wh$ ; $h \rightarrow$ $b\bar{b}$ ; $W \rightarrow \ell\nu$
		32	AABOUD	17B	ATLS	$WH, ZH$ resonance
		33	AAIJ	17BR	LHCB	$pp \rightarrow \pi_V\pi_V$ , $\pi_V \rightarrow jj$
		34	AAD	16O	ATLS	$\ell + (\ell s \text{ or jets})$
		35	AAD	16R	ATLS	$WW, WZ, ZZ$ resonance
		36	KRASZNAHO...	16		$p^7\text{Li} \rightarrow ^8\text{Be} \rightarrow X(17)N$ , $X(17) \rightarrow e^+e^-$
		37	LEES	15E	BABR	$e^+e^-$ collisions
		38	ADAMS	97B	KTEV	$m = 1.2\text{--}5$ GeV
$< 10^{-36}\text{--}10^{-33}$	90	39	GALLAS	95	TOF	$m = 0.5\text{--}20$ GeV
$< (4\text{--}0.3) \times 10^{-31}$	95	40	AKESSON	91	CNTR	$m = 0\text{--}5$ GeV
$< 2 \times 10^{-36}$	90	41	BADIER	86	BDMP	$\tau = (0.05\text{--}1.) \times 10^{-8}\text{s}$
$< 2.5 \times 10^{-35}$		42	GUSTAFSON	76	CNTR	$\tau > 10^{-7}$ s

<sup>1</sup> AAD 24AE search for heavy vector triplet production with decay to boson pairs. No signal was observed. Limits placed in  $\sigma$  vs  $m$  plane. Limits also placed in various two-dimensional coupling planes ( $g_F$ ,  $g_H$ ,  $g_\ell$ ,  $g_\ell$ (3rd Gen),  $g_q$ ,  $g_q$ (3rd Gen),  $g_q$ (1st/2nd Gen)).

<sup>2</sup> AAD 24BV search for VBF di-Higgs production with decay to boosted  $4b$  state. No signal observed. Limits placed in mass vs. cross section plane for various simplified models.

<sup>3</sup> AAD 24CD search for hadronically-decaying  $W/Z + \text{MET}$  events from new physics with  $140 \text{ fb}^{-1}$  at 13 TeV. No signal observed. Limits placed on various simplified new physics models.

<sup>4</sup> AAD 24E uses a new resonance search technique for two-body decays into any pair of  $\ell$ ,  $b$ , and jet with  $140 \text{ fb}^{-1}$  of data. No signal was observed. Limits placed in  $\sigma$  vs mass plane for various decay modes.

<sup>5</sup> AAD 24S search for quantum black hole (QBH) decay to lepton + jet in  $140 \text{ fb}^{-1}$  of data. No signal observed. Limits placed in  $\sigma \cdot \text{BF}$  vs mass plane for ADD and RS models.

<sup>6</sup> HAYRAPETYAN 24AB search for heavy neutral leptons  $N$  at CMS with  $138 \text{ fb}^{-1}$  of data. No signal observed. Limits placed in mixing angle vs  $m(N)$  plane.

<sup>7</sup> HAYRAPETYAN 24AC search for heavy long-lived neutrino  $N$  produced in  $B$ -decays in  $41.6 \text{ fb}^{-1}$  of data. No signal observed. Limits placed in mixing angle vs  $m(N)$  plane.

- <sup>8</sup> HAYRAPETYAN 24AZ search for various new bosons via production and decay to four muon states with 41.5 and 59.7 fb<sup>-1</sup> of data at 13 TeV. No signal observed. Limits placed usually in mass vs. cross section plane for a variety of new physics simplified models.
- <sup>9</sup> AAD 23P search in 22 channels for exotica produced in association with  $h \rightarrow \gamma\gamma$  in 139 fb<sup>-1</sup> of data. No signal observed. Limits placed on production cross section in various channels.
- <sup>10</sup> TUMASYAN 23BC search for  $\gamma$ -jet resonance at CMS with 138 fb<sup>-1</sup> of data. No signal observed. Limits placed on quantum black hole and excited quark models.
- <sup>11</sup> TUMASYAN 23BF search for  $pp \rightarrow pp + \gamma/Z + X$  search where  $X$  is missing particle using CMS-TOTEM with 37.2 fb<sup>-1</sup> of data. No signal observed. Limits placed on  $\sigma$  vs.  $m$  plane.
- <sup>12</sup> TUMASYAN 22AG search for strongly interacting neutral massive particles via trackless jets with 16.1 fb<sup>-1</sup> at 13 TeV; no signal detected; limits placed in mass vs. cross section plane for various simplified models.
- <sup>13</sup> AAD 21F search for hard monojet production at ATLAS with 139 fb<sup>-1</sup> of 13 TeV data. No signal observed. Limits placed on invisible production cross-section recoiling against ISR and interpreted in variety of BSM models.
- <sup>14</sup> AAIJ 20AL search for dimuon resonance from promptly decaying  $X$  particle. No signal detected. Limits placed on  $m(X)$  up to 60 GeV depending on mixing in 2HDM.
- <sup>15</sup> SIRUNYAN 20AY measured  $\Upsilon(1S)$  pair production cross section and searched for new states decaying into  $\Upsilon(1S)\mu^+\mu^-$  at CMS with 13 TeV with 35.9 fb<sup>-1</sup>. No signal is found and limits are set in  $\sigma \cdot \text{BF}$  vs. mass plane for tetra- $b$ -quarks with masses between 17.5 and 19 GeV and for generic search for narrow resonances with mass between 16.5 and 27 GeV.
- <sup>16</sup> SIRUNYAN 20Z search for BSM physics via multilepton production with CMS at 13 TeV with 137 fb<sup>-1</sup>; no signal is found and limits are set on type-III seesaw and other BSM models.
- <sup>17</sup> AABOUD 19H searches for di-photon-jet resonance at 13 TeV and 36.7 fb<sup>-1</sup> of data; no signal found and limits placed on  $\sigma \cdot \text{BR}$  vs. mass plane for various simplified models.
- <sup>18</sup> AABOUD 19V review ATLAS searches for mediator-based DM at 7, 8, and 13 TeV with up to 37 fb<sup>-1</sup> of data; no signal found and limits set for wide variety of simplified models of dark matter.
- <sup>19</sup> SIRUNYAN 19O search for  $pp \rightarrow \gamma \cancel{E}_T$  at 13 TeV with 36.1 fb<sup>-1</sup>; no signal found and limits set for various simplified models.
- <sup>20</sup> AABOUD 18CJ make multichannel search for  $pp \rightarrow VV/\ell\ell/\ell\nu$ ,  $V = W, Z, h$  at 13 TeV, 36.1 fb<sup>-1</sup>; no signal found; limits placed for several BSM models.
- <sup>21</sup> AABOUD 18CM search for lepton-flavor violating resonance in  $pp \rightarrow e\mu/e\tau/\mu\tau$  at 13 TeV, 36.1 fb<sup>-1</sup>; no signal is found and limits placed for various BSM models.
- <sup>22</sup> AAIJ 18AJ search for prompt and delayed dark photon decay  $A' \rightarrow \mu^+\mu^-$  at LHCb detector using 1.6 fb<sup>-1</sup> of  $pp$  collisions at 13 TeV; limits on  $m(A')$  vs. kinetic mixing are set.
- <sup>23</sup> BANERJEE 18 search for dark photon  $A'/16.7$  MeV boson  $X$  at NA64 via  $eZ \rightarrow eZX(A')$ ; no signal found and limits set on the  $X$ - $e^-$  coupling  $\epsilon_e$  in the range  $1.3 \times 10^{-4} \leq \epsilon_e \leq 4.2 \times 10^{-4}$  excluding part of the allowed parameter space.
- <sup>24</sup> BANERJEE 18A search for invisibly decaying dark photons in  $eZ \rightarrow eZA'$ ,  $A' \rightarrow$  invisible; no signal found and limits set on mixing for  $m(A') < 1$  GeV.
- <sup>25</sup> MARSICANO 18 search for dark photon  $e^+e^- \rightarrow A'(\gamma)$  visible decay in SLAC E137  $e$  beam dump data. No signal observed and limits set in  $\epsilon$  coupling vs  $m(A')$  plane, see their figure 7.
- <sup>26</sup> SIRUNYAN 18BB search for high mass dilepton resonance; no signal found and exclude portions of  $p$ -space of  $Z'$ , KK graviton models.
- <sup>27</sup> SIRUNYAN 18DA search for  $pp \rightarrow$  Black Hole, string ball, sphaleron via high multiplicity events at 13 TeV, 35.9 fb<sup>-1</sup>; no signal, require e.g.  $m(\text{BH}) > 10.1$  TeV.

- <sup>28</sup> SIRUNYAN 18DD search for  $pp \rightarrow jj$  deviations in dijet angular distribution. No signal observed. Set limits on large extra dimensions, black holes and DM mediators e.g.  $m(\text{BH}) > 5.9\text{--}8.2$  TeV.
- <sup>29</sup> SIRUNYAN 18DR search for dimuon resonance in  $pp \rightarrow b\mu\bar{\mu}$  at 8 and 13 TeV. Slight excess seen at  $m(\mu\bar{\mu}) \sim 28$  GeV in some channels.
- <sup>30</sup> SIRUNYAN 18DU search for high mass diphoton resonance in  $pp \rightarrow \gamma\gamma$  at 13 TeV using  $35.9 \text{ fb}^{-1}$ ; no signal; limits placed on RS Graviton, LED, and clockwork.
- <sup>31</sup> SIRUNYAN 18ED search for  $pp \rightarrow V \rightarrow Wh; h \rightarrow b\bar{b}; W \rightarrow \ell\nu$  at 13 TeV with  $35.9 \text{ fb}^{-1}$ ; no signal; limits set on  $m(W') > 2.9$  TeV.
- <sup>32</sup> AABOUD 17B exclude  $m(W', Z') < 1.49\text{--}2.31$  TeV depending on the couplings and  $W'/Z'$  degeneracy assumptions via  $WH, ZH$  search in  $pp$  collisions at 13 TeV with  $3.2 \text{ fb}^{-1}$  of data.
- <sup>33</sup> AAIJ 17BR search for long-lived hidden valley pions from Higgs decay. Limits are set on the signal strength as a function of the mass and lifetime of the long-lived particle in their Fig. 4 and Tab. 4.
- <sup>34</sup> AAD 16O search for high  $E_T \ell + (\ell\text{s or jets})$  with  $3.2 \text{ fb}^{-1}$  at 13 TeV; exclude micro black holes mass  $< 8$  TeV (Fig. 3) for models with two extra dimensions.
- <sup>35</sup> AAD 16R search for  $WW, WZ, ZZ$  resonance in  $20.3 \text{ fb}^{-1}$  at 8 TeV data; limits placed on massive RS graviton (Fig. 4).
- <sup>36</sup> KRASZNAHORKAY 16 report  $p\text{Li} \rightarrow \text{Be} \rightarrow e\bar{e}N$   $5\sigma$  resonance at 16.7 MeV— possible evidence for nuclear interference or new light boson. However, such nuclear interference was ruled out already by ZANG 17.
- <sup>37</sup> LEES 15E search for long-lived neutral particles produced in  $e^+e^-$  collisions in the Upsilon region, which decays into  $e^+e^-, \mu^+\mu^-, e^\pm\mu^\mp, \pi^+\pi^-, K^+K^-,$  or  $\pi^\pm K^\mp$ . See their Fig. 2 for cross section limits.
- <sup>38</sup> ADAMS 97B search for a hadron-like neutral particle produced in  $pN$  interactions, which decays into a  $\rho^0$  and a weakly interacting massive particle. Upper limits are given for the ratio to  $K_L$  production for the mass range 1.2–5 GeV and lifetime  $10^{-9}\text{--}10^{-4}$  s. See also our Light Gluino Section.
- <sup>39</sup> GALLAS 95 limit is for a weakly interacting neutral particle produced in 800 GeV/ $c$   $pN$  interactions decaying with a lifetime of  $10^{-4}\text{--}10^{-8}$  s. See their Figs. 8 and 9. Similar limits are obtained for a stable particle with interaction cross section  $10^{-29}\text{--}10^{-33} \text{ cm}^2$ . See Fig. 10.
- <sup>40</sup> AKESSON 91 limit is from weakly interacting neutral long-lived particles produced in  $pN$  reaction at 450 GeV/ $c$  performed at CERN SPS. Bourquin-Gaillard formula is used as the production model. The above limit is for  $\tau > 10^{-7}$  s. For  $\tau > 10^{-9}$  s,  $\sigma < 10^{-30} \text{ cm}^2/\text{nucleon}$  is obtained.
- <sup>41</sup> BADIER 86 looked for long-lived particles at 300 GeV  $\pi^-$  beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass  $> 2$  GeV. The limit applies for particle modes,  $\mu^+\pi^-, \mu^+\mu^-, \pi^+\pi^-X, \pi^+\pi^-\pi^\pm$  etc. See their figure 5 for the contours of limits in the mass- $\tau$  plane for each mode.
- <sup>42</sup> GUSTAFSON 76 is a 300 GeV FNAL experiment looking for heavy ( $m > 2$  GeV) long-lived neutral hadrons in the M4 neutral beam. The above typical value is for  $m = 3$  GeV and assumes an interaction cross section of 1 mb. Values as a function of mass and interaction cross section are given in figure 2.

## Production of New Penetrating Non- $\nu$ Like States in Beam Dump

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<sup>1</sup> ANDREEV	24	NA64	$\mu + E(\text{missing})$ search
<sup>2</sup> ABRATENKO	22A	MCBN	search for LLPs
<sup>3</sup> ANDREEV	22A	NA64	new boson $X$ in $eZ \rightarrow eZX$

<sup>4</sup> ANDREEV	21	NA64	new boson $X$ in $eZ \rightarrow eZX$
<sup>5</sup> LOSECCO	81	CALO	28 GeV protons

- <sup>1</sup> ANDREEV 24 search for  $\mu \rightarrow \mu + E(\text{missing})$ . No signal observed. Limits placed on  $Z'$  models and DM models.
- <sup>2</sup> ABRATENKO 22A search for LLPs from kaon decay in MicroBooNE absorber; no signal observed; limits placed for heavy neutral leptons (HNLs) and Higgs portal scalars (HPSs) in the MeV mass range.
- <sup>3</sup> ANDREEV 22A search for new light B-L gauge boson  $Z' \rightarrow \nu\bar{\nu}$  in electron beam dump at NA64; no signal observed; limits set in  $m(Z')$  vs coupling plane for  $m(Z') \sim 10^{-6}$ –1 GeV.
- <sup>4</sup> ANDREEV 21 search for new invisibly decaying boson  $X$  in  $eZ \rightarrow eZX$  at NA64. No signal observed. Limits set in coupling vs.  $m(X)$  plane for  $m(X) \sim 10^{-3}$  to 1 GeV.
- <sup>5</sup> No excess neutral-current events leads to  $\sigma(\text{production}) \times \sigma(\text{interaction}) \times \text{acceptance} < 2.26 \times 10^{-71} \text{ cm}^4/\text{nucleon}^2$  (CL = 90%) for light neutrals. Acceptance depends on models (0.1 to  $4. \times 10^{-4}$ ).

## LIMITS ON CHARGED PARTICLES IN $e^+e^-$

### Heavy Particle Production Cross Section in $e^+e^-$

Ratio to  $\sigma(e^+e^- \rightarrow \mu^+\mu^-)$  unless noted. See also entries in Free Quark Search and Magnetic Monopole Searches.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1 \times 10^{-3}$	90	<sup>1</sup> ADACHI	23K BELL	search for LLP in $B$ decays
		<sup>2</sup> KILE	18 ALEP	$e^+e^- \rightarrow 4 \text{ jets}$
		<sup>3</sup> ABLIKIM	17AA BES3	$e^+e^- \rightarrow \ell\bar{\ell}\gamma$
		<sup>4</sup> ACKERSTAFF	98P OPAL	$Q=1,2/3$ , $m=45$ –89.5 GeV
		<sup>5</sup> ABREU	97D DLPH	$Q=1,2/3$ , $m=45$ –84 GeV
$<2 \times 10^{-5}$	95	<sup>6</sup> BARATE	97K ALEP	$Q=1$ , $m=45$ –85 GeV
		<sup>7</sup> AKERS	95R OPAL	$Q=1$ , $m=5$ –45 GeV
$<1 \times 10^{-5}$	95	<sup>7</sup> AKERS	95R OPAL	$Q=2$ , $m=5$ –45 GeV
$<2 \times 10^{-3}$	90	<sup>8</sup> BUSKULIC	93C ALEP	$Q=1$ , $m=32$ –72 GeV
$<(10^{-2}-1)$	95	<sup>9</sup> ADACHI	90C TOPZ	$Q=1$ , $m=1$ –16, 18–27 GeV
$<7 \times 10^{-2}$	90	<sup>10</sup> ADACHI	90E TOPZ	$Q=1$ , $m=5$ –25 GeV
$<1.6 \times 10^{-2}$	95	<sup>11</sup> KINOSHITA	82 PLAS	$Q=3$ –180, $m < 14.5$ GeV
$<5.0 \times 10^{-2}$	90	<sup>12</sup> BARTEL	80 JADE	$Q=(3,4,5)/3$ 2–12 GeV

- <sup>1</sup> ADACHI 23K search for spin-0 LLP called  $S$  in  $B$  decays. No signal observed. Limits placed in branching fraction vs.  $m(S)$  plane.
- <sup>2</sup> KILE 18 investigate archived ALEPH  $e^+e^- \rightarrow 4 \text{ jets}$  data and see 4–5  $\sigma$  excess at 110 GeV.
- <sup>3</sup> ABLIKIM 17AA search for dark photon  $A \rightarrow \ell\bar{\ell}$  at 3.773 GeV with  $2.93 \text{ fb}^{-1}$ . Limits are set in  $\epsilon$  vs  $m(A)$  plane.
- <sup>4</sup> ACKERSTAFF 98P search for pair production of long-lived charged particles at  $E_{\text{cm}}$  between 130 and 183 GeV and give limits  $\sigma < (0.05\text{--}0.2) \text{ pb}$  (95%CL) for spin-0 and spin-1/2 particles with  $m=45$ –89.5 GeV, charge 1 and 2/3. The limit is translated to the cross section at  $E_{\text{cm}}=183$  GeV with the  $s$  dependence described in the paper. See their Figs. 2–4.
- <sup>5</sup> ABREU 97D search for pair production of long-lived particles and give limits  $\sigma < (0.4\text{--}2.3) \text{ pb}$  (95%CL) for various center-of-mass energies  $E_{\text{cm}}=130$ –136, 161, and 172 GeV, assuming an almost flat production distribution in  $\cos\theta$ .
- <sup>6</sup> BARATE 97K search for pair production of long-lived charged particles at  $E_{\text{cm}} = 130$ , 136, 161, and 172 GeV and give limits  $\sigma < (0.2\text{--}0.4) \text{ pb}$  (95%CL) for spin-0 and spin-1/2

particles with  $m=45\text{--}85$  GeV. The limit is translated to the cross section at  $E_{\text{cm}}=172$  GeV with the  $E_{\text{cm}}$  dependence described in the paper. See their Figs. 2 and 3 for limits on  $J = 1/2$  and  $J = 0$  cases.

- <sup>7</sup> AKERS 95R is a CERN-LEP experiment with  $W_{\text{cm}} \sim m_Z$ . The limit is for the production of a stable particle in multihadron events normalized to  $\sigma(e^+e^- \rightarrow \text{hadrons})$ . Constant phase space distribution is assumed. See their Fig. 3 for bounds for  $Q = \pm 2/3, \pm 4/3$ .
- <sup>8</sup> BUSKULIC 93C is a CERN-LEP experiment with  $W_{\text{cm}} = m_Z$ . The limit is for a pair or single production of heavy particles with unusual ionization loss in TPC. See their Fig. 5 and Table 1.
- <sup>9</sup> ADACHI 90C is a KEK-TRISTAN experiment with  $W_{\text{cm}} = 52\text{--}60$  GeV. The limit is for pair production of a scalar or spin-1/2 particle. See Figs. 3 and 4.
- <sup>10</sup> ADACHI 90E is KEK-TRISTAN experiment with  $W_{\text{cm}} = 52\text{--}61.4$  GeV. The above limit is for inclusive production cross section normalized to  $\sigma(e^+e^- \rightarrow \mu^+\mu^-) \cdot \beta(3 - \beta^2)/2$ , where  $\beta = (1 - 4m^2/W_{\text{cm}}^2)^{1/2}$ . See the paper for the assumption about the production mechanism.
- <sup>11</sup> KINOSHITA 82 is SLAC PEP experiment at  $W_{\text{cm}} = 29$  GeV using lexan and  $^{39}\text{Cr}$  plastic sheets sensitive to highly ionizing particles.
- <sup>12</sup> BARTEL 80 is DESY-PETRA experiment with  $W_{\text{cm}} = 27\text{--}35$  GeV. Above limit is for inclusive pair production and ranges between  $1. \times 10^{-1}$  and  $1. \times 10^{-2}$  depending on mass and production momentum distributions. (See their figures 9, 10, 11).

### Branching Fraction of $Z^0$ to a Pair of Stable Charged Heavy Fermions

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 5 \times 10^{-6}$	95	<sup>1</sup> AKERS	95R OPAL	$m = 40.4\text{--}45.6$ GeV
$< 1 \times 10^{-3}$	95	AKRAWY	900 OPAL	$m = 29\text{--}40$ GeV
<sup>1</sup> AKERS 95R give the 95% CL limit $\sigma(X\bar{X})/\sigma(\mu\mu) < 1.8 \times 10^{-4}$ for the pair production of singly- or doubly-charged stable particles. The limit applies for the mass range 40.4–45.6 GeV for $X^\pm$ and $< 45.6$ GeV for $X^{\pm\pm}$ . See the paper for bounds for $Q = \pm 2/3, \pm 4/3$ .				

## LIMITS ON CHARGED PARTICLES IN HADRONIC REACTIONS

### MASS LIMITS for Long-Lived Charged Heavy Fermions

Limits are for spin 1/2 particles with no color and  $SU(2)_L$  charge. The electric charge  $Q$  of the particle (in the unit of  $e$ ) is therefore equal to its weak hypercharge. Pair production by Drell-Yan like  $\gamma$  and  $Z$  exchange is assumed to derive the limits.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		<sup>1</sup> BARAK	24 SENS	millicharged particles
		<sup>2</sup> HAYRAPETY...	24AD CMS	emerging dark jet
		<sup>3</sup> AAD	23BT ATLS	multi-charged LLP
		<sup>4</sup> SIRUNYAN	20N CMS	disappearing track LLP
$> 660$	95	<sup>5</sup> AAD	15BJ ATLS	$ Q  = 2$
$> 200$	95	<sup>6</sup> CHATRCHYAN	13AB CMS	$ Q  = 1/3$
$> 480$	95	<sup>6</sup> CHATRCHYAN	13AB CMS	$ Q  = 2/3$
$> 574$	95	<sup>6</sup> CHATRCHYAN	13AB CMS	$ Q  = 1$
$> 685$	95	<sup>6</sup> CHATRCHYAN	13AB CMS	$ Q  = 2$
$> 140$	95	<sup>7</sup> CHATRCHYAN	13AR CMS	$ Q  = 1/3$
$> 310$	95	<sup>7</sup> CHATRCHYAN	13AR CMS	$ Q  = 2/3$

- <sup>1</sup> BARAK 24 search for milli-charged particle production in  $p$ -graphite collisions using skipper CCDs. No signal was observed. Limits at 95% placed in charge vs. mass plane in a wide range of masses in the MeV range.

- <sup>2</sup> HAYRAPETYAN 24AD search for emerging dark jets from dark mediator pair production in  $138 \text{ fb}^{-1}$  of data at 13 TeV. No signal observed. Limits placed in the dark  $\tau(\pi_d)$  vs  $m(\text{mediator})$  plane.
- <sup>3</sup> AAD 23BT search for multi-charged long-lived particles with ATLAS detector using  $139 \text{ fb}^{-1}$ . No signal observed. Limits placed on LLP mass vs. charge plane.
- <sup>4</sup> SIRUNYAN 20N search for LLPs using disappearing track signature at CMS at 13 TeV with  $101 \text{ fb}^{-1}$ ; no signal; limits placed on long-lived winos and higgsinos from SUSY depending on mass and lifetime: e.g. at 95% CL, for a purely higgsino neutralino,  $m(\text{chargino}) > 750$  (175) GeV for  $\tau = 3$  (0.05) ns, and for a purely wino neutralino,  $m(\text{chargino}) > 884$  (474) GeV for  $\tau = 3$  (0.2) ns.
- <sup>5</sup> AAD 15BJ use  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See paper for limits for  $|Q| = 3, 4, 5, 6$ .
- <sup>6</sup> CHATRCHYAN 13AB use  $5.0 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $18.8 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . See paper for limits for  $|Q| = 3, 4, \dots, 8$ .
- <sup>7</sup> CHATRCHYAN 13AR use  $5.0 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ .

## Heavy Particle Production Cross Section

VALUE (nb)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		<sup>1</sup> AAD	24AK ATLS	top + MET search
		<sup>2</sup> AAD	24B ATLS	non-resonant jet search
		<sup>3</sup> AAD	24CM ATLS	dark meson $\rightarrow t/b$ search
		<sup>4</sup> AAD	22G ATLS	vector-like matter search
		<sup>5</sup> TUMASYAN	22H CMS	search for new matter via multileptons
		<sup>6</sup> SIRUNYAN	21T CMS	model independent search
		<sup>7</sup> SIRUNYAN	20C CMS	$4t$ search via multileptons
		<sup>8</sup> AABOUD	19AA ATLS	BSM search
		<sup>9</sup> AABOUD	19Q ATLS	single top + MET
		<sup>10</sup> AABOUD	17D ATLS	anomalous $WWjj$ , $WZjj$
		<sup>11</sup> AABOUD	17L ATLS	$m > 870 \text{ GeV}$ , $Z(\rightarrow \nu\nu)tX$
		<sup>12</sup> SIRUNYAN	17B CMS	$tH$
		<sup>13</sup> SIRUNYAN	17C CMS	$Z + (t \text{ or } b)$
		<sup>14</sup> SIRUNYAN	17J CMS	$X_{5/3} \rightarrow tW$
		<sup>15</sup> AAIJ	15BD LHCb	$m=124\text{--}309 \text{ GeV}$
		<sup>16</sup> AAD	13AH ATLS	$ q =(2\text{--}6)e$ , $m=50\text{--}600 \text{ GeV}$
$<1.2 \times 10^{-3}$	95	<sup>17</sup> AAD	11I ATLS	$ q =10e$ , $m=0.2\text{--}1 \text{ TeV}$
$<1.0 \times 10^{-5}$	95	<sup>18,19</sup> AALTONEN	09Z CDF	$m > 100 \text{ GeV}$ , noncolored
$<4.8 \times 10^{-5}$	95	<sup>18,20</sup> AALTONEN	09Z CDF	$m > 100 \text{ GeV}$ , colored
$<0.31\text{--}0.04 \times 10^{-3}$	95	<sup>21</sup> ABAZOV	09M D0	pair production
$<0.19$	95	<sup>22</sup> AKTAS	04C H1	$m=3\text{--}10 \text{ GeV}$
$<0.05$	95	<sup>23</sup> ABE	92J CDF	$m=50\text{--}200 \text{ GeV}$
$<30\text{--}130$		<sup>24</sup> CARROLL	78 SPEC	$m=2\text{--}2.5 \text{ GeV}$
$<100$		<sup>25</sup> LEIPUNER	73 CNTR	$m=3\text{--}11 \text{ GeV}$

- <sup>1</sup> AAD 24AK search for single top + MET in  $140 \text{ fb}^{-1}$  of data. No signal observed. Limits placed in various planes such as  $\sigma$  vs  $m(\text{mediator})$  for assumed simplified models.
- <sup>2</sup> AAD 24B search for non-resonant jets + MET at  $\sqrt{s} = 13 \text{ TeV}$  with  $139 \text{ fb}^{-1}$ . No excess observed. Limits placed on dark sector model mediator mass and coupling.
- <sup>3</sup> AAD 24CM search for new dark mesons decaying to  $t/b$  states with  $140 \text{ fb}^{-1}$  at 13 TeV. No signal observed. Limits placed in mass vs.  $\sigma$  plane for various new physics simplified models.

- <sup>4</sup> AAD 22G search for single vector-like quark  $T$  with  $T \rightarrow t h$  in all hadronic mode with  $139 \text{ fb}^{-1}$  at 13 TeV; no signal observed; limits placed in mass vs. coupling plane.
- <sup>5</sup> TUMASYAN 22H search for new states of matter via non-resonant multilepton production based on a luminosity of  $138 \text{ fb}^{-1}$ ; no signal observed; limits placed on vector-like leptons, leptoquarks, and new fermions from type-III seesaw model.
- <sup>6</sup> SIRUNYAN 21T perform model unspecific search for deviations from SM with CMS at 13 TeV with  $35.9^{-1} \text{ fb}$  data in numerous signature channels. No deviations from SM found.
- <sup>7</sup> SIRUNYAN 20C search for four top-quark production with decay to multileptons at CMS at 13 TeV with  $137 \text{ fb}^{-1}$ ; no signal is found and limits are placed on the Higgs boson oblique parameter in the effective field theory framework (EFT) and the model parameters ( $\tan\beta$ ).
- <sup>8</sup> AABOUD 19AA search for BSM physics at 13 TeV with  $3.2 \text{ fb}^{-1}$  in  $> 10^5$  regions of  $> 700$  event classes; no significant signal found.
- <sup>9</sup> AABOUD 19Q search for single top+MET events at 13 TeV with  $36.1 \text{ fb}^{-1}$  of data; no signal found and limits set in  $\sigma$  or coupling vs. mass plane for variety of simplified models including DM and vector-like top quark  $T$ .
- <sup>10</sup> AABOUD 17D search for  $W W j j$ ,  $W Z j j$  in  $p p$  collisions at 8 TeV with  $3.2 \text{ fb}^{-1}$ ; set limits on anomalous couplings.
- <sup>11</sup> AABOUD 17L search for the pair production of heavy vector-like  $T$  quarks in the  $Z(\rightarrow \nu \nu) t X$  final state.
- <sup>12</sup> SIRUNYAN 17B search for vector-like quark  $p p \rightarrow T X \rightarrow t H X$  in  $2.3 \text{ fb}^{-1}$  at 13 TeV; no signal seen; limits placed.
- <sup>13</sup> SIRUNYAN 17C search for vector-like quark  $p p \rightarrow T X \rightarrow Z + (t \text{ or } b)$  in  $2.3 \text{ fb}^{-1}$  at 13 TeV; no signal seen; limits placed.
- <sup>14</sup> SIRUNYAN 17J search for  $p p \rightarrow X_{5/3} X_{5/3} \rightarrow t W t W$  with  $2.3 \text{ fb}^{-1}$  at 13 TeV. No signal seen:  $m(X) > 1020$  (990) GeV for RH (LH) new charge 5/3 quark.
- <sup>15</sup> AAIJ 15BD search for production of long-lived particles in  $p p$  collisions at  $E_{\text{cm}} = 7$  and 8 TeV. See their Table 6 for cross section limits.
- <sup>16</sup> AAD 13AH search for production of long-lived particles with  $|q|=(2-6)e$  in  $p p$  collisions at  $E_{\text{cm}} = 7$  TeV with  $4.4 \text{ fb}^{-1}$ . See their Fig. 8 for cross section limits.
- <sup>17</sup> AAD 11I search for production of highly ionizing massive particles in  $p p$  collisions at  $E_{\text{cm}} = 7$  TeV with  $L = 3.1 \text{ pb}^{-1}$ . See their Table 5 for similar limits for  $|q| = 6e$  and  $17e$ , Table 6 for limits on pair production cross section.
- <sup>18</sup> AALTONEN 09Z search for long-lived charged particles in  $p \bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 1.0 \text{ fb}^{-1}$ . The limits are on production cross section for a particle of mass above 100 GeV in the region  $|\eta| \lesssim 0.7$ ,  $p_T > 40$  GeV, and  $0.4 < \beta < 1.0$ .
- <sup>19</sup> Limit for weakly interacting charge-1 particle.
- <sup>20</sup> Limit for up-quark like particle.
- <sup>21</sup> ABAZOV 09M search for pair production of long-lived charged particles in  $p \bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV with  $L = 1.1 \text{ fb}^{-1}$ . Limit on the cross section of (0.31–0.04) pb (95% CL) is given for the mass range of 60–300 GeV, assuming the kinematics of stau pair production.
- <sup>22</sup> AKTAS 04C look for charged particle photoproduction at HERA with mean c.m. energy of 200 GeV.
- <sup>23</sup> ABE 92J look for pair production of unit-charged particles which leave detector before decaying. Limit shown here is for  $m=50$  GeV. See their Fig. 5 for different charges and stronger limits for higher mass.
- <sup>24</sup> CARROLL 78 look for neutral,  $S = -2$  dihyperon resonance in  $p p \rightarrow 2K^+ X$ . Cross section varies within above limits over mass range and  $p_{\text{lab}} = 5.1\text{--}5.9 \text{ GeV}/c$ .
- <sup>25</sup> LEIPUNER 73 is an NAL 300 GeV  $p$  experiment. Would have detected particles with lifetime greater than 200 ns.

**Heavy Particle Production Differential Cross Section**

VALUE ( $\text{cm}^2\text{sr}^{-1}\text{GeV}^{-1}$ )	CL%	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
		<sup>1</sup> HAYRAPETY...23F	CMS		top $\rightarrow$ $\ell$ s via EFT ops.
$<2.6 \times 10^{-36}$	90	<sup>2</sup> BALDIN 76	CNTR	–	$Q=1$ , $m=2.1\text{--}9.4$ GeV
$<2.2 \times 10^{-33}$	90	<sup>3</sup> ALBROW 75	SPEC	$\pm$	$Q= \pm 1$ , $m=4\text{--}15$ GeV
$<1.1 \times 10^{-33}$	90	<sup>3</sup> ALBROW 75	SPEC	$\pm$	$Q= \pm 2$ , $m=6\text{--}27$ GeV
$<8. \times 10^{-35}$	90	<sup>4</sup> JOVANO... 75	CNTR	$\pm$	$m=15\text{--}26$ GeV
$<1.5 \times 10^{-34}$	90	<sup>4</sup> JOVANO... 75	CNTR	$\pm$	$Q= \pm 2$ , $m=3\text{--}10$ GeV
$<6. \times 10^{-35}$	90	<sup>4</sup> JOVANO... 75	CNTR	$\pm$	$Q= \pm 2$ , $m=10\text{--}26$ GeV
$<1. \times 10^{-31}$	90	<sup>5</sup> APPEL 74	CNTR	$\pm$	$m=3.2\text{--}7.2$ GeV
$<5.8 \times 10^{-34}$	90	<sup>6</sup> ALPER 73	SPEC	$\pm$	$m=1.5\text{--}24$ GeV
$<1.2 \times 10^{-35}$	90	<sup>7</sup> ANTIPOV 71B	CNTR	–	$Q=-$ , $m=2.2\text{--}2.8$
$<2.4 \times 10^{-35}$	90	<sup>8</sup> ANTIPOV 71C	CNTR	–	$Q=-$ , $m=1.2\text{--}1.7$ , 2.1–4
$<2.4 \times 10^{-35}$	90	BINON 69	CNTR	–	$Q=-$ , $m=1\text{--}1.8$ GeV
$<1.5 \times 10^{-36}$		<sup>9</sup> DORFAN 65	CNTR		Be target $m=3\text{--}7$ GeV
$<3.0 \times 10^{-36}$		<sup>9</sup> DORFAN 65	CNTR		Fe target $m=3\text{--}7$ GeV

<sup>1</sup> HAYRAPETYAN 23F search for anomalous top  $\rightarrow$  leptons decay via effective operators. No signal observed. Limits placed on EFT operators.

<sup>2</sup> BALDIN 76 is a 70 GeV Serpukhov experiment. Value is per Al nucleus at  $\theta = 0$ . For other charges in range  $-0.5$  to  $-3.0$ , CL = 90% limit is  $(2.6 \times 10^{-36})/|(\text{charge})|$  for mass range  $(2.1\text{--}9.4 \text{ GeV}) \times |(\text{charge})|$ . Assumes stable particle interacting with matter as do antiprotons.

<sup>3</sup> ALBROW 75 is a CERN ISR experiment with  $E_{\text{cm}} = 53$  GeV.  $\theta = 40$  mr. See figure 5 for mass ranges up to 35 GeV.

<sup>4</sup> JOVANOVIH 75 is a CERN ISR 26+26 and 15+15 GeV  $pp$  experiment. Figure 4 covers ranges  $Q = 1/3$  to 2 and  $m = 3$  to 26 GeV. Value is per GeV momentum.

<sup>5</sup> APPEL 74 is NAL 300 GeV  $pW$  experiment. Studies forward production of heavy (up to 24 GeV) charged particles with momenta 24–200 GeV ( $-$ charge) and 40–150 GeV ( $+$ charge). Above typical value is for 75 GeV and is per GeV momentum per nucleon.

<sup>6</sup> ALPER 73 is CERN ISR 26+26 GeV  $pp$  experiment.  $p > 0.9$  GeV,  $0.2 < \beta < 0.65$ .

<sup>7</sup> ANTIPOV 71B is from same 70 GeV  $p$  experiment as ANTIPOV 71C and BINON 69.

<sup>8</sup> ANTIPOV 71C limit inferred from flux ratio. 70 GeV  $p$  experiment.

<sup>9</sup> DORFAN 65 is a 30 GeV/ $c$   $p$  experiment at BNL. Units are per GeV momentum per nucleus.

**Long-Lived Heavy Particle Invariant Cross Section**

VALUE ( $\text{cm}^2/\text{GeV}^2/N$ )	CL%	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<5\text{--}700 \times 10^{-35}$	90	<sup>1</sup> BERNSTEIN 88	CNTR		
$<5\text{--}700 \times 10^{-37}$	90	<sup>1</sup> BERNSTEIN 88	CNTR		
$<2.5 \times 10^{-36}$	90	<sup>2</sup> THRON 85	CNTR	–	$Q=1$ , $m=4\text{--}12$ GeV
$<1. \times 10^{-35}$	90	<sup>2</sup> THRON 85	CNTR	+	$Q=1$ , $m=4\text{--}12$ GeV
$<6. \times 10^{-33}$	90	<sup>3</sup> ARMITAGE 79	SPEC		$m=1.87$ GeV
$<1.5 \times 10^{-33}$	90	<sup>3</sup> ARMITAGE 79	SPEC		$m=1.5\text{--}3.0$ GeV
		<sup>4</sup> BOZZOLI 79	CNTR	$\pm$	$Q = (2/3, 1, 4/3, 2)$
$<1.1 \times 10^{-37}$	90	<sup>5</sup> CUTTS 78	CNTR		$m=4\text{--}10$ GeV
$<3.0 \times 10^{-37}$	90	<sup>6</sup> VIDAL 78	CNTR		$m=4.5\text{--}6$ GeV

- <sup>1</sup>BERNSTEIN 88 limits apply at  $x = 0.2$  and  $p_T = 0$ . Mass and lifetime dependence of limits are shown in the regions:  $m = 1.5\text{--}7.5$  GeV and  $\tau = 10^{-8}\text{--}2 \times 10^{-6}$  s. First number is for hadrons; second is for weakly interacting particles.
- <sup>2</sup>THRON 85 is FNAL 400 GeV proton experiment. Mass determined from measured velocity and momentum. Limits are for  $\tau > 3 \times 10^{-9}$  s.
- <sup>3</sup>ARMITAGE 79 is CERN-ISR experiment at  $E_{\text{cm}} = 53$  GeV. Value is for  $x = 0.1$  and  $p_T = 0.15$ . Observed particles at  $m = 1.87$  GeV are found all consistent with being antideuterons.
- <sup>4</sup>BOZZOLI 79 is CERN-SPS 200 GeV  $pN$  experiment. Looks for particle with  $\tau$  larger than  $10^{-8}$  s. See their figure 11–18 for production cross-section upper limits vs mass.
- <sup>5</sup>CUTTS 78 is  $p\text{Be}$  experiment at FNAL sensitive to particles of  $\tau > 5 \times 10^{-8}$  s. Value is for  $-0.3 < x < 0$  and  $p_T = 0.175$ .
- <sup>6</sup>VIDAL 78 is FNAL 400 GeV proton experiment. Value is for  $x = 0$  and  $p_T = 0$ . Puts lifetime limit of  $< 5 \times 10^{-8}$  s on particle in this mass range.

### Long-Lived Heavy Particle Production ( $\sigma(\text{Heavy Particle}) / \sigma(\pi)$ )

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 10^{-8}$		<sup>1</sup> NAKAMURA 89	SPEC	$\pm$	$Q = (-5/3, \pm 2)$
	0	<sup>2</sup> BUSSIÈRE 80	CNTR	$\pm$	$Q = (2/3, 1, 4/3, 2)$
<sup>1</sup> NAKAMURA 89 is KEK experiment with 12 GeV protons on Pt target. The limit applies for mass $\lesssim 1.6$ GeV and lifetime $\gtrsim 10^{-7}$ s.					
<sup>2</sup> BUSSIÈRE 80 is CERN-SPS experiment with 200–240 GeV protons on Be and Al target. See their figures 6 and 7 for cross-section ratio vs mass.					

### Production and Capture of Long-Lived Massive Particles

VALUE ( $10^{-36} \text{ cm}^2$ )	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$< 20$ to 800	<sup>1</sup> AAD	21X	ATLAS search for captured LLPs
	<sup>2</sup> ACHARYA	21	INDU dyons production, capture
	<sup>3</sup> ALEKSEEV	76	ELEC $\tau = 5$ ms to 1 day
$< 200$ to 2000	<sup>3</sup> ALEKSEEV	76B	ELEC $\tau = 100$ ms to 1 day
$< 1.4$ to 9	<sup>4</sup> FRANKEL	75	CNTR $\tau = 50$ ms to 10 hours
$< 0.1$ to 9	<sup>5</sup> FRANKEL	74	CNTR $\tau = 1$ to 1000 hours
<sup>1</sup> AAD 21X search for LLPs which come to rest in ATLAS detector to deposit energy between collisions. No signal observed in $111 \text{ fb}^{-1}$ of data. Limits placed in lifetime vs. mass plane assuming model with gluino hadrons: e.g. $m > 1.4$ TeV for $\tau \sim 10^{-5}$ to $10^3$ sec.			
<sup>2</sup> ACHARYA 21 search for dyons (carrying electric and magnetic charge) and monopoles via production and capture in $6.46 \text{ fb}^{-1}$ of 13 TeV LHC data. No signal observed. Limits placed in mass vs. magnetic charge plane.			
<sup>3</sup> ALEKSEEV 76 and ALEKSEEV 76B are 61–70 GeV $p$ Serpukhov experiment. Cross section is per Pb nucleus.			
<sup>4</sup> FRANKEL 75 is extension of FRANKEL 74.			
<sup>5</sup> FRANKEL 74 looks for particles produced in thick Al targets by 300–400 GeV/ $c$ protons.			

**Long-Lived Particle (LLP) Search at Hadron Collisions**

Limits are for cross section times branching ratio.

VALUE (fb)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<	0.07	95	1 AAD	24AS ATLS long-lived dark photon search
			2 AAD	24BN ATLS hadronic LLP search
			3 AAD	24BZ ATLS LLP + lepton/jet search
			4 HAYRAPETY...24AI	CMS LLP search
			5 HAYRAPETY...24M	CMS long-lived SUSY
			6 HAYRAPETY...24P	CMS LLP from SUSY search
			7 HAYRAPETY...24S	CMS long-lived HNL search
			8 HAYRAPETY...24Y	CMS LLP → dimuon search
			9 AAD	23AM ATLS LLP higgsino search
			10 AAD	23AR ATLS LLP search via displaced $\gamma$
			11 AAD	23BQ ATLS displaced dimuon search
			12 AAD	23CO ATLS highly ionizing LLP/monopole
			13 AAD	23G ATLS heavy highly ionizing LLP search
			14 AAD	23I ATLS light LLP via collimated decays
			15 TUMASYAN	23AO CMS LLP search via trackless jets
			16 TUMASYAN	23G CMS LLP search via displaced dimuons
			17 AAD	22H ATLS LLP search with $\mu$ spectrometer
			18 AAD	22K ATLS LLP search via displaced jets in the calorimeter
			19 AAD	22U ATLS LLP/chargino search via tracklet
			20 AAIJ	22U LHCb LLP semileptonic decay to muon
			21 ACHARYA	22A MOED monopoles/HECOs at LHC
			22 TUMASYAN	22AD CMS heavy neutral lepton LLP search
			23 TUMASYAN	22AF CMS LLP search via displaced lepton tracks
			24 TUMASYAN	22M CMS LLP search via $ZH$ production
			25 TUMASYAN	22N CMS LLP search via dimuons
			26 AAD	21AL ATLS charged LLPs search
			27 AAD	21BA ATLS LLP from higgs decay search
			28 AAIJ	21V LHCb LLP → $e\mu\nu$ search
			29 SIRUNYAN	21AF CMS LLP search via displaced jets
			30 SIRUNYAN	21U CMS LLP search via displaced jets
			31 TUMASYAN	21 CMS LLP endcap muon detector searches
			32 AAD	20D ATLS $pp \rightarrow$ LLPs at 13 TeV
			33 AAD	20J ATLS scalar boson decay to LLPs
			34 AAD	20M ATLS LLP top squark decay to $\mu$
			35 AAD	20P ATLS LLP dark photon search
			36 AAIJ	20AL LHCb $pp$ dimuon resonance
			37 BALL	20 LLP milli-charged particles at LHC
			38 AABOUD	19AE ATLS $pp$ at 13 TeV
			39 AABOUD	19AK ATLS $pp \rightarrow \Phi \rightarrow ZZ_d$
			40 AABOUD	19AM ATLS DY multi-charged LLP production
			41 AABOUD	19AO ATLS LLP via displaced jets
			42 AABOUD	19AT ATLS heavy, charged LLPs
			43 AABOUD	19G ATLS LLP decay to $\mu^+\mu^-$
			44 SIRUNYAN	19BH CMS LLP via displaced jets

		45	SIRUNYAN	19BT CMS	LLP via displaced jets+MET
		46	SIRUNYAN	19CA CMS	LLP $\rightarrow \gamma$ search
		47	SIRUNYAN	19Q CMS	$pp \rightarrow j + \text{displaced dark quark jet}$
		48	SIRUNYAN	18AW CMS	Long-lived particle search
		49	AAIJ	16AR LHCb	$H \rightarrow XX$ LLPs
		50	KHACHATRY...	16BW CMS	direct production: HSCPs
<2000	90	51	BADIER	86 BDMP	$\tau = (0.05-1.) \times 10^{-8} \text{s}$
<sup>1</sup> AAD 24AS search for long-lived dark photons (DP) produced from dark sector. No signal observed. Limits placed in portal coupling vs $m(\text{DP})$ plane.					
<sup>2</sup> AAD 24BN report on ATLAS search for hadronically-decaying long-lived particles that decay in the tracking detector. No signal was observed. Limits placed in $c\tau$ vs cross section or branching fraction plane for various simplified new physics models.					
<sup>3</sup> AAD 24BZ search for long-lived particle decaying to jets in hadronic calorimeter in association with lepton/jet. No signal observed. Limits placed in $c\tau$ vs. cross section plane for various simplified new physics models.					
<sup>4</sup> HAYRAPETYAN 24AI search for long-lived particles (LLPs) using a Muon Detector Shower with a high hit multiplicity in the muon chambers with $138 \text{ fb}^{-1}$ of data at 13 TeV. No signal was observed. Limits placed in Higgs BF vs. $c\tau$ (lifetime) plane.					
<sup>5</sup> HAYRAPETYAN 24M search for long-lived charginos in SUSY decays with $138 \text{ fb}^{-1}$ of data at 13 TeV. No signal observed. Limits placed in mass vs mass difference plane for various simplified models.					
<sup>6</sup> HAYRAPETYAN 24P search for long-lived particles in SUSY models via displaced vertices with $138 \text{ fb}^{-1}$ of data at 13 TeV. No signal observed. Limits set in split SUSY and gauge mediated SUSY breaking models.					
<sup>7</sup> HAYRAPETYAN 24S search for long-lived heavy neutral lepton decaying in muon chambers with $138 \text{ fb}^{-1}$ of data at 13 TeV. No signal observed. Limits placed in mass vs. mixing angle plane.					
<sup>8</sup> HAYRAPETYAN 24Y search for displaced dimuons in CMS with $36 \text{ fb}^{-1}$ of data at 13.6 TeV. No signal observed. Limits placed for various models including Abelian dark Higgs and RPV SUSY.					
<sup>9</sup> AAD 23AM search for long-lived higgsinos from gauge-mediation which decay to $Z$ or $H$ . No signal observed. Limits placed in $c\tau$ vs $m(\chi)$ plane for various simplified models.					
<sup>10</sup> AAD 23AR search for long-lived particles via decay to displaced $\gamma$ with $139 \text{ fb}^{-1}$ of data. No signal observed. Limits placed in $m$ vs. $\tau$ and BF vs. $\tau$ planes for gauge-mediated SUSY model.					
<sup>11</sup> AAD 23BQ search for displaced dimuon events in ATLAS detector. No signal observed. Limits placed in smuon lifetime vs. mass plane for long-lived smuon model.					
<sup>12</sup> AAD 23CO search for monopoles and high-electric-charge LLPs in ATLAS with $139 \text{ fb}^{-1}$ of data. No signal observed. Limits placed in mass vs. charge plane.					
<sup>13</sup> AAD 23G search for heavy highly ionizing long-lived particles with $139 \text{ fb}^{-1}$ of data. No signal observed. Limits placed in $m$ vs. $\tau$ plane for several SUSY models.					
<sup>14</sup> AAD 23I search for light long-lived particles decaying to collimated decay products (e.g. dileptons). No signal observed. Limits placed in BF vs. $\tau$ plane.					
<sup>15</sup> TUMASYAN 23AO search for trackless jets from LLP production at CMS. No signal observed. Limits placed for SUSY model with long-lived neutralino in $m(\chi)$ vs. $c\tau$ plane.					
<sup>16</sup> TUMASYAN 23G search for LLP decaying to displaced dimuons at CM with $97.6 \text{ fb}^{-1}$ of data. No signal observed. Limits placed in $c\tau$ vs. $m$ plane for hidden Abelian Higgs simplified model.					
<sup>17</sup> AAD 22H search for scalar mediator decay to two LLPs which decay in muon chambers with $139 \text{ fb}^{-1}$ at 13 TeV; no signal detected; limits placed on various simplified models.					
<sup>18</sup> AAD 22K search for LLP pair production via scalar mediator with LLP decay in hadron calorimeter; no signal detected; limits placed for various simplified models.					

- 19 AAD 22U search for chargino LLP via disappearing tracks; no signal observed; limits placed in  $m(\text{chargino})$  vs lifetime plane for cases of higgsino- or wino-like chargino.
- 20 AAIJ 22U reports search for LLP production at LHCb with  $5.4 \text{ fb}^{-1}$  at 13 TeV followed by semileptonic decay to muon; no signal detected; limits placed in mass or lifetime vs. cross section plane for several simplified models.
- 21 ACHARYA 22A report search for monople and HECO production via DY at 8 TeV LHC with  $2.2 \text{ fb}^{-1}$  with MoEDAL detector; no signal detected; limits placed in mass vs. cross section plane for various electric/magnetic charge scenarios.
- 22 TUMASYAN 22AD search for heavy neutral lepton which decays as LLP to trilepton state with  $138 \text{ fb}^{-1}$  at 13 TeV; no signal detected; limits placed in mass vs. coupling plane.
- 23 TUMASYAN 22AF search for LLPs via displaced lepton vertices. The analysis is performed with an integrated luminosity of 118 (113)  $\text{fb}^{-1}$  when analyzing the  $ee$  ( $e\mu$ ,  $\mu\mu$ ) channel; no signal detected; limits placed for a variety of simplified models.
- 24 TUMASYAN 22M search in  $117 \text{ fb}^{-1}$  of 13 TeV data for  $ZH$  production with  $H \rightarrow SS$  where  $S$  is a LLP; no signal observed; limits placed in decay length vs. branching fraction plane.
- 25 TUMASYAN 22N search in  $101 \text{ fb}^{-1}$  of 13 TeV data for LLP production via decay to dimuons; no signal observed; limits placed on mass vs. coupling or lifetime for a variety of simplified models.
- 26 AAD 21AL reports on ATLAS search for long-lived charged particles with  $139 \text{ fb}^{-1}$  at 13 TeV. No signal observed. Limits placed in lifetime vs. mass plane: e.g. for  $\tau(\text{LLP}) \sim 0.1 \text{ ns}$ ,  $m(\text{selectron}) > 720 \text{ GeV}$ .
- 27 AAD 21BA search for long-lived particles from  $ZH$  production ( $H \rightarrow b\bar{b}$ ) with 2 displaced vertices in  $139 \text{ fb}^{-1}$  of data at 13 TeV. No signal detected. Limits placed in branching fraction vs. lifetime plane.
- 28 AAIJ 21V search for  $pp \rightarrow \text{LLP} + \text{LLP}$  with  $\text{LLP} \rightarrow e\mu\nu$  in the lifetime range between 2 and 50 ps at LHCb with  $5.4 \text{ fb}^{-1}$  at 13 TeV. No signal observed. Limits placed in LLP cross section vs. mass or lifetime plane for  $m(\text{LLP}) \sim 7$  to 50 GeV.
- 29 SIRUNYAN 21AF search for LLPs at CMS via jets with 2 displaced vertices in  $140 \text{ fb}^{-1}$  of data at 13 TeV. No signal observed. Limits placed for RPV SUSY models in which a long-lived neutralino or gluino decays into a multijet final state with top, bottom, and strange quarks.
- 30 SIRUNYAN 21U search for long-lived particles (LLPs) via displaced jets at CMS with LHC13 and  $132 \text{ fb}^{-1}$ . No signal detected. Limits placed on simplified model production of  $\text{LLP } X \rightarrow q\bar{q}$  with  $\sigma < 0.07 \text{ fb}$  for  $m(X) > 500 \text{ GeV}$  and  $c\tau \sim 2$  to 250 mm.
- 31 TUMASYAN 21 search for long-lived particles in CMS muon endcap detector in  $137 \text{ fb}^{-1}$  of data at 13 TeV. No signal detected. Limits are placed depending on the branching fraction of Higgs boson to LLP decaying to  $dd$ ,  $bb$ , and  $\tau^+\tau^-$ , depending on proper decay length, and LLP masses.
- 32 AAD 20D search for opposite-sign dileptons originating from long-lived particles in  $pp$  collisions at 13 TeV with  $32.8 \text{ fb}^{-1}$ ; limits placed in squark cross section vs.  $c\tau$  plane for RPV SUSY.
- 33 AAD 20J search for scalar boson decay to two long-lived particles; no signal; limits placed in BF vs  $c\tau$  plane for various mass hypotheses. This search is also combined with other ATLAS displaced-jet searches.
- 34 AAD 20M search for long-lived top-squarks decay to  $\mu$  and hadrons; no signal; limits placed in cross section vs. mass and mass vs. lifetime planes.
- 35 AAD 20P search for long-lived dark photons produced from the decay of a scalar boson, with each dark photon decaying into displaced collimated leptons or light hadrons at 13 TeV with  $36 \text{ fb}^{-1}$ ; no signal; limits placed in  $\sigma \cdot \text{BF}$  vs.  $c\tau$  and other planes.
- 36 AAIJ 20AL search for long-lived  $X \rightarrow \mu^+\mu^-$  decays in  $5.1 \text{ fb}^{-1}$  of LHCb data at 13 TeV; no signal; limits placed on  $m(X)$  up to 3 GeV depending on kinetic mixing.
- 37 BALL 20 search for long-lived milli-charged particles produced at LHC; limits placed in charge vs. mass plane (Fig. 8).

- <sup>38</sup> AABOUD 19AE search for long-lived particles via displaced jets using  $10.8 \text{ fb}^{-1}$  or  $33.0 \text{ fb}^{-1}$  data (depending on a trigger) at 13 TeV; no signal found and limits set in branching ratio vs. decay length plane.
- <sup>39</sup> AABOUD 19AK searches for long-lived particle  $Z_d$  via  $pp \rightarrow \Phi \rightarrow ZZ_d$  at 13 TeV with  $36.1 \text{ fb}^{-1}$ ; no signal found and limits set in  $\sigma \times \text{BR}$  vs. lifetime plane for simplified model.
- <sup>40</sup> AABOUD 19AM search for Drell-Yan (DY) production of long-lived multi-charge particles at 13 TeV with  $36.1 \text{ fb}^{-1}$  of data; no signal found and exclude  $50 \text{ GeV} < m(\text{LLMCP}) < 980\text{--}1220 \text{ GeV}$  for electric charge  $|q| = (2\text{--}7)e$ .
- <sup>41</sup> AABOUD 19AO search for neutral long-lived particles producing displaced jets at 13 TeV with  $36.1 \text{ fb}^{-1}$  of data; no signal found and exclude regions of  $\sigma \cdot \text{BR}$  vs. lifetime plane for various models.
- <sup>42</sup> AABOUD 19AT search for heavy, charged long-lived particles at 13 TeV with  $36.1 \text{ fb}^{-1}$ ; no signal found and upper limits set on masses of various hypothetical particles.
- <sup>43</sup> AABOUD 19G search for long-lived particle with decay to  $\mu^+ \mu^-$  at 13 TeV with  $32.9 \text{ fb}^{-1}$ ; no signal found and limits set in combinations of lifetime, mass and coupling planes for various simplified models.
- <sup>44</sup> SIRUNYAN 19BH search for long-lived SUSY particles via displaced jets at 13 TeV with  $35.9 \text{ fb}^{-1}$ ; no signal found and limits placed in mass vs lifetime plane for various hypothetical models.
- <sup>45</sup> SIRUNYAN 19BT search for displaced jet(s)+ $\cancel{E}_T$  at 13 TeV with  $137 \text{ fb}^{-1}$ ; no signal found and limits placed in mass vs lifetime plane for gauge mediated SUSY breaking models.
- <sup>46</sup> SIRUNYAN 19CA search for gluino/squark decay to long-lived neutralino, decay to  $\gamma$  in GMSB; no signal, limits placed in  $m(\chi)$  vs. lifetime plane for SPS8 GMSB benchmark point.
- <sup>47</sup> SIRUNYAN 19Q search for  $pp \rightarrow j + \text{displaced jet}$  via dark quark with 13 TeV at  $16.1 \text{ fb}^{-1}$ ; no signal found and limits set in mass vs lifetime plane for dark quark/dark pion model.
- <sup>48</sup> SIRUNYAN 18AW search for very long lived particles (LLPs) decaying hadronically or to  $\mu \bar{\mu}$  in CMS detector; none seen/limits set on lifetime vs. cross section.
- <sup>49</sup> AAIJ 16AR search for long lived particles from  $H \rightarrow XX$  with displaced X decay vertex using  $0.62 \text{ fb}^{-1}$  at 7 TeV; limits set in Fig. 7.
- <sup>50</sup> KHACHATRYAN 16BW search for heavy stable charged particles via ToF with  $2.5 \text{ fb}^{-1}$  at 13 TeV; require stable  $m(\text{gluinoball}) > 1610 \text{ GeV}$ .
- <sup>51</sup> BADIÉ 86 looked for long-lived particles at 300 GeV  $\pi^-$  beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass  $> 2 \text{ GeV}$ . The limit applies for particle modes,  $\mu^+ \pi^-$ ,  $\mu^+ \mu^-$ ,  $\pi^+ \pi^- X$ ,  $\pi^+ \pi^- \pi^\pm$  etc. See their figure 5 for the contours of limits in the mass- $\tau$  plane for each mode.

## Long-Lived Heavy Particle Cross Section

VALUE (pb/sr)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<34	95	<sup>1</sup> RAM	94	SPEC 1015< $m_{X^{++}}$ <1085 MeV
<75	95	<sup>1</sup> RAM	94	SPEC 920< $m_{X^{++}}$ <1025 MeV
<sup>1</sup> RAM 94 search for a long-lived doubly-charged fermion $X^{++}$ with mass between $m_N$ and $m_N + m_\pi$ and baryon number +1 in the reaction $pp \rightarrow X^{++} n$ . No candidate is found. The limit is for the cross section at $15^\circ$ scattering angle at 460 MeV incident energy and applies for $\tau(X^{++}) \gg 0.1 \mu\text{s}$ .				

## LIMITS ON CHARGED PARTICLES IN COSMIC RAYS

## Heavy Particle Flux in Cosmic Rays

VALUE ( $\text{cm}^{-2}\text{sr}^{-1}\text{s}^{-1}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6.2	$\times 10^{-10}$	90	0	<sup>1</sup> ALEMANN0 22	DAMP fractionally charged particles in space
				<sup>2</sup> CAO 22	superheavy DM $\rightarrow \gamma$ rays
				<sup>3</sup> ALVIS 18	MAJD Fractionally charged
< 1	$\times 10^{-8}$	90		<sup>4</sup> AGNESE 15	CDM2 $Q = 1/6$
$\sim 6$	$\times 10^{-9}$	2	<sup>5</sup> SAITO 90		$Q \simeq 14, m \simeq 370m_p$
< 1.4	$\times 10^{-12}$	90	0	<sup>6</sup> MINCER 85	CALO $m \geq 1 \text{ TeV}$
				<sup>7</sup> SAKUYAMA 83B	PLAS $m \sim 1 \text{ TeV}$
< 1.7	$\times 10^{-11}$	99	0	<sup>8</sup> BHAT 82	CC
< 1.	$\times 10^{-9}$	90	0	<sup>9</sup> MARINI 82	CNTR $Q = 1, m \sim 4.5m_p$
2.	$\times 10^{-9}$	3	<sup>10</sup> YOCK 81	SPRK	$Q = 1, m \sim 4.5m_p$
		3	<sup>10</sup> YOCK 81	SPRK	Fractionally charged
3.0	$\times 10^{-9}$	3	<sup>11</sup> YOCK 80	SPRK	$m \sim 4.5 m_p$
$(4 \pm 1) \times 10^{-11}$		3	GOODMAN 79	ELEC	$m \geq 5 \text{ GeV}$
< 1.3	$\times 10^{-9}$	90	<sup>12</sup> BHAT 78	CNTR	$m > 1 \text{ GeV}$
< 1.0	$\times 10^{-9}$	0	BRIATORE 76	ELEC	
< 7.	$\times 10^{-10}$	90	0	YOCK 75	ELEC $Q > 7e \text{ or } < -7e$
> 6.	$\times 10^{-9}$	5	<sup>13</sup> YOCK 74	CNTR	$m > 6 \text{ GeV}$
< 3.0	$\times 10^{-8}$	0	DARDO 72	CNTR	
< 1.5	$\times 10^{-9}$	0	TONWAR 72	CNTR	$m > 10 \text{ GeV}$
< 3.0	$\times 10^{-10}$	0	BJORNBOE 68	CNTR	$m > 5 \text{ GeV}$
< 5.0	$\times 10^{-11}$	90	0	JONES 67	ELEC $m = 5\text{--}15 \text{ GeV}$

<sup>1</sup> ALEMANN0 22 search for flux of fractionally charged particles (FCPs) in space; no signal observed; limits set in flux vs charge plane for mass as low as GeV.

<sup>2</sup> CAO 22 search for superheavy DM decaying to gamma rays; no signal observed; limits placed in mass vs. lifetime plane for  $m \sim 10^5\text{--}10^9 \text{ GeV}$  for DM decays to  $b\bar{b}$  or  $\tau\bar{\tau}$ .

<sup>3</sup> ALVIS 18 search for fractional charged flux of cosmic matter at Majorana demonstrator; no signal observed and limits are set on the flux of lightly ionizing particles for charge as low as  $e/1000$ .

<sup>4</sup> See AGNESE 15 Fig. 6 for limits extending down to  $Q = 1/200$ .

<sup>5</sup> SAITO 90 candidates carry about 450 MeV/nucleon. Cannot be accounted for by conventional backgrounds. Consistent with strange quark matter hypothesis.

<sup>6</sup> MINCER 85 is high statistics study of calorimeter signals delayed by 20–200 ns. Calibration with AGS beam shows they can be accounted for by rare fluctuations in signals from low-energy hadrons in the shower. Claim that previous delayed signals including BJORNBOE 68, DARDO 72, BHAT 82, SAKUYAMA 83B below may be due to this fake effect.

<sup>7</sup> SAKUYAMA 83B analyzed 6000 extended air shower events. Increase of delayed particles and change of lateral distribution above  $10^{17} \text{ eV}$  may indicate production of very heavy parent at top of atmosphere.

<sup>8</sup> BHAT 82 observed 12 events with delay  $> 2. \times 10^{-8} \text{ s}$  and with more than 40 particles. 1 eV has good hadron shower. However all events are delayed in only one of two detectors in cloud chamber, and could not be due to strongly interacting massive particle.

- <sup>9</sup> MARINI 82 applied PEP-counter for TOF. Above limit is for velocity = 0.54 of light. Limit is inconsistent with YOCK 80 YOCK 81 events if isotropic dependence on zenith angle is assumed.
- <sup>10</sup> YOCK 81 saw another 3 events with  $Q = \pm 1$  and  $m$  about  $4.5m_p$  as well as 2 events with  $m > 5.3m_p$ ,  $Q = \pm 0.75 \pm 0.05$  and  $m > 2.8m_p$ ,  $Q = \pm 0.70 \pm 0.05$  and 1 event with  $m = (9.3 \pm 3.)m_p$ ,  $Q = \pm 0.89 \pm 0.06$  as possible heavy candidates.
- <sup>11</sup> YOCK 80 events are with charge exactly or approximately equal to unity.
- <sup>12</sup> BHAT 78 is at Kolar gold fields. Limit is for  $\tau > 10^{-6}$  s.
- <sup>13</sup> YOCK 74 events could be tritons.

## Superheavy Particle (Quark Matter) Flux in Cosmic Rays

VALUE ( $\text{cm}^{-2}\text{sr}^{-1}\text{s}^{-1}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		<sup>1</sup> ADRIANI	15	PMLA $4 < m < 1.2 \times 10^5 m_p$
$< 5 \times 10^{-16}$	90	<sup>2</sup> AMBROSIO	00B	MCRO $m > 5 \times 10^{14}$ GeV
$< 1.8 \times 10^{-12}$	90	<sup>3</sup> ASTONE	93	CNTR $m \geq 1.5 \times 10^{-13}$ gram
$< 1.1 \times 10^{-14}$	90	<sup>4</sup> AHLEN	92	MCRO $10^{-10} < m < 0.1$ gram
$< 2.2 \times 10^{-14}$	90	<sup>5</sup> NAKAMURA	91	PLAS $m > 10^{11}$ GeV
$< 6.4 \times 10^{-16}$	90	<sup>6</sup> ORITO	91	PLAS $m > 10^{12}$ GeV
$< 2.0 \times 10^{-11}$	90	<sup>7</sup> LIU	88	BOLO $m > 1.5 \times 10^{-13}$ gram
$< 4.7 \times 10^{-12}$	90	<sup>8</sup> BARISH	87	CNTR $1.4 \times 10^8 < m < 10^{12}$ GeV
$< 3.2 \times 10^{-11}$	90	<sup>9</sup> NAKAMURA	85	CNTR $m > 1.5 \times 10^{-13}$ gram
$< 3.5 \times 10^{-11}$	90	<sup>10</sup> ULLMAN	81	CNTR Planck-mass $10^{19}$ GeV
$< 7. \times 10^{-11}$	90	<sup>10</sup> ULLMAN	81	CNTR $m \leq 10^{16}$ GeV

- <sup>1</sup> ADRIANI 15 search for relatively light quark matter with charge  $Z = 1-8$ . See their Figs. 2 and 3 for flux upper limits.
- <sup>2</sup> AMBROSIO 00B searched for quark matter ("nuclearites") in the velocity range  $(10^{-5}-1) c$ . The listed limit is for  $2 \times 10^{-3} c$ .
- <sup>3</sup> ASTONE 93 searched for quark matter ("nuclearites") in the velocity range  $(10^{-3}-1) c$ . Their Table 1 gives a compilation of searches for nuclearites.
- <sup>4</sup> AHLEN 92 searched for quark matter ("nuclearites"). The bound applies to velocity  $< 2.5 \times 10^{-3} c$ . See their Fig. 3 for other velocity/ $c$  and heavier mass range.
- <sup>5</sup> NAKAMURA 91 searched for quark matter in the velocity range  $(4 \times 10^{-5}-1) c$ .
- <sup>6</sup> ORITO 91 searched for quark matter. The limit is for the velocity range  $(10^{-4}-10^{-3}) c$ .
- <sup>7</sup> LIU 88 searched for quark matter ("nuclearites") in the velocity range  $(2.5 \times 10^{-3}-1) c$ . A less stringent limit of  $5.8 \times 10^{-11}$  applies for  $(1-2.5) \times 10^{-3} c$ .
- <sup>8</sup> BARISH 87 searched for quark matter ("nuclearites") in the velocity range  $(2.7 \times 10^{-4}-5 \times 10^{-3}) c$ .
- <sup>9</sup> NAKAMURA 85 at KEK searched for quark-matter. These might be lumps of strange quark matter with roughly equal numbers of  $u$ ,  $d$ ,  $s$  quarks. These lumps or nuclearites were assumed to have velocity of  $(10^{-4}-10^{-3}) c$ .
- <sup>10</sup> ULLMAN 81 is sensitive for heavy slow singly charge particle reaching earth with vertical velocity 100–350 km/s.

**Highly Ionizing Particle Flux**

<u>VALUE</u> ( $\text{m}^{-2}\text{yr}^{-1}$ )	<u>CL%</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.4	95	0	KINOSHITA	81B PLAS	$Z/\beta$ 30–100

**SEARCHES FOR BLACK HOLE PRODUCTION**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	<sup>1</sup> AABOUD 16P	ATLS	13 TeV $pp \rightarrow e\mu, e\tau, \mu\tau$
	<sup>2</sup> AAD 15AN	ATLS	8 TeV $pp \rightarrow$ multijets
	<sup>3</sup> AAD 14A	ATLS	8 TeV $pp \rightarrow \gamma + \text{jet}$
	<sup>4</sup> AAD 14AL	ATLS	8 TeV $pp \rightarrow \ell + \text{jet}$
	<sup>5</sup> AAD 14C	ATLS	8 TeV $pp \rightarrow \ell + (\ell \text{ or jets})$
	<sup>6</sup> AAD 13D	ATLS	7 TeV $pp \rightarrow 2 \text{ jets}$
	<sup>7</sup> CHATRCHYAN 13A	CMS	7 TeV $pp \rightarrow 2 \text{ jets}$
	<sup>8</sup> CHATRCHYAN 13AD	CMS	8 TeV $pp \rightarrow$ multijets
	<sup>9</sup> AAD 12AK	ATLS	7 TeV $pp \rightarrow \ell + (\ell \text{ or jets})$
	<sup>10</sup> CHATRCHYAN 12W	CMS	7 TeV $pp \rightarrow$ multijets
	<sup>11</sup> AAD 11AG	ATLS	7 TeV $pp \rightarrow 2 \text{ jets}$

<sup>1</sup> AABOUD 16P set limits on quantum BH production in  $n = 6$  ADD or  $n = 1$  RS models.

<sup>2</sup> AAD 15AN search for black hole or string ball formation followed by its decay to multijet final states, in  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $L = 20.3 \text{ fb}^{-1}$ . See their Figs. 6–8 for limits.

<sup>3</sup> AAD 14A search for quantum black hole formation followed by its decay to a  $\gamma$  and a jet, in  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $L = 20 \text{ fb}^{-1}$ . See their Fig. 3 for limits.

<sup>4</sup> AAD 14AL search for quantum black hole formation followed by its decay to a lepton and a jet, in  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $L = 20.3 \text{ fb}^{-1}$ . See their Fig. 2 for limits.

<sup>5</sup> AAD 14C search for microscopic (semiclassical) black hole formation followed by its decay to final states with a lepton and  $\geq 2$  (leptons or jets), in  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $L = 20.3 \text{ fb}^{-1}$ . See their Figures 8–11, Tables 7, 8 for limits.

<sup>6</sup> AAD 13D search for quantum black hole formation followed by its decay to two jets, in  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  with  $L = 4.8 \text{ fb}^{-1}$ . See their Fig. 8 and Table 3 for limits.

<sup>7</sup> CHATRCHYAN 13A search for quantum black hole formation followed by its decay to two jets, in  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  with  $L = 5 \text{ fb}^{-1}$ . See their Figs. 5 and 6 for limits.

<sup>8</sup> CHATRCHYAN 13AD search for microscopic (semiclassical) black hole formation followed by its evaporation to multiparticle final states, in multijet (including  $\gamma, \ell$ ) events in  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $L = 12 \text{ fb}^{-1}$ . See their Figs. 5–7 for limits.

<sup>9</sup> AAD 12AK search for microscopic (semiclassical) black hole formation followed by its decay to final states with a lepton and  $\geq 2$  (leptons or jets), in  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  with  $L = 1.04 \text{ fb}^{-1}$ . See their Fig. 4 and 5 for limits.

<sup>10</sup> CHATRCHYAN 12W search for microscopic (semiclassical) black hole formation followed by its evaporation to multiparticle final states, in multijet (including  $\gamma, \ell$ ) events in  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  with  $L = 4.7 \text{ fb}^{-1}$ . See their Figs. 5–8 for limits.

<sup>11</sup> AAD 11AG search for quantum black hole formation followed by its decay to two jets, in  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  with  $L = 36 \text{ pb}^{-1}$ . See their Fig. 11 and Table 4 for limits.

## REFERENCES FOR Other Particle Searches

AAD	24AE	JHEP 2404 118	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24AK	JHEP 2405 263	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24AS	EPJ C84 719	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24AT	EPJ C84 742	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24B	PL B848 138324	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24BN	PRL 133 161803	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24BV	PL B858 139007	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24BZ	JHEP 2411 036	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24CD	JHEP 2411 126	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24CM	JHEP 2409 005	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24E	PRL 132 081801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	24S	PR D109 032010	G. Aad <i>et al.</i>	(ATLAS Collab.)
ANDREEV	24	PRL 132 211803	Yu.M. Andreev <i>et al.</i>	(NA64 Collab.)
BARAK	24	PRL 133 071801	L. Barak <i>et al.</i>	(SENSEI Collab., +)
HAYRAPETY...	24AB	JHEP 2406 123	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24AC	JHEP 2406 183	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24AD	JHEP 2407 142	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24AI	PR D110 032007	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24AO	PRL 133 191902	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24AZ	JHEP 2412 172	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24G	PRL 133 011801	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24M	PR D109 072007	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24P	PR D109 112005	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24S	PR D110 012004	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
HAYRAPETY...	24Y	JHEP 2405 047	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
AAD	23AM	PR D108 012012	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23AR	PR D108 032016	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23BQ	PL B846 138172	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23BT	PL B847 138316	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23CO	JHEP 2311 112	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23G	JHEP 2306 158	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23I	JHEP 2306 153	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23P	JHEP 2307 176	G. Aad <i>et al.</i>	(ATLAS Collab.)
ADACHI	23K	PR D108 L111104	I. Adachi <i>et al.</i>	(BELLE II Collab.)
HAYRAPETY...	23F	JHEP 2312 068	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	23AO	JHEP 2307 210	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	23BC	JHEP 2312 189	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	23BF	EPJ C83 827	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	23G	JHEP 2305 228	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	23L	JHEP 2307 161	A. Tumasyan <i>et al.</i>	(CMS Collab.)
AAD	22G	PR D105 092012	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	22H	PR D106 032005	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	22K	JHEP 2206 005	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	22U	EPJ C82 606	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	22U	EPJ C82 373	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABRATENKO	22A	PR D106 092006	P. Abratenko <i>et al.</i>	(MicroBooNE Collab.)
ACHARYA	22A	EPJ C82 694	B. Acharya <i>et al.</i>	(MoEDAL Collab.)
ALEMANNO	22	PR D106 063026	F. Alemanno <i>et al.</i>	(DAMPE Collab.)
ANDREEV	22A	PRL 129 161801	Yu.M. Andreev <i>et al.</i>	(NA64 Collab.)
CAO	22	PRL 129 261103	Z. Cao <i>et al.</i>	(LHAASO Collab.)
TUMASYAN	22AD	JHEP 2207 081	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	22AF	EPJ C82 153	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	22AG	EPJ C82 213	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	22H	PR D105 112007	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	22M	JHEP 2203 160	A. Tumasyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	22N	JHEP 2204 062	A. Tumasyan <i>et al.</i>	(CMS Collab.)
AAD	21AL	PRL 127 051802	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	21BA	JHEP 2111 229	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	21F	PR D103 112006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	21X	JHEP 2107 173	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	21V	EPJ C81 261	R. Aaij <i>et al.</i>	(LHCb Collab.)
ACHARYA	21	PRL 126 071801	B. Acharya <i>et al.</i>	(MoEDAL Collab.)
AFEK	21	PR D104 012004	G. Afek <i>et al.</i>	(YALE)
ALKHATIB	21A	PRL 127 081802	I. Alkhatib <i>et al.</i>	(SuperCDMS Collab.)
ANDREEV	21	PRL 126 211802	Yu.M. Andreev <i>et al.</i>	(NA64 Collab.)
SIRUNYAN	21AF	PR D104 052011	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	21T	EPJ C81 629	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	21U	PR D104 012015	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
TUMASYAN	21	PRL 127 261804	A. Tumasyan <i>et al.</i>	(CMS Collab.)

AAD	20AD	PRL 125 131801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20D	PL B801 135114	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20J	PR D101 052013	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20M	PR D102 032006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20P	EPJ C80 450	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20T	JHEP 2003 145	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	20W	JHEP 2006 151	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	20AL	JHEP 2010 156	R. Aaij <i>et al.</i>	(LHCb Collab.)
AGUILAR-AR...	20B	JHEP 2004 054	A. Aguilar-Arevalo <i>et al.</i>	(CONNIE Collab.)
BALL	20	PR D102 032002	A.H. Ball <i>et al.</i>	(milliQan)
FEDDERKE	20	PR D101 115021	M.A. Fedderke, P.W. Graham, S. Rajendran	(STAN+)
SIRUNYAN	20A	EPJ C80 3	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	20AI	JHEP 2005 033	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	20AY	PL B808 135578	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	20C	EPJ C80 75	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	20N	PL B806 135502	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	20Z	JHEP 2003 051	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	19AA	EPJ C79 120	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AE	EPJ C79 481	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AJ	PL B795 56	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AK	PRL 122 151801	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AM	PR D99 052003	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AO	PR D99 052005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19AT	PR D99 092007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19G	PR D99 012001	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19H	PR D99 012008	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19Q	JHEP 1905 041	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19V	JHEP 1905 142	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
ALCANTARA	19	PR D99 103016	E. Alcantara, L.A. Anchordoqui, J.F. Soriano	
SIRUNYAN	19B	PR D99 012005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19BH	PR D99 032011	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19BT	PL B797 134876	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19CA	PR D100 112003	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19CD	PRL 123 231803	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19O	JHEP 1902 074	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19Q	JHEP 1902 179	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	18AD	PL B779 24	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18CJ	PR D98 052008	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18CK	PR D98 092002	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18CL	PR D98 092005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18CM	PR D98 092008	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18N	PRL 121 081801	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ	18AJ	PRL 120 061801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ALBERT	18C	PR D98 123012	A. Albert <i>et al.</i>	(HAWC Collab.)
ALVIS	18	PRL 120 211804	S.I. Alvis <i>et al.</i>	(MAJORANA Collab.)
BANERJEE	18	PRL 120 231802	D. Banerjee <i>et al.</i>	(NA64 Collab.)
BANERJEE	18A	PR D97 072002	D. Banerjee <i>et al.</i>	(NA64 Collab.)
KILE	18	JHEP 1810 116	J. Kile, J. von Wimmersperg-Toeller	(LISBT)
MARSICANO	18	PR D98 015031	L. Marsicano <i>et al.</i>	
PORAYKO	18	PR D98 102002	N.K. Porayako <i>et al.</i>	(PPTA Collab.)
SIRUNYAN	18AW	JHEP 1805 127	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18BB	JHEP 1806 120	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DA	JHEP 1811 042	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DD	EPJ C78 789	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DJ	JHEP 1809 101	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DR	JHEP 1811 161	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DU	PR D98 092001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DY	PR D98 112014	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18ED	JHEP 1811 172	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	17B	PL B765 32	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17D	PR D95 032001	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17L	JHEP 1708 052	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ	17BR	EPJ C77 812	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AA	PL B774 252	M. Ablikim <i>et al.</i>	(BESIII Collab.)
KHACHATRY...	17D	JHEP 1701 076	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17W	PL B769 520	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	17Y	PL B770 257	V. Khachatryan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17B	JHEP 1704 136	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17C	JHEP 1705 029	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17F	JHEP 1707 013	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17J	JHEP 1708 073	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)

ZANG	17	PL B773 159	X. Zang, G.A. Miller	(WASH)
AABOUD	16	PL B759 229	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16P	EPJ C76 541	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	16AI	JHEP 1603 041	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16N	JHEP 1603 026	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16O	PL B760 520	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16R	PL B755 285	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16S	PL B754 302	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	16AR	EPJ C76 664	R. Aaij <i>et al.</i>	(LHCb Collab.)
KHACHATRY...	16BW	PR D94 112004	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16K	PRL 116 071801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16L	PRL 117 031802	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16M	PRL 117 051802	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KRASZNAHO...	16	PRL 116 042501	A.J. Krasznahorkay <i>et al.</i>	(HINR, ANIK+)
AAD	15AN	JHEP 1507 032	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AT	EPJ C75 79	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15BJ	EPJ C75 362	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	15BD	EPJ C75 595	R. Aaij <i>et al.</i>	(LHCb Collab.)
ADRIANI	15	PRL 115 111101	O. Adriani <i>et al.</i>	(PAMELA Collab.)
AGNESE	15	PRL 114 111302	R. Agnese <i>et al.</i>	(CDMS Collab.)
KHACHATRY...	15F	PRL 114 101801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
LEES	15E	PRL 114 171801	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAD	14A	PL B728 562	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14AL	PRL 112 091804	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14C	JHEP 1408 103	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	14J	PR D89 092001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAD	13A	PL B718 860	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13AH	PL B722 305	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13C	PRL 110 011802	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13D	JHEP 1301 029	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	13I	PR D88 031103	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	13R	PRL 111 031802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
CHATRCHYAN	13	PL B718 815	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13A	JHEP 1301 013	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AB	JHEP 1307 122	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
Also		JHEP 2211 149 (errat.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AD	JHEP 1307 178	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13AR	PR D87 092008	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
Also		PR D106 099903 (errat.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAD	12AK	PL B716 122	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12C	PRL 108 041805	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12S	PL B708 37	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	12M	PRL 108 211804	T. Aaltonen <i>et al.</i>	(CDF Collab.)
CHATRCHYAN	12AP	JHEP 1209 094	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12BL	JHEP 1212 015	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12Q	PL B716 260	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12T	PRL 108 261803	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12W	JHEP 1204 061	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAD	11AG	NJP 13 053044	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11I	PL B698 353	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11S	PL B705 294	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	11AF	PRL 107 181801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11M	PRL 106 171801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	11I	PRL 107 011804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	11C	JHEP 1106 026	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	11U	PRL 107 201804	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAD	10	PRL 105 161801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	10AF	PR D82 052005	T. Aaltonen <i>et al.</i>	(CDF Collab.)
KHACHATRY...	10	PRL 105 211801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
Also		PRL 106 029902	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AALTONEN	09AF	PR D80 011102	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09G	PR D79 052004	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09Z	PRL 103 021802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	09M	PRL 102 161802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AKTAS	04C	EPJ C36 413	A. Atkas <i>et al.</i>	(H1 Collab.)
JAVORSEK	02	PR D65 072003	D. Javorsek II <i>et al.</i>	
JAVORSEK	01	PR D64 012005	D. Javorsek II <i>et al.</i>	
JAVORSEK	01B	PRL 87 231804	D. Javorsek II <i>et al.</i>	
ABBIENDI	00D	EPJ C13 197	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AMBROSIO	00B	EPJ C13 453	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
ABE	99F	PRL 82 2038	F. Abe <i>et al.</i>	(CDF Collab.)

ACKERSTAFF	98P	PL B433 195	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ABE	97G	PR D55 5263	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	97D	PL B396 315	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97B	PL B391 210	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ADAMS	97B	PRL 79 4083	J. Adams <i>et al.</i>	(FNAL KTeV Collab.)
BARATE	97K	PL B405 379	R. Barate <i>et al.</i>	(ALEPH Collab.)
AKERS	95R	ZPHY C67 203	R. Akers <i>et al.</i>	(OPAL Collab.)
GALLAS	95	PR D52 6	E. Gallas <i>et al.</i>	(MSU, FNAL, MIT, FLOR)
RAM	94	PR D49 3120	S. Ram <i>et al.</i>	(TELA, TRIU)
ABE	93G	PRL 71 2542	F. Abe <i>et al.</i>	(CDF Collab.)
ASTONE	93	PR D47 4770	P. Astone <i>et al.</i>	(ROMA, ROMAI, CATA, FRAS)
BUSKULIC	93C	PL B303 198	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
YAMAGATA	93	PR D47 1231	T. Yamagata, Y. Takamori, H. Utsunomiya	(KONAN)
ABE	92J	PR D46 1889	F. Abe <i>et al.</i>	(CDF Collab.)
AHLEN	92	PRL 69 1860	S.P. Ahlen <i>et al.</i>	(MACRO Collab.)
VERKERK	92	PRL 68 1116	P. Verkerk <i>et al.</i>	(ENSP, SACL, PAST)
AKESSON	91	ZPHY C52 219	T. Akesson <i>et al.</i>	(HELIOS Collab.)
NAKAMURA	91	PL B263 529	S. Nakamura <i>et al.</i>	
ORITO	91	PRL 66 1951	S. Orito <i>et al.</i>	(ICEPP, WASCOR, NIHO, ICRR)
ADACHI	90C	PL B244 352	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ADACHI	90E	PL B249 336	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
AKRAWY	90O	PL B252 290	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
HEMMICK	90	PR D41 2074	T.K. Hemmick <i>et al.</i>	(ROCH, MICH, OHIO+)
SAITO	90	PRL 65 2094	T. Saito <i>et al.</i>	(ICRR, KOBE)
NAKAMURA	89	PR D39 1261	T.T. Nakamura <i>et al.</i>	(KYOT, TMT)
NORMAN	89	PR D39 2499	E.B. Norman <i>et al.</i>	(LBL)
BERNSTEIN	88	PR D37 3103	R.M. Bernstein <i>et al.</i>	(STAN, WISC)
LIU	88	PRL 61 271	G. Liu, B. Barish	
BARISH	87	PR D36 2641	B.C. Barish, G. Liu, C. Lane	(CIT)
NORMAN	87	PRL 58 1403	E.B. Norman, S.B. Gazes, D.A. Bennett	(LBL)
BADIER	86	ZPHY C31 21	J. Badier <i>et al.</i>	(NA3 Collab.)
MINCER	85	PR D32 541	A. Mincer <i>et al.</i>	(UMD, GMAS, NSF)
NAKAMURA	85	PL 161B 417	K. Nakamura <i>et al.</i>	(KEK, INUS)
THRON	85	PR D31 451	J.L. Thron <i>et al.</i>	(YALE, FNAL, IOWA)
SAKUYAMA	83B	LNC 37 17	H. Sakuyama, N. Suzuki	(MEIS)
Also		LNC 36 389	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 78A 147	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 6C 371	H. Sakuyama, K. Watanabe	(MEIS)
BHAT	82	PR D25 2820	P.N. Bhat <i>et al.</i>	(TATA)
KINOSHITA	82	PRL 48 77	K. Kinoshita, P.B. Price, D. Fryberger	(UCB+)
MARINI	82	PR D26 1777	A. Marini <i>et al.</i>	(FRAS, LBL, NWES, STAN+)
SMITH	82B	NP B206 333	P.F. Smith <i>et al.</i>	(RAL)
KINOSHITA	81B	PR D24 1707	K. Kinoshita, P.B. Price	(UCB)
LOSECCO	81	PL 102B 209	J.M. LoSecco <i>et al.</i>	(MICH, PENN, BNL)
ULLMAN	81	PRL 47 289	J.D. Ullman	(LEHM, BNL)
YOCK	81	PR D23 1207	P.C.M. Yock	(AUCK)
BARTEL	80	ZPHY C6 295	W. Bartel <i>et al.</i>	(JADE Collab.)
BUSSIERE	80	NP B174 1	A. Bussiere <i>et al.</i>	(BGNA, SACL, LAPP)
YOCK	80	PR D22 61	P.C.M. Yock	(AUCK)
ARMITAGE	79	NP B150 87	J.C.M. Armitage <i>et al.</i>	(CERN, DARE, FOM+)
BOZZOLI	79	NP B159 363	W. Bozzoli <i>et al.</i>	(BGNA, LAPP, SACL+)
GOODMAN	79	PR D19 2572	J.A. Goodman <i>et al.</i>	(UMD)
SMITH	79	NP B149 525	P.F. Smith, J.R.J. Bennett	(RHEL)
BHAT	78	PRAM 10 115	P.N. Bhat, P.V. Ramana Murthy	(TATA)
CARROLL	78	PRL 41 777	A.S. Carroll <i>et al.</i>	(BNL, PRIN)
CUTTS	78	PRL 41 363	D. Cutts <i>et al.</i>	(BROW, FNAL, ILL, BARI+)
VIDAL	78	PL 77B 344	R.A. Vidal <i>et al.</i>	(COLU, FNAL, STON+)
ALEKSEEV	76	SJNP 22 531	G.D. Alekseev <i>et al.</i>	(JINR)
ALEKSEEV	76B	Translated from YAF 22 1021.	G.D. Alekseev <i>et al.</i>	(JINR)
ALEKSEEV	76B	SJNP 23 633	G.D. Alekseev <i>et al.</i>	(JINR)
ALEKSEEV	76B	Translated from YAF 23 1190.	G.D. Alekseev <i>et al.</i>	(JINR)
BALDIN	76	SJNP 22 264	B.Y. Baldin <i>et al.</i>	(JINR)
BALDIN	76	Translated from YAF 22 512.	B.Y. Baldin <i>et al.</i>	(JINR)
BRIATORE	76	NC 31A 553	L. Briatore <i>et al.</i>	(LCGT, FRAS, FREIB)
GUSTAFSON	76	PRL 37 474	H.R. Gustafson <i>et al.</i>	(MICH)
ALBROW	75	NP B97 189	M.G. Albrow <i>et al.</i>	(CERN, DARE, FOM+)
FRANKEL	75	PR D12 2561	S. Frankel <i>et al.</i>	(PENN, FNAL)
JOVANOV...	75	PL 56B 105	J.V. Jovanovich <i>et al.</i>	(MANI, AACH, CERN+)
YOCK	75	NP B86 216	P.C.M. Yock	(AUCK, SLAC)
APPEL	74	PRL 32 428	J.A. Appel <i>et al.</i>	(COLU, FNAL)
FRANKEL	74	PR D9 1932	S. Frankel <i>et al.</i>	(PENN, FNAL)
YOCK	74	NP B76 175	P.C.M. Yock	(AUCK)

ALPER	73	PL 46B 265	B. Alper <i>et al.</i>	(CERN, LIPV, LUND, BOHR+)
LEIPUNER	73	PRL 31 1226	L.B. Leipuner <i>et al.</i>	(BNL, YALE)
DARDO	72	NC 9A 319	M. Dardo <i>et al.</i>	(TORI)
TONWAR	72	JP A5 569	S.C. Tonwar, S. Naranan, B.V. Sreekantan	(TATA)
ANTIPOV	71B	NP B31 235	Y.M. Antipov <i>et al.</i>	(SERP)
ANTIPOV	71C	PL 34B 164	Y.M. Antipov <i>et al.</i>	(SERP)
BINON	69	PL 30B 510	F.G. Binon <i>et al.</i>	(SERP)
BJORNBOE	68	NC B53 241	J. Bjornboe <i>et al.</i>	(BOHR, TATA, BERN+)
JONES	67	PR 164 1584	L.W. Jones	(MICH, WISC, LBL, UCLA, MINN+)
DORFAN	65	PRL 14 999	D.E. Dorfman <i>et al.</i>	(COLU)

---