

$\phi(1680)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

 **$\phi(1680)$  MASS** **$e^+e^-$  PRODUCTION**

| VALUE (MeV)                                                                   | EVTS | DOCUMENT ID              | TECN      | COMMENT                                                         |
|-------------------------------------------------------------------------------|------|--------------------------|-----------|-----------------------------------------------------------------|
| <b>1680 ±20 OUR ESTIMATE</b>                                                  |      |                          |           |                                                                 |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                          |           |                                                                 |
| 1663 ± 5 $^{+16}_{-4}$                                                        |      | ABLIKIM                  | 24CB BES3 | $J/\psi \rightarrow \phi \eta \pi^0$                            |
| 1656.8 ± 4.9                                                                  |      | <sup>1</sup> LICHARD     | 23 RVUE   | $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi \eta \gamma$  |
| 1683 ± 7 ± 9                                                                  |      | <sup>2</sup> ZHU         | 23 BELL   | $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi \eta \gamma$  |
| 1678 $^{+5}_{-3}$ ± 7                                                         |      | <sup>3</sup> ZHU         | 23A RVUE  | $e^+e^- \rightarrow \eta \phi$                                  |
| 1673 ± 5                                                                      |      | <sup>4</sup> ABLIKIM     | 22L BES3  | $2.0\text{--}3.08 e^+e^- \rightarrow K^+K^-\pi^0$               |
| 1680 $^{+12}_{-13}$ ±21 1.8k                                                  |      | <sup>5</sup> ABLIKIM     | 20F BES3  | $\psi(2S) \rightarrow K^+K^-\eta$                               |
| 1662 ±20                                                                      |      | <sup>6</sup> ACHASOV     | 20C SND   | $1.3\text{--}2.0 e^+e^- \rightarrow K^+K^-\pi^0$                |
| 1641 $^{+24}_{-18}$                                                           |      | ACHASOV                  | 19 SND    | $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$                        |
| 1667 ± 5 ±11 3k                                                               |      | <sup>7</sup> IVANOV      | 19A CMD3  | $1.59\text{--}2.007 e^+e^- \rightarrow K^+K^-\eta$              |
| 1700 ±23 2k                                                                   |      | <sup>8</sup> ACHASOV     | 18A SND   | $1.3\text{--}2.0 e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$          |
| 1674 ±12 ± 6 6.2k                                                             |      | <sup>9</sup> LEES        | 14H BABR  | $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$                         |
| 1733 ±10 ±10                                                                  |      | <sup>10</sup> LEES       | 12F BABR  | $10.6 e^+e^- \rightarrow \phi \pi^+ \pi^- \gamma$               |
| 1689 ± 7 ±10 4.8k                                                             |      | <sup>11</sup> SHEN       | 09 BELL   | $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$                |
| 1709 ±20 ±43                                                                  |      | <sup>12</sup> AUBERT     | 08S BABR  | $10.6 e^+e^- \rightarrow \text{hadrons}$                        |
| 1623 ±20 948                                                                  |      | <sup>13</sup> AKHMETSHIN | 03 CMD2   | $1.05\text{--}1.38 e^+e^- \rightarrow K_L^0 K_S^0$              |
| ~ 1500                                                                        |      | <sup>14</sup> ACHASOV    | 98H RVUE  | $e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega \pi^+\pi^-, K^+K^-$ |
| ~ 1900                                                                        |      | <sup>15</sup> ACHASOV    | 98H RVUE  | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$                        |
| 1700 ±20                                                                      |      | <sup>16</sup> CLEGG      | 94 RVUE   | $e^+e^- \rightarrow K^+K^-, K_S^0 K \pi$                        |
| 1657 ±27 367                                                                  |      | BISELLO                  | 91C DM2   | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$                        |
| 1655 ±17                                                                      |      | <sup>17</sup> BISELLO    | 88B DM2   | $e^+e^- \rightarrow K^+K^-$                                     |
| 1680 ±10                                                                      |      | <sup>18</sup> BUON       | 82 DM1    | $e^+e^- \rightarrow \text{hadrons}$                             |
| 1677 ±12                                                                      |      | <sup>19</sup> MANE       | 82 DM1    | $e^+e^- \rightarrow K_S^0 K \pi$                                |

<sup>1</sup> From a VDM fit to ZHU 23  $\eta \phi \gamma$  data with two resonances,  $\phi(1680)$ ,  $\phi(2170)$ , and a third resonance with mass  $1850.7 \pm 5.3$  MeV and width  $25 \pm 35$  MeV of  $1.7 \sigma$  statistical evidence.

<sup>2</sup> From a fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution.

<sup>3</sup> From the analysis of the combined measurements of  $\sigma(e^+e^- \rightarrow \eta \phi)$  from BaBar, Belle, BESIII, CMD3.

<sup>4</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

- <sup>5</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .
- <sup>6</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .
- <sup>7</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.
- <sup>8</sup> Assuming the  $K\bar{K}^*(892) + \text{c.c.}$  dynamics. Systematic uncertainties not estimated.
- <sup>9</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .
- <sup>10</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.
- <sup>11</sup> From a fit with two incoherent Breit-Wigners.
- <sup>12</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.
- <sup>13</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.
- <sup>14</sup> Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.
- <sup>15</sup> Using the data from BISELLO 91C.
- <sup>16</sup> Using BISELLO 88B and MANE 82 data.
- <sup>17</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$  assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitation.
- <sup>18</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0K_L^0$ ,  $K_S^0K^\pm\pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.
- <sup>19</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

 **$p\bar{p}$  ANNIHILATION**

| VALUE (MeV)                                                                   | DOCUMENT ID         | TECN | COMMENT                                     |
|-------------------------------------------------------------------------------|---------------------|------|---------------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |      |                                             |
| $1700 \pm 8$                                                                  | <sup>1</sup> AMSLER | 06   | CBAR $0.9 \bar{p}p \rightarrow K^+K^-\pi^0$ |
| <sup>1</sup> Could also be $\rho(1700)$ .                                     |                     |      |                                             |

 **$\phi(1680)$  WIDTH** **$e^+e^-$  PRODUCTION**

| VALUE (MeV)                                                                                                                                                  | EVTS | DOCUMENT ID          | TECN      | COMMENT                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------|------|----------------------|-----------|--------------------------------------------------------------|
| <b>150 <math>\pm</math> 50 OUR ESTIMATE</b> This is only an educated guess; the error given is larger than the error on the average of the published values. |      |                      |           |                                                              |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                                                |      |                      |           |                                                              |
| $159 \pm 15 \pm 11$                                                                                                                                          |      | ABLIKIM              | 24CB BES3 | $J/\psi \rightarrow \phi\eta\pi^0$                           |
| $150.8 \pm 7.0$                                                                                                                                              |      | <sup>1</sup> LICHARD | 23 RVUE   | $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$ |
| $149 \pm 12 \pm 13$                                                                                                                                          |      | <sup>2</sup> ZHU     | 23 BELL   | $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$ |
| $156 \pm 5 \pm 9$                                                                                                                                            |      | <sup>3</sup> ZHU     | 23A RVUE  | $e^+e^- \rightarrow \eta\phi$                                |
| $172 \pm 8$                                                                                                                                                  |      | <sup>4</sup> ABLIKIM | 22L BES3  | $2.0\text{--}3.08 e^+e^- \rightarrow K^+K^-\pi^0$            |
| $185^{+30}_{-26} {}^{+25}_{-47}$                                                                                                                             | 1.8k | <sup>5</sup> ABLIKIM | 20F BES3  | $\psi(2S) \rightarrow K^+K^-\eta$                            |
| $159 \pm 32$                                                                                                                                                 |      | <sup>6</sup> ACHASOV | 20C SND   | $1.3\text{--}2.0 e^+e^- \rightarrow K^+K^-\pi^0$             |
| $103^{+26}_{-24}$                                                                                                                                            |      | ACHASOV              | 19 SND    | $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$                     |
| $176 \pm 23 \pm 38$                                                                                                                                          | 3k   | <sup>7</sup> IVANOV  | 19A CMD3  | $1.59\text{--}2.007 e^+e^- \rightarrow K^+K^-\eta$           |

|     |          |           |                      |                          |          |                                                         |
|-----|----------|-----------|----------------------|--------------------------|----------|---------------------------------------------------------|
| 300 | $\pm 50$ | 2k        | <sup>8</sup> ACHASOV | 18A                      | SND      | $1.3\text{--}2.0\ e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$ |
| 165 | $\pm 38$ | $\pm 70$  | 6.2k                 | <sup>9</sup> LEES        | 14H BABR | $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$                 |
| 300 | $\pm 15$ | $\pm 37$  |                      | <sup>10</sup> LEES       | 12F BABR | $10.6\ e^+e^- \rightarrow \phi \pi^+ \pi^- \gamma$      |
| 211 | $\pm 14$ | $\pm 19$  | 4.8k                 | <sup>11</sup> SHEN       | 09 BELL  | $10.6\ e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$   |
| 322 | $\pm 77$ | $\pm 160$ |                      | <sup>12</sup> AUBERT     | 08S BABR | $10.6\ e^+e^- \rightarrow \text{hadrons}$               |
| 139 | $\pm 60$ |           | 948                  | <sup>13</sup> AKHMETSHIN | 03 CMD2  | $1.05\text{--}1.38\ e^+e^- \rightarrow K_L^0 K_S^0$     |
| 300 | $\pm 60$ |           |                      | <sup>14</sup> CLEGG      | 94 RVUE  | $e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$               |
| 146 | $\pm 55$ |           | 367                  | BISELLO                  | 91C DM2  | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$                |
| 207 | $\pm 45$ |           |                      | <sup>15</sup> BISELLO    | 88B DM2  | $e^+e^- \rightarrow K^+ K^-$                            |
| 185 | $\pm 22$ |           |                      | <sup>16</sup> BUON       | 82 DM1   | $e^+e^- \rightarrow \text{hadrons}$                     |
| 102 | $\pm 36$ |           |                      | <sup>17</sup> MANE       | 82 DM1   | $e^+e^- \rightarrow K_S^0 K \pi$                        |

<sup>1</sup> From a VDM fit to ZHU 23  $\eta\phi\gamma$  data with two resonances,  $\phi(1680)$ ,  $\phi(2170)$ , and a third resonance with mass  $1850.7 \pm 5.3$  MeV and width  $25 \pm 35$  MeV of 1.7  $\sigma$  statistical evidence.

<sup>2</sup> From a fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution.

<sup>3</sup> From the analysis of the combined measurements of  $\sigma(e^+e^- \rightarrow \eta\phi)$  from BaBar, Belle, BESIII, CMD3.

<sup>4</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

<sup>5</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .

<sup>6</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .

<sup>7</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

<sup>8</sup> Assuming the  $K\bar{K}^*(892) + \text{c.c.}$  dynamics. Systematic uncertainties not estimated.

<sup>9</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

<sup>10</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.

<sup>11</sup> From a fit with two incoherent Breit-Wigners.

<sup>12</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>13</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.

<sup>14</sup> Using BISELLO 88B and MANE 82 data.

<sup>15</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$

<sup>16</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm \pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.

<sup>17</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

## **$p\bar{p}$ ANNIHILATION**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|---------|
|-------------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|              |                     |    |                                                |
|--------------|---------------------|----|------------------------------------------------|
| 143 $\pm$ 24 | <sup>1</sup> AMSLER | 06 | CBAR $0.9\ \bar{p}p \rightarrow K^+ K^- \pi^0$ |
|--------------|---------------------|----|------------------------------------------------|

<sup>1</sup> Could also be  $\rho(1700)$ .

**$\phi(1680)$  DECAY MODES**

| Mode                               | Fraction ( $\Gamma_i/\Gamma$ ) |
|------------------------------------|--------------------------------|
| $\Gamma_1$ $K\bar{K}^*(892)+$ c.c. | seen                           |
| $\Gamma_2$ $K_S^0 K\pi$            | seen                           |
| $\Gamma_3$ $K\bar{K}$              | seen                           |
| $\Gamma_4$ $K_L^0 K_S^0$           |                                |
| $\Gamma_5$ $e^+e^-$                | seen                           |
| $\Gamma_6$ $\omega\pi\pi$          | not seen                       |
| $\Gamma_7$ $\phi\pi\pi$            |                                |
| $\Gamma_8$ $K^+K^-\pi^+\pi^-$      | seen                           |
| $\Gamma_9$ $\eta\phi$              | seen                           |
| $\Gamma_{10}$ $\eta\gamma$         | seen                           |
| $\Gamma_{11}$ $K^+K^-\pi^0$        |                                |
| $\Gamma_{12}$ $f'_2(1525)\gamma$   | not seen                       |

 **$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the integrated cross section into channel (I) in  $e^+e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(I)$  or the branching ratio  $\Gamma(I)/\text{total}$ .

 **$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_4\Gamma_5/\Gamma$** 

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.3 $\pm$ 2.4 $\pm$ 6.2      6.2k      <sup>1</sup> LEES      14H BABR       $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$

<sup>1</sup>Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

 **$\Gamma(\phi\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_7\Gamma_5/\Gamma$** 

| VALUE ( $10^{-2}$ keV) | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|---------|
|------------------------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 $\pm$ 0.2 $\pm$ 0.3      LEES      12F BABR      10.6  $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$

 **$\Gamma(\eta\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_9\Gamma_5/\Gamma$** 

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

122 $\pm$  6 $\pm$ 13      <sup>1</sup> ZHU      23 BELL       $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$

65 $\pm$  5 $\pm$ 13 to      <sup>2</sup> ZHU      23A RVUE       $e^+e^- \rightarrow \eta\phi$

215 $\pm$  8 $\pm$ 11

94 $\pm$ 13 $\pm$ 15      3k      <sup>3</sup> IVANOV      19A CMD3      1.59–2.007  $e^+e^- \rightarrow K^+K^-\eta$

<sup>1</sup>From a solution of the fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution. Other solutions with equal fit quality give (219  $\pm$  15  $\pm$  18) eV, (163  $\pm$  11  $\pm$  13) eV and (203  $\pm$  12  $\pm$  18) eV.

<sup>2</sup>From the analysis of the combined measurements of  $\sigma(e^+e^- \rightarrow \eta\phi)$  from BaBar, Belle, BESIII, CMD3. Four solutions are found, with equal fit quality:  $(79 \pm 4 \pm 16)$  eV,  $(127 \pm 5 \pm 12)$  eV,  $(65^{+5}_{-4} \pm 13)$  eV,  $(215^{+8}_{-5} \pm 11)$  eV.

<sup>3</sup>From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

### $\phi(1680) \Gamma(i) \Gamma(e^+e^-) / \Gamma^2(\text{total})$

This combination of a branching ratio into channel (*i*) and branching ratio into  $e^+e^-$  is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (*i*) or  $e^+e^-$ .

$$\Gamma(K_L^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \qquad \Gamma_4 / \Gamma \times \Gamma_5 / \Gamma$$

| VALUE (units $10^{-6}$ )                                                      | EVTS | DOCUMENT ID                | TECN | COMMENT                                              |
|-------------------------------------------------------------------------------|------|----------------------------|------|------------------------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                            |      |                                                      |
| $0.131 \pm 0.059$                                                             | 948  | <sup>1</sup> AKHMETSHIN 03 | CMD2 | $1.05\text{--}1.38 \ e^+e^- \rightarrow K_L^0 K_S^0$ |

<sup>1</sup>From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known. Recalculated by us.

$$\Gamma(K \bar{K}^*(892) + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \qquad \Gamma_1 / \Gamma \times \Gamma_5 / \Gamma$$

| VALUE (units $10^{-6}$ )                                                      | EVTS | DOCUMENT ID              | TECN | COMMENT                                                           |
|-------------------------------------------------------------------------------|------|--------------------------|------|-------------------------------------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                          |      |                                                                   |
| $1.15 \pm 0.16 \pm 0.01$                                                      |      | <sup>1</sup> AUBERT 08S  | BABR | $10.6 \ e^+e^- \rightarrow K \bar{K}^*(892) \gamma + \text{c.c.}$ |
| $3.29 \pm 1.57$                                                               | 367  | <sup>2</sup> BISELLO 91C | DM2  | $1.35\text{--}2.40 \ e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$      |

<sup>1</sup>From the simultaneous fit to the  $K \bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>2</sup>Recalculated by us with the published value of  $B(K \bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)$ .

$$\Gamma(\phi\pi\pi) / \Gamma_{\text{total}} \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \qquad \Gamma_7 / \Gamma \times \Gamma_5 / \Gamma$$

| VALUE (units $10^{-7}$ )                                                                                                                | EVTS | DOCUMENT ID          | TECN | COMMENT                                                |
|-----------------------------------------------------------------------------------------------------------------------------------------|------|----------------------|------|--------------------------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                           |      |                      |      |                                                        |
| $1.86 \pm 0.14 \pm 0.21$                                                                                                                | 4.8k | <sup>1</sup> SHEN 09 | BELL | $10.6 \ e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ |
| <sup>1</sup> Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ . |      |                      |      |                                                        |

$$\Gamma(\eta\phi) / \Gamma_{\text{total}} \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \qquad \Gamma_9 / \Gamma \times \Gamma_5 / \Gamma$$

| VALUE (units $10^{-7}$ )                                                      | EVTS | DOCUMENT ID             | TECN | COMMENT                                                |
|-------------------------------------------------------------------------------|------|-------------------------|------|--------------------------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                         |      |                                                        |
| $5.64^{+1.74}_{-1.80}$                                                        |      | ACHASOV 19              | SND  | $e^+e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$            |
| $5.3 \pm 0.6 \pm 0.9$                                                         | 3k   | <sup>1</sup> IVANOV 19A | CMD3 | $1.59\text{--}2.007 \ e^+e^- \rightarrow K^+ K^- \eta$ |
| $4.3 \pm 1.0 \pm 0.9$                                                         |      | <sup>2</sup> AUBERT 08S | BABR | $10.6 \ e^+e^- \rightarrow \phi\eta\gamma$             |

<sup>1</sup>From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

<sup>2</sup>From the simultaneous fit to the  $K \bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

$\phi(1680)$  BRANCHING RATIOS

| $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(K_S^0 K\pi)$ |             |      |         | $\Gamma_1/\Gamma_2$                      |
|----------------------------------------------------------|-------------|------|---------|------------------------------------------|
| VALUE                                                    | DOCUMENT ID | TECN | COMMENT |                                          |
| dominant                                                 | MANE        | 82   | DM1     | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$ |

| $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+\text{c.c.})$                        |             |      |         | $\Gamma_3/\Gamma_1$ |
|-------------------------------------------------------------------------------|-------------|------|---------|---------------------|
| VALUE                                                                         | DOCUMENT ID | TECN | COMMENT |                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |         |                     |
| $0.07 \pm 0.01$                                                               | BUON        | 82   | DM1     | $e^+e^-$            |

| $\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892)+\text{c.c.})$ |             |      |         | $\Gamma_6/\Gamma_1$ |
|------------------------------------------------------------|-------------|------|---------|---------------------|
| VALUE                                                      | DOCUMENT ID | TECN | COMMENT |                     |
| <b>&lt;0.10</b>                                            | BUON        | 82   | DM1     | $e^+e^-$            |

| $\Gamma(\eta\phi)/\Gamma_{\text{total}}$ |      |                      |      | $\Gamma_9/\Gamma$                             |
|------------------------------------------|------|----------------------|------|-----------------------------------------------|
| VALUE                                    | EVTS | DOCUMENT ID          | TECN | COMMENT                                       |
| seen                                     | 35   | <sup>1</sup> ACHASOV | 14   | SND 1.15–2.00 $e^+e^- \rightarrow \eta\gamma$ |

<sup>1</sup>From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

| $\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892)+\text{c.c.})$                        |                     |      |         | $\Gamma_9/\Gamma_1$                      |
|-------------------------------------------------------------------------------|---------------------|------|---------|------------------------------------------|
| VALUE                                                                         | DOCUMENT ID         | TECN | COMMENT |                                          |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |      |         |                                          |
| $\approx 0.37$                                                                | <sup>1</sup> AUBERT | 08S  | BABR    | 10.6 $e^+e^- \rightarrow \text{hadrons}$ |

<sup>1</sup>From the fit including data from AUBERT 07AK.

| $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ |      |                      |      | $\Gamma_{10}/\Gamma$                          |
|--------------------------------------------|------|----------------------|------|-----------------------------------------------|
| VALUE                                      | EVTS | DOCUMENT ID          | TECN | COMMENT                                       |
| seen                                       | 35   | <sup>1</sup> ACHASOV | 14   | SND 1.15–2.00 $e^+e^- \rightarrow \eta\gamma$ |

<sup>1</sup>From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

| $\Gamma(f_2'(1525)\gamma)/\Gamma_{\text{total}}$ |                      |      |         | $\Gamma_{12}/\Gamma$                          |
|--------------------------------------------------|----------------------|------|---------|-----------------------------------------------|
| VALUE                                            | DOCUMENT ID          | TECN | COMMENT |                                               |
| not seen                                         | <sup>1</sup> ACHASOV | 22   | SND     | 1.17–2.00 $e^+e^- \rightarrow \eta\eta\gamma$ |

<sup>1</sup>The 90% CL upper limit on the Born cross sections  $\sigma(e^+e^- \rightarrow \phi(1680) \rightarrow f_2'(1525)\gamma \rightarrow \eta\eta\gamma)$  and  $\sigma(e^+e^- \rightarrow \rho(1700) \rightarrow f_0(1500)\gamma \rightarrow \eta\eta\gamma)$  is 10.6 pb.

 $\phi(1680)$  REFERENCES

|         |      |                |                            |                  |
|---------|------|----------------|----------------------------|------------------|
| ABLIKIM | 24CB | PR D110 112014 | M. Ablikim <i>et al.</i>   | (BESIII Collab.) |
| LICHARD | 23   | PR D108 092005 | P. Lichard                 | (OPAV, CTUP)     |
| ZHU     | 23   | PR D107 012006 | W. Zhu <i>et al.</i>       | (BELLE Collab.)  |
| ZHU     | 23A  | CP C47 113003  | W. Zhu, X. Wang            | (RVUE)           |
| ABLIKIM | 22L  | JHEP 2207 045  | M. Ablikim <i>et al.</i>   | (BESIII Collab.) |
| ACHASOV | 22   | EPJ C82 168    | M.N. Achasov <i>et al.</i> | (SND Collab.)    |
| ABLIKIM | 20F  | PR D101 032008 | M. Ablikim <i>et al.</i>   | (BESIII Collab.) |
| ACHASOV | 20C  | EPJ C80 1139   | M.N. Achasov <i>et al.</i> | (SND Collab.)    |
| ACHASOV | 19   | PR D99 112004  | M.N. Achasov <i>et al.</i> | (SND Collab.)    |
| IVANOV  | 19A  | PL B798 134946 | V.L. Ivanov <i>et al.</i>  | (CMD-3 Collab.)  |

|            |      |                               |                                                |                             |
|------------|------|-------------------------------|------------------------------------------------|-----------------------------|
| ACHASOV    | 18A  | PR D97 032011                 | M.N. Achasov <i>et al.</i>                     | (SND Collab.)               |
| ACHASOV    | 14   | PR D90 032002                 | M.N. Achasov <i>et al.</i>                     | (SND Collab.)               |
| LEES       | 14H  | PR D89 092002                 | J.P. Lees <i>et al.</i>                        | (BABAR Collab.)             |
| LEES       | 12F  | PR D86 012008                 | J.P. Lees <i>et al.</i>                        | (BABAR Collab.)             |
| PDG        | 12   | PR D86 010001                 | J. Beringer <i>et al.</i>                      | (PDG Collab.)               |
| SHEN       | 09   | PR D80 031101                 | C.P. Shen <i>et al.</i>                        | (BELLE Collab.)             |
| AUBERT     | 08S  | PR D77 092002                 | B. Aubert <i>et al.</i>                        | (BABAR Collab.)             |
| AUBERT     | 07AK | PR D76 012008                 | B. Aubert <i>et al.</i>                        | (BABAR Collab.)             |
| AMSLER     | 06   | PL B639 165                   | C. Amsler <i>et al.</i>                        | (Crystal Barrel Collab.)    |
| AKHMETSHIN | 03   | PL B551 27                    | R.R. Akhmetshin <i>et al.</i>                  | (Novosibirsk CMD-2 Collab.) |
| Also       |      | PAN 65 1222                   | E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin |                             |
| ACHASOV    | 98H  | PR D57 4334                   | N.N. Achasov, A.A. Kozhevnikov                 |                             |
| CLEGG      | 94   | ZPHY C62 455                  | A.B. Clegg, A. Donnachie                       | (LANC, MCHS)                |
| ANTONELLI  | 92   | ZPHY C56 15                   | A. Antonelli <i>et al.</i>                     | (DM2 Collab.)               |
| BISELLO    | 91C  | ZPHY C52 227                  | D. Bisello <i>et al.</i>                       | (DM2 Collab.)               |
| DOLINSKY   | 91   | PRPL 202 99                   | S.I. Dolinsky <i>et al.</i>                    | (NOVO)                      |
| BISELLO    | 88B  | ZPHY C39 13                   | D. Bisello <i>et al.</i>                       | (PADO, CLER, FRAS+)         |
| BARKOV     | 87   | JETPL 46 164                  | L.M. Barkov <i>et al.</i>                      | (NOVO)                      |
|            |      | Translated from ZETFP 46 132. |                                                |                             |
| BUON       | 82   | PL 118B 221                   | J. Buon <i>et al.</i>                          | (LALO, MONP)                |
| MANE       | 82   | PL 112B 178                   | F. Mane <i>et al.</i>                          | (LALO)                      |
| IVANOV     | 81   | PL 107B 297                   | P.M. Ivanov <i>et al.</i>                      | (NOVO)                      |
| MANE       | 81   | PL 99B 261                    | F. Mane <i>et al.</i>                          | (ORSAY)                     |