

$\Upsilon_2(1D)$

$$J^{PC} = 0^-(2^--)$$

was $\Upsilon(1D)$

First observed by BONVICINI 04 in the decay to $\gamma\gamma\Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+\pi^-\Upsilon(1S)$.
Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

$\Upsilon_2(1D)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10163.7±1.4 OUR AVERAGE		Error includes scale factor of 1.7.		
10164.5±0.8±0.5		DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1±0.6±1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Upsilon_2(1D)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma\gamma\Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta\Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+\pi^-\Upsilon(1S)$	$(6.6\pm1.6)\times10^{-3}$

$\Upsilon_2(1D)$ BRANCHING RATIOS

$\Gamma(\eta\Upsilon(1S))/\Gamma(\gamma\gamma\Upsilon(1S))$				Γ_3/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Gamma(\pi^+\pi^-\Upsilon(1S))/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	
$0.66^{+0.15}_{-0.14} \pm 0.06$	¹ DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$	

¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon_2(1D))$.

$\Gamma(\pi^+\pi^-\Upsilon(1S))/\Gamma(\gamma\gamma\Upsilon(1S))$				Γ_4/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	² BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

² Assuming $J = 2$.

$\Upsilon_2(1D)$ REFERENCES

DEL-AMO-SA... 10R	PR D82 111102	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
BONVICINI 04	PR D70 032001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)