

$\Upsilon(3S)$


$$J^{PC} = 0^{--}(1^{--})$$

$\Upsilon(3S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.1±0.5	¹ SHAMOV 23	RVUE	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10355.2±0.5	^{2,3} ARTAMONOV 00	MD1	$e^+e^- \rightarrow \text{hadrons}$
10355.3±0.5	^{4,5} BARU 86B	MD1	$e^+e^- \rightarrow \text{hadrons}$
¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.			
² Reanalysis of BARU 86B using new electron mass (COHEN 87).			
³ Superseded by SHAMOV 23.			
⁴ Reanalysis of ARTAMONOV 84.			
⁵ Superseded by ARTAMONOV 00.			

$m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
331.82±0.08 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
331.77±0.07±0.10	1.8M	⁶ AAIJ	24AC LHCb	$\Upsilon \rightarrow \mu^+\mu^-$
331.88±0.03±0.04	1.5k	⁶ AAIJ	24AC LHCb	$\Upsilon(3S) \rightarrow \Upsilon(2S)\pi^+\pi^-$
331.50±0.02±0.13		LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
⁶ Observed in prompt pp production.				



$\Upsilon(3S)$ WIDTH

VALUE (keV)

DOCUMENT ID

 20.32 ± 1.85 OUR EVALUATIONSee the Note on “Width Determinations of the Υ States” **$\Upsilon(3S)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(2S)$ anything	(10.6 $\pm$ 0.8) %	
Γ_2 $\Upsilon(2S) \pi^+ \pi^-$	(2.82 $\pm$ 0.18) %	S=1.6
Γ_3 $\Upsilon(2S) \pi^0 \pi^0$	(1.85 $\pm$ 0.14) %	
Γ_4 $\Upsilon(2S) \gamma \gamma$	(5.0 $\pm$ 0.7) %	
Γ_5 $\Upsilon(2S) \pi^0$	< 5.1 $\times 10^{-4}$	CL=90%
Γ_6 $\Upsilon(1S) \pi^+ \pi^-$	(4.37 $\pm$ 0.08) %	
Γ_7 $\Upsilon(1S) \pi^0 \pi^0$	(2.20 $\pm$ 0.13) %	
Γ_8 $\Upsilon(1S) \eta$	< 1 $\times 10^{-4}$	CL=90%
Γ_9 $\Upsilon(1S) \pi^0$	< 7 $\times 10^{-5}$	CL=90%
Γ_{10} $h_b(1P) \pi^0$	< 1.2 $\times 10^{-3}$	CL=90%
Γ_{11} $h_b(1P) \pi^0 \rightarrow \gamma \eta_b(1S) \pi^0$	(4.3 $\pm$ 1.4) $\times 10^{-4}$	
Γ_{12} $h_b(1P) \pi^+ \pi^-$	< 1.2 $\times 10^{-4}$	CL=90%
Γ_{13} $\tau^+ \tau^-$	(2.29 $\pm$ 0.30) %	
Γ_{14} $\mu^+ \mu^-$	(2.18 $\pm$ 0.21) %	S=2.1
Γ_{15} $e^+ e^-$	(2.18 $\pm$ 0.20) %	
Γ_{16} hadrons	(93 $\pm$ 12) %	
Γ_{17} $g g g$	(35.7 $\pm$ 2.6) %	
Γ_{18} $\gamma g g$	(9.7 $\pm$ 1.8) $\times 10^{-3}$	
Γ_{19} 2H anything	(2.33 $\pm$ 0.33) $\times 10^{-5}$	

Radiative decays

Γ_{20} $\gamma \chi_{b2}(2P)$	(13.1 $\pm$ 1.6) %	S=3.4
Γ_{21} $\gamma \chi_{b1}(2P)$	(12.6 $\pm$ 1.2) %	S=2.4
Γ_{22} $\gamma \chi_{b0}(2P)$	(5.9 $\pm$ 0.6) %	S=1.4
Γ_{23} $\gamma \chi_{b2}(1P)$	(10.0 $\pm$ 1.0) $\times 10^{-3}$	S=1.7
Γ_{24} $\gamma \chi_{b1}(1P)$	(9 $\pm$ 5) $\times 10^{-4}$	S=1.8
Γ_{25} $\gamma \chi_{b0}(1P)$	(2.7 $\pm$ 0.4) $\times 10^{-3}$	
Γ_{26} $\gamma \eta_b(2S)$	< 6.2 $\times 10^{-4}$	CL=90%
Γ_{27} $\gamma \eta_b(1S)$	(5.1 $\pm$ 0.7) $\times 10^{-4}$	
Γ_{28} $\gamma A^0 \rightarrow \gamma$ hadrons	< 8 $\times 10^{-5}$	CL=90%
Γ_{29} $\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 $\times 10^{-4}$	CL=95%
Γ_{30} $\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	< 5.5 $\times 10^{-6}$	CL=90%
Γ_{31} $\gamma A^0 \rightarrow \gamma \tau^+ \tau^-$	[b] < 1.6 $\times 10^{-4}$	CL=90%

Lepton Family number (LF) violating modes

Γ_{32} $e^\pm \tau^\mp$	LF	< 4.2 $\times 10^{-6}$	CL=90%
--------------------------------	----	------------------------	--------

Γ_{33}	$e^\pm \mu^\mp$	LF	< 3.6	$\times 10^{-7}$	CL=90%
Γ_{34}	$\mu^\pm \tau^\mp$	LF	< 3.1	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ [b] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. **$\Upsilon(3S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$** **$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$** **$\Gamma_{16} \Gamma_{15} / \Gamma$**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.414 ± 0.007 OUR AVERAGE0.413 ± 0.004 ± 0.006 ROSNER 06 CLEO 10.4 $e^+ e^- \rightarrow \text{hadrons}$ 0.45 ± 0.03 ± 0.03 ⁷ GILES 84B CLEO $e^+ e^- \rightarrow \text{hadrons}$ ⁷ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85. **$\Gamma(\Upsilon(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$** **$\Gamma_6 \Gamma_{15} / \Gamma$**

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

18.46 ± 0.27 ± 0.77 6.4k ⁸ AUBERT 08BP BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$ ⁸ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$. **$\Upsilon(3S)$ PARTIAL WIDTHS** **$\Gamma(e^+ e^-)$** **Γ_{15}**

VALUE (keV)	DOCUMENT ID
-------------	-------------

0.443 ± 0.008 OUR EVALUATION **$\Upsilon(3S)$ BRANCHING RATIOS** **$\Gamma(\Upsilon(2S) \text{anything}) / \Gamma_{\text{total}}$** **$\Gamma_1 / \Gamma$**

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.106 ± 0.008 OUR AVERAGE0.1023 ± 0.0105 4625 ^{9,10,11} BUTLER 94B CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- X$ 0.111 ± 0.012 4891 ^{10,11,12} BROCK 91 CLEO $e^+ e^- \rightarrow \pi^+ \pi^- X,$
 $\pi^+ \pi^- \ell^+ \ell^-$ ⁹ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-)$.¹⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.¹¹ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-) = (18.5 \pm 0.8)\%$.¹² Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality. **$\Gamma(\Upsilon(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$** **$\Gamma_2 / \Gamma$**

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.82 ± 0.18 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.3.00 ± 0.02 ± 0.14 543k LEES 11C BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$ 2.40 ± 0.10 ± 0.26 800 ¹³ AUBERT 08BP BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- e^+ e^-$

3.12 ± 0.49	980	^{14,15} BUTLER	94B	CLE2	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13 ± 0.38	974	¹⁶ BROCK	91	CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.82 \pm 0.65 \pm 0.53$	138	¹⁶ WU	93	CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.1 ± 2.0	5	MAGERAS	82	CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$


¹³ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

¹⁴ From the exclusive mode.

¹⁵ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

¹⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

WEIGHTED AVERAGE
 2.82 ± 0.18 (Error scaled by 1.6)



$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85 ± 0.14 OUR AVERAGE				
$1.82 \pm 0.09 \pm 0.12$	4391	¹⁷ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16 ± 0.39		^{18,19} BUTLER	94B	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
$1.7 \pm 0.5 \pm 0.2$	10	²⁰ HEINTZ	92	CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹⁷ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

¹⁸ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

¹⁹ From the exclusive mode.

²⁰ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0502±0.0069	²¹ BUTLER	94B	CLE2 $e^+e^- \rightarrow \ell^+\ell^-2\gamma$

²¹ From the exclusive mode. $\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	²² HE	08A	CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

²² Authors assume $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$. $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.37±0.08 OUR AVERAGE				
4.32±0.07±0.13	90k	²³ LEES	11L	BABR $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.01±0.13	190k	²⁴ BHARI	09	CLEO $e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17±0.06±0.19	6.4k	²⁵ AUBERT	08BP	BABR $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52±0.35	11830	²⁶ BUTLER	94B	CLE2 $e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$
4.46±0.34±0.50	451	²⁶ WU	93	CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.30	11221	²⁶ BROCK	91	CLEO $e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ±1.0 22 GREEN 82 CLEO $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ 3.9 ±1.3 26 MAGERAS 82 CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ ²³ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.²⁴ A weighted average of the inclusive and exclusive results.²⁵ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.²⁶ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality. $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.577±0.026±0.060	800	²⁷ AUBERT	08BP	BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

²⁷ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.

Not independent of other values reported by AUBERT 08BP.

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.13 OUR AVERAGE				
2.24±0.09±0.11	6584	²⁸ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99±0.34	56	²⁹ BUTLER	94B	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 ±0.4 ±0.3	33	³⁰ HEINTZ	92	CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

²⁸ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.²⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.³⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.501 ± 0.043 ³¹ BHARI 09 CLEO $e^+e^- \rightarrow \Upsilon(3S)$

³¹ Not independent of other values reported by BHARI 09.

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.1 90 ³² LEES 11L BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8 90 ^{32,33} AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$

<0.18 90 ³⁴ HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$

<2.2 90 BROCK 91 CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$

³² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³³ Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

³⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.23 90 ³⁵ LEES 11L BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9 90 ³⁶ AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

³⁵ Not independent of other values reported by LEES 11L.

³⁶ Not independent of other values reported by AUBERT 08BP.

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.07 90 ³⁷ HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

³⁷ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

$\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.2 \times 10^{-3}$ 90 ³⁸ GE 11 CLEO $\Upsilon(3S) \rightarrow \pi^0$ anything

³⁸ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.

$\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.3 \pm 1.1 \pm 0.9$ LEES 11K BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.2 90 ³⁹ LEES 11C BABR $e^+e^- \rightarrow \pi^+\pi^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18 ³⁹ BUTLER 94B CLE2 $e^+e^- \rightarrow \pi^+\pi^-X$

<15 ³⁹ BROCK 91 CLEO $e^+e^- \rightarrow \pi^+\pi^-X$

³⁹ For $M(h_b(1P)) = 9900$ MeV.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.29 \pm 0.21 \pm 0.22$	15k	⁴⁰ BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$

⁴⁰ BESSON 07 reports $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$


Γ_{13}/Γ_{14}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.968 ± 0.016 OUR AVERAGE				
$0.966 \pm 0.008 \pm 0.014$	2.2M	LEES	20E	BABR $e^+e^- \rightarrow \Upsilon(3S)$
$1.05 \pm 0.08 \pm 0.05$	15k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(3S)$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0218 ± 0.0021 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
$0.0239 \pm 0.0007 \pm 0.0010$	81k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
$0.0202 \pm 0.0019 \pm 0.0033$		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
$0.0173 \pm 0.0015 \pm 0.0011$		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
$0.033 \pm 0.013 \pm 0.007$	1096	ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$



$\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
35.7 ± 2.6	3M	⁴¹ BESSON	06A	CLEO $\Upsilon(3S) \rightarrow \text{hadrons}$

⁴¹ Calculated using BESSON 06A value of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\Upsilon(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+ \pi^- \Upsilon(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0 \pi^0 \Upsilon(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma \chi_{b2}(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma \chi_{b1}(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma \chi_{b0}(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+ \mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ BESSON 06A value.

 $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.18	60k	⁴² BESSON	06A	CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

⁴² Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(g g g)/\Gamma_{\text{total}}$ BESSON 06A value.

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_{18}/Γ_{17}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.72 \pm 0.06 \pm 0.49$	3M	BESSON	06A	CLEO $\Upsilon(3S) \rightarrow (\gamma +) \text{hadrons}$


 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.33 \pm 0.15^{+0.31}_{-0.28}$	LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$


 $\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.131 ± 0.016 OUR AVERAGE		Error includes scale factor of 3.4. See the ideogram below.		
$0.1579 \pm 0.0017 \pm 0.0073$	568k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
$0.111 \pm 0.005 \pm 0.004$	10319	⁴³ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$
$0.135 \pm 0.003 \pm 0.017$	30741	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$

⁴³ Supersedes NARAIN 91.




$\Gamma(\gamma\chi_{b1}(2P))/\Gamma_{\text{total}}$						Γ_{21}/Γ
VALUE			EVTS	DOCUMENT ID	TECN	COMMENT
0.126 ± 0.012	OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.				
$0.1454 \pm 0.0018 \pm 0.0073$		537k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$
$0.115 \pm 0.005 \pm 0.005$		11147	44 HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma X$
$0.105 \pm 0.003 \pm 0.013$		25759	MORRISON	91	CLE2	$e^+e^- \rightarrow \gamma X$



⁴⁴Supersedes NARAIN 91.

$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{\text{total}}$					Γ_{22}/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.059 ±0.006 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.					
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.060 ±0.004 ±0.006	4959	⁴⁵ HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma X$	
0.049 ^{+0.003} _{-0.004} ±0.006	9903	MORRISON	91	CLE2	$e^+e^- \rightarrow \gamma X$	

⁴⁵Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$					Γ_{23}/Γ	
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
10.0 ± 1.0 OUR AVERAGE	Error includes scale factor of 1.7.					
8.0 ± 1.3 ± 0.4	126	^{46,47} KORNICER	11	CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$	
10.5 ± 0.3 $^{+0.7}_{-0.6}$	9.7k	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	⁴⁸ ASNER	08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen		⁴⁹ HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

⁴⁶ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁴⁷ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S))]$
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S)) = (18.0 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁸ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))]$
 $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) =$
 7.15×10^{-2} .

⁴⁹ HEINTZ 92, while unable to distinguish between different J states, measures
 $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for J
 $= 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using
 $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.5 OUR AVERAGE Error includes scale factor of 1.8.					
$1.5 \pm 0.4 \pm 0.1$		50	^{50,51} KORNICER	11 CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
$0.5 \pm 0.3^{+0.2}_{-0.1}$			LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.7 seen	90	⁵² ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
		⁵³ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

⁵⁰ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁵¹ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S))] =$
 $(5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S)) =$
 $(35.2 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error
is the systematic error from using our best value.

⁵² ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] <$
 2.5×10^{-2} which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

⁵³ HEINTZ 92, while unable to distinguish between different J states, measures
 $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for J
 $= 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using
 $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.27±0.04 OUR AVERAGE					
$0.27 \pm 0.04 \pm 0.02$		2.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
$0.30 \pm 0.04 \pm 0.10$		8.7k	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.8	90	⁵⁴ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
---------	----	---------------------	----------	--

⁵⁴ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] <$
 21.9×10^{-2} which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$
 3.8×10^{-2} .

$\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.2				
	90	ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 19	90	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.1 ± 0.7					OUR AVERAGE
$7.1 \pm 1.8 \pm 1.3$		$2.3 \pm 0.5\text{k}$	⁵⁵ BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
$4.8 \pm 0.5 \pm 0.6$		$19 \pm 3\text{k}$	⁵⁵ AUBERT	09AQ	BABR $\Upsilon(3S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.5	90		LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
$4.8 \pm 0.5 \pm 1.2$		$19 \pm 3\text{k}$	^{55,56} AUBERT	08V	BABR $\Upsilon(3S) \rightarrow \gamma X$
< 4.3	90		⁵⁷ ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$

⁵⁵ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

⁵⁶ Systematic error re-evaluated by AUBERT 09AQ.

⁵⁷ Superseded by BONVICINI 10.

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ Γ_{28}/Γ ($0.3 \text{ GeV} < m_{A^0} < 7 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	⁵⁸ LEES	11H	BABR $\Upsilon(3S) \rightarrow \gamma \text{hadrons}$

⁵⁸ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{29}/Γ ($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.5	90	⁵⁹ AUBERT	09Z	BABR $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

⁵⁹ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from 0.27 – 5.5×10^{-6} .

$\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-4}$	90	⁶⁰ AUBERT	09P	BABR $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

⁶⁰ For a narrow scalar or pseudoscalar, A^0 , with $M(\tau^+ \tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from 1.5 – 16×10^{-5} .

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.2	90	LEES	10B	BABR $e^+e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					Γ_{33}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 3.6 \times 10^{-7}$	90	LEES	22A	BABR	$e^+ e^- \rightarrow e^\pm \mu^\mp$
$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{34}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
< 3.1	90	LEES	10B	BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 20.3	95	LOVE	08A	CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

$\tau(3S)$ REFERENCES

AAIJ	24AC	JHEP 2410 122	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
LEES	22A	PRL 128 091804	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	20E	PRL 125 241801	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+ e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86B	ZPHY C32 622 (errat.)	S.E. Baru <i>et al.</i>	(NOVO)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)