



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$  Status: \* \* \* \*

Neither  $J$  or  $P$  has actually been measured.

Ξ\_c^0 MASS

The fit uses the Ξ\_c^0 and Ξ\_c^+ mass and mass-difference measurements.

| VALUE (MeV)                                                                   | EVTS | DOCUMENT ID           | TECN     | COMMENT                                 |
|-------------------------------------------------------------------------------|------|-----------------------|----------|-----------------------------------------|
| <b>2470.44±0.28 OUR FIT</b> Error includes scale factor of 1.2.               |      |                       |          |                                         |
| <b>2470.99<sup>+0.30</sup><sub>-0.50</sub> OUR AVERAGE</b>                    |      |                       |          |                                         |
| 2470.85±0.24±0.55                                                             | 3.4k | AALTONEN              | 14B CDF  | $p\bar{p}$ at 1.96 TeV                  |
| 2471.0 ±0.3 <sup>+0.2</sup> <sub>-1.4</sub>                                   | 8.6k | <sup>1</sup> LESIAK   | 05 BELL  | $e^+e^-$ , $\Upsilon(4S)$               |
| 2470.0 ±2.8 ±2.6                                                              | 85   | FRABETTI              | 98B E687 | $\gamma$ Be, $\bar{E}_\gamma = 220$ GeV |
| 2469 ±2 ±3                                                                    | 9    | HENDERSON             | 92B CLEO | $\Omega^- K^+$                          |
| 2472.1 ±2.7 ±1.6                                                              | 54   | ALBRECHT              | 90F ARG  | $e^+e^-$ at $\Upsilon(4S)$              |
| 2473.3 ±1.9 ±1.2                                                              | 4    | BARLAG                | 90 ACCM  | $\pi^- (K^-)$ Cu 230 GeV                |
| 2472 ±3 ±4                                                                    | 19   | ALAM                  | 89 CLEO  | $e^+e^-$ 10.6 GeV                       |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                       |          |                                         |
| 2462.1 ±3.1 ±1.4                                                              | 42   | <sup>2</sup> FRABETTI | 93C E687 | See FRABETTI 98B                        |
| 2471 ±3 ±4                                                                    | 14   | AVERY                 | 89 CLEO  | See ALAM 89                             |

<sup>1</sup> The systematic error was (wrongly) given the other way round in LESIAK 05.

<sup>2</sup> The FRABETTI 93C mass is well below the other measurements.

Ξ\_c^0 - Ξ\_c^+ MASS DIFFERENCE

| VALUE (MeV)                                                  | EVTS     | DOCUMENT ID | TECN    | COMMENT                                                                     |
|--------------------------------------------------------------|----------|-------------|---------|-----------------------------------------------------------------------------|
| <b>2.72±0.23 OUR FIT</b> Error includes scale factor of 1.1. |          |             |         |                                                                             |
| <b>2.91±0.26 OUR AVERAGE</b>                                 |          |             |         |                                                                             |
| 2.85±0.30±0.04                                               | 5.1/3.4k | AALTONEN    | 14B CDF | $p\bar{p}$ at 1.96 TeV                                                      |
| 2.9 ±0.5                                                     |          | LESIAK      | 05 BELL | $e^+e^-$ , $\Upsilon(4S)$                                                   |
| 7.0 ±4.5 ±2.2                                                |          | ALBRECHT    | 90F ARG | $e^+e^-$ at $\Upsilon(4S)$                                                  |
| 6.8 ±3.3 ±0.5                                                |          | BARLAG      | 90 ACCM | $\pi^- (K^-)$ Cu 230 GeV                                                    |
| 5 ±4 ±1                                                      |          | ALAM        | 89 CLEO | $\Xi_c^0 \rightarrow \Xi^- \pi^+$ , $\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$ |

Ξ\_c^0 MEAN LIFE

| VALUE (10 <sup>-15</sup> s)                                       | EVTS | DOCUMENT ID         | TECN      | COMMENT                                                                                       |
|-------------------------------------------------------------------|------|---------------------|-----------|-----------------------------------------------------------------------------------------------|
| <b>150.4± 2.8 OUR AVERAGE</b> Error includes scale factor of 1.4. |      |                     |           |                                                                                               |
| 148.0± 2.3±2.2                                                    |      | <sup>1</sup> AAIJ   | 22Y LHCb  | $pp \rightarrow \Xi_c^0 + X$ , $\Xi_c^0 \rightarrow p K^- K^- \pi^+$                          |
| 153.4± 2.4±0.7                                                    | 22k  | <sup>2,3</sup> AAIJ | 19AG LHCb | $\Xi_b^- \rightarrow \Xi_c^0 \mu^- \bar{\nu}_\mu + X$ , $\Xi_c^0 \rightarrow p K^- K^- \pi^+$ |

|     |                |         |     |          |          |                                              |
|-----|----------------|---------|-----|----------|----------|----------------------------------------------|
| 118 | $+14$<br>$-12$ | $\pm 5$ | 110 | LINK     | 02H FOCS | $\gamma$ nucleus, $\approx 180$ GeV          |
| 101 | $+25$<br>$-17$ | $\pm 5$ | 42  | FRABETTI | 93C E687 | $\gamma$ Be, $\overline{E}_\gamma = 220$ GeV |
| 82  | $+59$<br>$-30$ |         | 4   | BARLAG   | 90 ACCM  | $\pi^- (K^-)$ Cu 230 GeV                     |

<sup>1</sup> Measured in  $\Xi_c^0$  produced promptly in  $pp$  collisions, using  $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$  as normalisation mode. AAIJ 22Y reports this lifetime value as  $(148.0 \pm 2.3 \pm 2.2 \pm 0.2) \times 10^{-15}$  s where the last uncertainty is due to the uncertainty on the  $D^0$  lifetime value from PDG 20 average,  $\tau_{D^0} = (410.1 \pm 1.5)$  fs.

<sup>2</sup> AAIJ 19AG reports  $[\Xi_c^0 \text{ MEAN LIFE}] / [D^\pm \text{ MEAN LIFE}] = 0.1485 \pm 0.0017 \pm 0.0016$  which we multiply by our best value  $D^\pm \text{ MEAN LIFE} = (1.033 \pm 0.005) \times 10^{-12}$  s. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Measured in  $\Xi_c^0$  produced in semileptonic  $\Xi_b^-$  decays.

## $\Xi_c^0$ DECAY MODES

| Mode                                                                              | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level |
|-----------------------------------------------------------------------------------|--------------------------------|------------------|
| <b>Cabibbo-favored decays</b>                                                     |                                |                  |
| $\Gamma_1$ $p K^- K^- \pi^+$                                                      | $(4.9 \pm 1.0) \times 10^{-3}$ |                  |
| $\Gamma_2$ $p K^- \overline{K}^*(892)^0, \overline{K}^{*0} \rightarrow K^- \pi^+$ | $(2.0 \pm 0.6) \times 10^{-3}$ |                  |
| $\Gamma_3$ $p K^- K^- \pi^+$ (no $\overline{K}^{*0}$ )                            | $(3.0 \pm 0.8) \times 10^{-3}$ |                  |
| $\Gamma_4$ $\Lambda K_S^0$                                                        | $(3.2 \pm 0.6) \times 10^{-3}$ |                  |
| $\Gamma_5$ $\Lambda K^- \pi^+$                                                    | $(1.45 \pm 0.28) \%$           |                  |
| $\Gamma_6$ $\Lambda \overline{K}^*(892)^0$                                        | $(2.6 \pm 0.6) \times 10^{-3}$ |                  |
| $\Gamma_7$ $\Lambda \overline{K}^0 \pi^+ \pi^-$                                   | seen                           |                  |
| $\Gamma_8$ $\Lambda K^- \pi^+ \pi^+ \pi^-$                                        | seen                           |                  |
| $\Gamma_9$ $\Sigma^0 K_S^0$                                                       | $(5.4 \pm 1.4) \times 10^{-4}$ |                  |
| $\Gamma_{10}$ $\Sigma^+ K^-$                                                      | $(1.8 \pm 0.4) \times 10^{-3}$ |                  |
| $\Gamma_{11}$ $\Sigma^0 \overline{K}^*(892)^0$                                    | $(9.9 \pm 1.9) \times 10^{-3}$ |                  |
| $\Gamma_{12}$ $\Sigma^+ K^*(892)^-$                                               | $(4.9 \pm 1.3) \times 10^{-3}$ |                  |
| $\Gamma_{13}$ $\Xi^- \pi^+$                                                       | $(1.43 \pm 0.27) \%$           |                  |
| $\Gamma_{14}$ $\Xi^- \pi^+ \pi^+ \pi^-$                                           | $(4.8 \pm 2.3) \%$             |                  |
| $\Gamma_{15}$ $\Xi^0 \pi^0$                                                       | $(6.9 \pm 1.4) \times 10^{-3}$ |                  |
| $\Gamma_{16}$ $\Xi^0 \eta$                                                        | $(1.6 \pm 0.4) \times 10^{-3}$ |                  |
| $\Gamma_{17}$ $\Xi^0 \eta'$                                                       | $(1.1 \pm 0.4) \times 10^{-3}$ |                  |
| $\Gamma_{18}$ $\Xi^0 K^+ K^-$                                                     |                                |                  |
| $\Gamma_{19}$ $\Xi^0 \phi, \phi \rightarrow K^+ K^-$                              | $(5.2 \pm 1.2) \times 10^{-4}$ |                  |
| $\Gamma_{20}$ $\Xi^0 K^+ K^-$ nonresonant                                         | $(5.6 \pm 1.2) \times 10^{-4}$ |                  |
| $\Gamma_{21}$ $\Omega^- K^+$                                                      | $(4.2 \pm 0.9) \times 10^{-3}$ |                  |
| $\Gamma_{22}$ $\Xi^- e^+ \nu_e$                                                   | $(1.05 \pm 0.20) \%$           |                  |
| $\Gamma_{23}$ $\Xi^- \mu^+ \nu_\mu$                                               | $(1.01 \pm 0.21) \%$           |                  |
| $\Gamma_{24}$ $\Xi^0 \gamma$                                                      | $< 1.7 \times 10^{-4}$         | 90%              |
| $\Gamma_{25}$ $\Xi^0 \mu^+ \mu^-$                                                 | $< 6 \times 10^{-5}$           | 90%              |
| $\Gamma_{26}$ $\Xi^0 e^+ e^-$                                                     | $< 1.0 \times 10^{-4}$         | 90%              |

**Cabibbo-suppressed decays**

|               |                                |                                |
|---------------|--------------------------------|--------------------------------|
| $\Gamma_{27}$ | $\Lambda_c^+ \pi^-$            | $(5.5 \pm 1.1) \times 10^{-3}$ |
| $\Gamma_{28}$ | $\Xi^- K^+$                    | $(3.9 \pm 1.1) \times 10^{-4}$ |
| $\Gamma_{29}$ | $\Lambda K^+ K^-$ (no $\phi$ ) | $(4.1 \pm 1.3) \times 10^{-4}$ |
| $\Gamma_{30}$ | $\Lambda \phi$                 | $(4.9 \pm 1.3) \times 10^{-4}$ |

**FIT INFORMATION**

An overall fit to 7 branching ratios uses 8 measurements to determine 4 parameters. The overall fit has a  $\chi^2 = 1.4$  for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

|          |       |       |          |
|----------|-------|-------|----------|
| $x_5$    | 64    |       |          |
| $x_{13}$ | 86    | 74    |          |
| $x_{27}$ | 64    | 55    | 75       |
|          | $x_1$ | $x_5$ | $x_{13}$ |

 **$\Xi_c^0$  BRANCHING RATIOS****Cabibbo-favored ( $S = -2$ ) decays**

| $\Gamma(p K^- K^- \pi^+) / \Gamma_{\text{total}}$ |            | $\Gamma_1 / \Gamma$ |      |         |                             |
|---------------------------------------------------|------------|---------------------|------|---------|-----------------------------|
| VALUE (%)                                         | EVTS       | DOCUMENT ID         | TECN | COMMENT |                             |
| <b><math>0.49 \pm 0.10</math> OUR FIT</b>         |            |                     |      |         |                             |
| <b><math>0.58 \pm 0.23 \pm 0.05</math></b>        | $17 \pm 5$ | LI                  | 19A  | BELL    | $e^+ e^-$ at $\Upsilon(4S)$ |

| $\Gamma(p K^- K^- \pi^+) / \Gamma(\Xi^- \pi^+)$ |               | $\Gamma_1 / \Gamma_{13}$ |      |         |                            |
|-------------------------------------------------|---------------|--------------------------|------|---------|----------------------------|
| VALUE                                           | EVTS          | DOCUMENT ID              | TECN | COMMENT |                            |
| <b><math>0.339 \pm 0.035</math> OUR FIT</b>     |               |                          |      |         |                            |
| <b><math>0.34 \pm 0.04</math> OUR AVERAGE</b>   |               |                          |      |         |                            |
| $0.33 \pm 0.03 \pm 0.03$                        | $1908 \pm 62$ | LESLIAK                  | 05   | BELL    | $e^+ e^-$ , $\Upsilon(4S)$ |
| $0.35 \pm 0.06 \pm 0.03$                        | $148 \pm 18$  | DANKO                    | 04   | CLEO    | $e^+ e^-$                  |

| $\Gamma(p K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+) / \Gamma(\Xi^- \pi^+)$ |  | $\Gamma_2 / \Gamma_{13}$ |      |         |           |
|--------------------------------------------------------------------------------------------|--|--------------------------|------|---------|-----------|
| VALUE                                                                                      |  | DOCUMENT ID              | TECN | COMMENT |           |
| <b><math>0.14 \pm 0.03 \pm 0.01</math></b>                                                 |  | DANKO                    | 04   | CLEO    | $e^+ e^-$ |

| $\Gamma(p K^- K^- \pi^+ (\text{no } \bar{K}^{*0})) / \Gamma(\Xi^- \pi^+)$ |  | $\Gamma_3 / \Gamma_{13}$ |      |         |           |
|---------------------------------------------------------------------------|--|--------------------------|------|---------|-----------|
| VALUE                                                                     |  | DOCUMENT ID              | TECN | COMMENT |           |
| <b><math>0.21 \pm 0.04 \pm 0.02</math></b>                                |  | DANKO                    | 04   | CLEO    | $e^+ e^-$ |

| $\Gamma(\Lambda K_S^0) / \Gamma(\Xi^- \pi^+)$   |              | $\Gamma_4 / \Gamma_{13}$ |      |         |                             |
|-------------------------------------------------|--------------|--------------------------|------|---------|-----------------------------|
| VALUE                                           | EVTS         | DOCUMENT ID              | TECN | COMMENT |                             |
| <b><math>0.225 \pm 0.013</math> OUR AVERAGE</b> |              |                          |      |         |                             |
| $0.229 \pm 0.008 \pm 0.012$                     | 5.6k         | LI                       | 21F  | BELL    | $e^+ e^-$ at $\Upsilon(nS)$ |
| $0.21 \pm 0.02 \pm 0.02$                        | $465 \pm 37$ | LESLIAK                  | 05   | BELL    | $e^+ e^-$ , $\Upsilon(4S)$  |

| $\Gamma(\Lambda K^- \pi^+)/\Gamma_{\text{total}}$             |                                     |             |      |         | $\Gamma_5/\Gamma$                       |
|---------------------------------------------------------------|-------------------------------------|-------------|------|---------|-----------------------------------------|
| VALUE (%)                                                     | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>1.45±0.28 OUR FIT</b>                                      |                                     |             |      |         |                                         |
| <b>1.17±0.37±0.09</b>                                         | 24 ± 6                              | LI          | 19A  | BELL    | $e^+e^-$ at $\Upsilon(4S)$              |
| $\Gamma(\Lambda K^- \pi^+)/\Gamma(\Xi^- \pi^+)$               |                                     |             |      |         | $\Gamma_5/\Gamma_{13}$                  |
| VALUE                                                         | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>1.02±0.14 OUR FIT</b>                                      | Error includes scale factor of 1.1. |             |      |         |                                         |
| <b>1.07±0.12±0.07</b>                                         | 2979 ± 211                          | LESIKAK     | 05   | BELL    | $e^+e^-$ , $\Upsilon(4S)$               |
| $\Gamma(\Lambda \bar{K}^*(892)^0)/\Gamma(\Xi^- \pi^+)$        |                                     |             |      |         | $\Gamma_6/\Gamma_{13}$                  |
| VALUE                                                         | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>0.18±0.02±0.01</b>                                         | 4k                                  | JIA         | 21   | BELL    | $e^+e^-$ at $\Upsilon(nS)$              |
| $\Gamma(\Lambda \bar{K}^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ |                                     |             |      |         | $\Gamma_7/\Gamma$                       |
| VALUE                                                         |                                     | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>seen</b>                                                   |                                     | FRABETTI    | 98B  | E687    | $\gamma$ Be, $\bar{E}_\gamma = 220$ GeV |
| $\Gamma(\Lambda K^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ |                                     |             |      |         | $\Gamma_8/\Gamma$                       |
| VALUE                                                         |                                     | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>seen</b>                                                   |                                     | FRABETTI    | 98B  | E687    | $\gamma$ Be, $\bar{E}_\gamma = 220$ GeV |
| $\Gamma(\Sigma^0 K_S^0)/\Gamma(\Xi^- \pi^+)$                  |                                     |             |      |         | $\Gamma_9/\Gamma_{13}$                  |
| VALUE (units $10^{-2}$ )                                      | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>3.8±0.6±0.4</b>                                            | 279                                 | LI          | 21F  | BELL    | $e^+e^-$ at $\Upsilon(nS)$              |
| $\Gamma(\Sigma^+ K^-)/\Gamma(\Xi^- \pi^+)$                    |                                     |             |      |         | $\Gamma_{10}/\Gamma_{13}$               |
| VALUE (units $10^{-2}$ )                                      | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>12.3±0.7±1.0</b>                                           | 889                                 | LI          | 21F  | BELL    | $e^+e^-$ at $\Upsilon(nS)$              |
| $\Gamma(\Sigma^0 \bar{K}^*(892)^0)/\Gamma(\Xi^- \pi^+)$       |                                     |             |      |         | $\Gamma_{11}/\Gamma_{13}$               |
| VALUE                                                         | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>0.69±0.03±0.03</b>                                         | 6.3k                                | JIA         | 21   | BELL    | $e^+e^-$ at $\Upsilon(nS)$              |
| $\Gamma(\Sigma^+ K^*(892)^-)/\Gamma(\Xi^- \pi^+)$             |                                     |             |      |         | $\Gamma_{12}/\Gamma_{13}$               |
| VALUE                                                         | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>0.34±0.06±0.02</b>                                         | 373                                 | JIA         | 21   | BELL    | $e^+e^-$ at $\Upsilon(nS)$              |
| $\Gamma(\Xi^- \pi^+)/\Gamma_{\text{total}}$                   |                                     |             |      |         | $\Gamma_{13}/\Gamma$                    |
| VALUE (%)                                                     | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>1.43±0.27 OUR FIT</b>                                      |                                     |             |      |         |                                         |
| <b>1.80±0.50±0.14</b>                                         | 45 ± 7                              | LI          | 19A  | BELL    | $e^+e^-$ at $\Upsilon(4S)$              |
| $\Gamma(\Xi^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+ \pi^-)$         |                                     |             |      |         | $\Gamma_{13}/\Gamma_{14}$               |
| VALUE                                                         |                                     | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>0.30±0.12±0.05</b>                                         |                                     | ALBRECHT    | 90F  | ARG     | $e^+e^-$ at $\Upsilon(4S)$              |
| $\Gamma(\Omega^- K^+)/\Gamma(\Xi^- \pi^+)$                    |                                     |             |      |         | $\Gamma_{21}/\Gamma_{13}$               |
| VALUE                                                         | EVTS                                | DOCUMENT ID | TECN | COMMENT |                                         |
| <b>0.294±0.018±0.016</b>                                      | 650                                 | AUBERT,B    | 05M  | BABR    | $e^+e^- \approx \Upsilon(4S)$           |

$$\Gamma(\Xi^0 \phi, \phi \rightarrow K^+ K^-) / \Gamma(\Xi^- \pi^+) \quad \Gamma_{19} / \Gamma_{13}$$

| VALUE                                         | EVTs | DOCUMENT ID         | TECN | COMMENT                          |
|-----------------------------------------------|------|---------------------|------|----------------------------------|
| <b><math>0.036 \pm 0.004 \pm 0.002</math></b> | 311  | <sup>1</sup> MCNEIL | 21   | BELL $e^+ e^-$ at $\Upsilon(nS)$ |

<sup>1</sup> MCNEIL 21 assumes an azimuthally symmetric amplitude model to recover resonant and nonresonant contributions to  $\Xi_c^0 \rightarrow \Xi_c^0 K^+ K^-$ .

$$\Gamma(\Xi^0 \pi^0) / \Gamma(\Xi^- \pi^+) \quad \Gamma_{15} / \Gamma_{13}$$

| VALUE                                      | EVTs | DOCUMENT ID         | TECN | COMMENT                               |
|--------------------------------------------|------|---------------------|------|---------------------------------------|
| <b><math>0.48 \pm 0.02 \pm 0.03</math></b> | 2.2k | <sup>1</sup> ADACHI | 24s  | BEL2 $e^+ e^-$ at $\sim \Upsilon(nS)$ |

<sup>1</sup> Analysis of Belle and Belle II data samples.

$$\Gamma(\Xi^0 \eta) / \Gamma(\Xi^- \pi^+) \quad \Gamma_{16} / \Gamma_{13}$$

| VALUE                                      | EVTs  | DOCUMENT ID         | TECN | COMMENT                               |
|--------------------------------------------|-------|---------------------|------|---------------------------------------|
| <b><math>0.11 \pm 0.01 \pm 0.01</math></b> | 0.14k | <sup>1</sup> ADACHI | 24s  | BEL2 $e^+ e^-$ at $\sim \Upsilon(nS)$ |

<sup>1</sup> Analysis of Belle and Belle II data samples.

$$\Gamma(\Xi^0 \eta') / \Gamma(\Xi^- \pi^+) \quad \Gamma_{17} / \Gamma_{13}$$

| VALUE                                      | EVTs | DOCUMENT ID         | TECN | COMMENT                               |
|--------------------------------------------|------|---------------------|------|---------------------------------------|
| <b><math>0.08 \pm 0.02 \pm 0.01</math></b> | 31   | <sup>1</sup> ADACHI | 24s  | BEL2 $e^+ e^-$ at $\sim \Upsilon(nS)$ |

<sup>1</sup> Analysis of Belle and Belle II data samples.

$$\Gamma(\Xi^0 K^+ K^- \text{ nonresonant}) / \Gamma(\Xi^- \pi^+) \quad \Gamma_{20} / \Gamma_{13}$$

| VALUE                                         | EVTs | DOCUMENT ID         | TECN | COMMENT                          |
|-----------------------------------------------|------|---------------------|------|----------------------------------|
| <b><math>0.039 \pm 0.004 \pm 0.002</math></b> | 311  | <sup>1</sup> MCNEIL | 21   | BELL $e^+ e^-$ at $\Upsilon(nS)$ |

<sup>1</sup> MCNEIL 21 assumes an azimuthally symmetric amplitude model to recover resonant and nonresonant contributions to  $\Xi_c^0 \rightarrow \Xi_c^0 K^+ K^-$ .

$$\Gamma(\Xi^- e^+ \nu_e) / \Gamma(\Xi^- \pi^+) \quad \Gamma_{22} / \Gamma_{13}$$

| VALUE                                         | EVTs | DOCUMENT ID     | TECN | COMMENT                            |
|-----------------------------------------------|------|-----------------|------|------------------------------------|
| <b><math>0.730 \pm 0.021 \pm 0.039</math></b> |      | <sup>1</sup> LI | 21c  | BELL $e^+ e^-$ at 10.52, 10.58 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.38  $\pm 0.14 \pm 0.22$  ACHARYA 21A ALCE  $pp$  at 13 TeV

3.1  $\pm 1.0 \begin{smallmatrix} +0.3 \\ -0.5 \end{smallmatrix}$  54 ALEXANDER 95B CLE2  $e^+ e^- \approx \Upsilon(4S)$

0.96  $\pm 0.43 \pm 0.18$  18 <sup>2</sup> ALBRECHT 93B ARG  $e^+ e^- \approx 10.4$  GeV

<sup>1</sup> LI 21c measures ratio  $B(\Xi_c^0 \rightarrow \Xi^- e^+ \nu_e) / B(\Xi_c^0 \rightarrow \Xi^- \mu^+ \nu_\mu) = 1.03 \pm 0.05 \pm 0.07$ .

<sup>2</sup> This ALBRECHT 93B value is the average of the  $(\Xi^- e^+ \text{ anything}) / \Xi^- \pi^+$  and  $(\Xi^- \mu^+ \text{ anything}) / \Xi^- \pi^+$  ratios. Here we average it with the  $\Xi^- e^+ \nu_e / \Xi^- \pi^+$  ratio.

$$\Gamma(\Xi^- e^+ \nu_e) / \Gamma(\Xi^- \mu^+ \nu_\mu) \quad \Gamma_{22} / \Gamma_{23}$$

| VALUE                                                                         | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|-------------|------|---------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |         |

1.03  $\pm 0.05 \pm 0.07$  <sup>1</sup> LI 21c BELL  $e^+ e^-$  at 10.52, 10.58 GeV

<sup>1</sup> LI 21c value is not independent from other quoted measurements.

$$\Gamma(\Xi^- \mu^+ \nu_\mu) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{23}/\Gamma_{13}$ 

| VALUE                                                                                                                                               | DOCUMENT ID     | TECN     | COMMENT                       |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|----------|-------------------------------|
| <b>0.708 ± 0.033 ± 0.056</b>                                                                                                                        | <sup>1</sup> LI | 21C BELL | $e^+ e^-$ at 10.52, 10.58 GeV |
| <sup>1</sup> LI 21C measures ratio $B(\Xi_c^0 \rightarrow \Xi^- e^+ \nu_e) / B(\Xi_c^0 \rightarrow \Xi^- \mu^+ \nu_\mu) = 1.03 \pm 0.05 \pm 0.07$ . |                 |          |                               |

$$\Gamma(\Xi^0 \gamma) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{24}/\Gamma_{13}$ 

| VALUE                             | CL% | DOCUMENT ID | TECN    | COMMENT                          |
|-----------------------------------|-----|-------------|---------|----------------------------------|
| <b>&lt; 1.2 × 10<sup>-2</sup></b> | 90  | LI          | 23 BELL | $e^+ e^- \rightarrow \gamma(nS)$ |

$$\Gamma(\Xi^0 \mu^+ \mu^-) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{25}/\Gamma_{13}$ 

| VALUE                             | CL% | DOCUMENT ID | TECN    | COMMENT                                  |
|-----------------------------------|-----|-------------|---------|------------------------------------------|
| <b>&lt; 4.3 × 10<sup>-3</sup></b> | 90  | CUI         | 24 BELL | 980fb <sup>-1</sup> , $e^+ e^-$ at Y(4S) |

$$\Gamma(\Xi^0 e^+ e^-) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{26}/\Gamma_{13}$ 

| VALUE                             | CL% | DOCUMENT ID | TECN    | COMMENT                                  |
|-----------------------------------|-----|-------------|---------|------------------------------------------|
| <b>&lt; 6.7 × 10<sup>-3</sup></b> | 90  | CUI         | 24 BELL | 980fb <sup>-1</sup> , $e^+ e^-$ at Y(4S) |

### Cabibbo-suppressed decays

$$\Gamma(\Lambda_c^+ \pi^-) / \Gamma_{\text{total}}$$

 $\Gamma_{27}/\Gamma$ 

| VALUE (units 10 <sup>-3</sup> ) | EVTS | DOCUMENT ID       | TECN      | COMMENT        |
|---------------------------------|------|-------------------|-----------|----------------|
| <b>5.5 ± 1.1 OUR FIT</b>        |      |                   |           |                |
| <b>5.5 ± 0.2 ± 1.8</b>          | 6.3k | <sup>1</sup> AAIJ | 20AH LHCB | $pp$ at 13 TeV |

<sup>1</sup> AAIJ 20AH extracts  $B(\Xi_c^0 \rightarrow \Lambda_c^+ \pi^-)$  using two different normalization modes:  $\Lambda_c^+ \rightarrow p K^- \pi^+$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$ . The mean value of both results, taking their correlations into account, is presented as the final result. The measurement assumes production fraction ratios  $f_{\Xi_c^0}/f_{\Lambda_c^+} = (9.7 \pm 0.9 \pm 3.1) \times 10^{-2}$  (from AAIJ 19AB plus heavy quark symmetry arguments) as well as  $f_{\Xi_c^0}/f_{\Xi_c^+} = 1.00 \pm 0.01$ . It further uses the inputs  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.23 \pm 0.33) \times 10^{-2}$  and  $B(\Xi_c^+ \rightarrow p K^- \pi^+) = (4.5 \pm 2.1 \pm 0.7) \times 10^{-3}$  (from LI 19C). Its correlation with  $B(\Xi_c^+ \rightarrow p K^- \pi^+)$ , as measured in AAIJ 20AH, is 0.414.

$$\Gamma(\Lambda_c^+ \pi^-) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{27}/\Gamma_{13}$ 

| VALUE                                                                                                                         | EVTS | DOCUMENT ID       | TECN    | COMMENT                          |
|-------------------------------------------------------------------------------------------------------------------------------|------|-------------------|---------|----------------------------------|
| <b>0.38 ± 0.05 OUR FIT</b>                                                                                                    |      |                   |         |                                  |
| <b>0.38 ± 0.04 ± 0.04</b>                                                                                                     | 1468 | <sup>1</sup> TANG | 23 BELL | $e^+ e^- \rightarrow \gamma(nS)$ |
| <sup>1</sup> TANG 23 reports fitted masses $m_{\Lambda_c^+} = 2286.55 \pm 0.03$ MeV and $m_{\Xi_c^0} = 2470.43 \pm 0.06$ MeV. |      |                   |         |                                  |

$$\Gamma(\Xi^- K^+) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{28}/\Gamma_{13}$ 

| VALUE (units 10 <sup>-2</sup> ) | EVTS     | DOCUMENT ID | TECN    | COMMENT                      |
|---------------------------------|----------|-------------|---------|------------------------------|
| <b>2.75 ± 0.51 ± 0.25</b>       | 314 ± 58 | CHISTOV     | 13 BELL | $e^+ e^- \approx \gamma(4S)$ |

$$\Gamma(\Lambda K^+ K^- (\text{no } \phi)) / \Gamma(\Xi^- \pi^+)$$

 $\Gamma_{29}/\Gamma_{13}$ 

| VALUE (units 10 <sup>-2</sup> ) | EVTS      | DOCUMENT ID | TECN    | COMMENT                      |
|---------------------------------|-----------|-------------|---------|------------------------------|
| <b>2.86 ± 0.61 ± 0.37</b>       | 510 ± 110 | CHISTOV     | 13 BELL | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Lambda\phi)/\Gamma(\Xi^-\pi^+)$  $\Gamma_{30}/\Gamma_{13}$ 

| VALUE (units $10^{-2}$ )                   | EVTS         | DOCUMENT ID | TECN | COMMENT                            |
|--------------------------------------------|--------------|-------------|------|------------------------------------|
| <b><math>3.43 \pm 0.58 \pm 0.32</math></b> | $316 \pm 54$ | CHISTOV     | 13   | BELL $e^+e^- \approx \Upsilon(4S)$ |

 $\Xi_c^0$  DECAY PARAMETERS

See the note on “Baryon Decay Parameters” in the neutron Listings.

 $\alpha$  FOR  $\Xi_c^0 \rightarrow \Xi^-\pi^+$ 

| VALUE                                       | EVTS | DOCUMENT ID | TECN | COMMENT                           |
|---------------------------------------------|------|-------------|------|-----------------------------------|
| <b><math>-0.64 \pm 0.05 \pm 0.01</math></b> |      | LI          | 21C  | BELL $e^+e^-$ at 10.52, 10.58 GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                  |     |      |    |                                    |
|----------------------------------|-----|------|----|------------------------------------|
| $-0.56 \pm 0.39^{+0.10}_{-0.09}$ | 138 | CHAN | 01 | CLE2 $e^+e^- \approx \Upsilon(4S)$ |
|----------------------------------|-----|------|----|------------------------------------|

 $\alpha$  FOR  $\Xi_c^0 \rightarrow \Xi^+\pi^-$ 

| VALUE                                      | DOCUMENT ID | TECN | COMMENT                           |
|--------------------------------------------|-------------|------|-----------------------------------|
| <b><math>0.61 \pm 0.05 \pm 0.01</math></b> | LI          | 21C  | BELL $e^+e^-$ at 10.52, 10.58 GeV |

 $\alpha$  FOR  $\Xi_c^0 \rightarrow \Lambda \bar{K}^*(892)^0$ 

| VALUE                                      | EVTS | DOCUMENT ID      | TECN | COMMENT                         |
|--------------------------------------------|------|------------------|------|---------------------------------|
| <b><math>0.15 \pm 0.22 \pm 0.04</math></b> | 4k   | <sup>1</sup> JIA | 21   | BELL $e^+e^-$ at $\Upsilon(nS)$ |

<sup>1</sup> JIA 21 measures  $\alpha(\Xi_c^0 \rightarrow \Lambda \bar{K}^*(892)^0)$   $\alpha(\Lambda \rightarrow p\pi^-) = 0.115 \pm 0.164 \pm 0.031$ , and uses  $\alpha(\Lambda \rightarrow p\pi^-) = 0.747 \pm 0.010$ .

 $\alpha$  FOR  $\Xi_c^0 \rightarrow \Sigma^+ K^*(892)^-$ 

| VALUE                                       | EVTS | DOCUMENT ID      | TECN | COMMENT                         |
|---------------------------------------------|------|------------------|------|---------------------------------|
| <b><math>-0.52 \pm 0.30 \pm 0.02</math></b> | 373  | <sup>1</sup> JIA | 21   | BELL $e^+e^-$ at $\Upsilon(nS)$ |

<sup>1</sup> JIA 21 measures  $\alpha(\Xi_c^0 \rightarrow \Sigma^+ \bar{K}^*(892)^-)$   $\alpha(\Sigma^+ \rightarrow p\pi^0) = 0.514 \pm 0.295 \pm 0.012$ , and uses  $\alpha(\Sigma^+ \rightarrow p\pi^0) = -0.980 \pm 0.017$ .

 $\alpha$  FOR  $\Xi_c^0 \rightarrow \Xi^0\pi^0$ 

| VALUE                                       | EVTS | DOCUMENT ID         | TECN | COMMENT                              |
|---------------------------------------------|------|---------------------|------|--------------------------------------|
| <b><math>-0.90 \pm 0.15 \pm 0.23</math></b> | 2.2k | <sup>1</sup> ADACHI | 24S  | BEL2 $e^+e^-$ at $\sim \Upsilon(nS)$ |

<sup>1</sup> Analysis of Belle and Belle II data samples.

 $\Xi_c^0$  Tests of Baryon Number Violation $\tau_{mix}$ ,  $\Xi_c^0 - \Xi_c^{\prime 0}$  oscillation period

| VALUE ( $10^{-12}$ s)       | DOCUMENT ID     | TECN | COMMENT                                |
|-----------------------------|-----------------|------|----------------------------------------|
| <b><math>&gt;1.3</math></b> | <sup>1</sup> GU | 24   | BELL $e^+e^- \rightarrow \Upsilon(4S)$ |

<sup>1</sup> Search for baryon-number violating decay  $B^- \rightarrow \Xi_c^0 \bar{\Lambda}_c^-$ , which can be interpreted as a search for  $B^- \rightarrow \Xi_c^0 \bar{\Lambda}_c^-$  followed by  $\Xi_c^0 - \Xi_c^{\prime 0}$  baryon-number violating oscillation, from which a bound on the oscillation period  $\tau_{mix}$  can be inferred, assuming no direct  $B^- \rightarrow \Xi_c^0 \bar{\Lambda}_c^-$  decay.

## REFERENCES

|           |      |                      |                             |                              |
|-----------|------|----------------------|-----------------------------|------------------------------|
| ADACHI    | 24S  | JHEP 2410 045        | I. Adachi <i>et al.</i>     | (BELLE and BELLE II Collab.) |
| CUI       | 24   | PR D109 052003       | J.X.Cui <i>et al.</i>       | (BELLE Collab.)              |
| GU        | 24   | PRL 133 071802       | T. Gu <i>et al.</i>         | (BELLE Collab.)              |
| LI        | 23   | PR D107 032001       | Y. Li <i>et al.</i>         | (BELLE Collab.)              |
| TANG      | 23   | PR D107 032005       | S.S. Tang <i>et al.</i>     | (BELLE Collab.)              |
| AAIJ      | 22Y  | SCIB 67 479          | R. Aaij <i>et al.</i>       | (LHCb Collab.)               |
| ACHARYA   | 21A  | PRL 127 272001       | S. Acharya <i>et al.</i>    | (ALICE Collab.)              |
| JIA       | 21   | JHEP 2106 160        | S. Jia <i>et al.</i>        | (BELLE Collab.)              |
| LI        | 21C  | PRL 127 121803       | Y.B. Li <i>et al.</i>       | (BELLE Collab.)              |
| LI        | 21F  | PR D105 L011102      | Y. Li <i>et al.</i>         | (BELLE Collab.)              |
| MCNEIL    | 21   | PR D103 112002       | J.T. McNeil <i>et al.</i>   | (BELLE Collab.)              |
| AAIJ      | 20AH | PR D102 071101       | R. Aaij <i>et al.</i>       | (LHCb Collab.)               |
| PDG       | 20   | PTEP 2020 083C01     | P.A. Zyla <i>et al.</i>     | (PDG Collab.)                |
| AAIJ      | 19AB | PR D99 052006        | R. Aaij <i>et al.</i>       | (LHCb Collab.)               |
| AAIJ      | 19AG | PR D100 032001       | R. Aaij <i>et al.</i>       | (LHCb Collab.)               |
| LI        | 19A  | PRL 122 082001       | Y.B. Li <i>et al.</i>       | (BELLE Collab.)              |
| LI        | 19C  | PR D100 031101       | Y.B. Li <i>et al.</i>       | (BELLE Collab.)              |
| AALTONEN  | 14B  | PR D89 072014        | T. Aaltonen <i>et al.</i>   | (CDF Collab.)                |
| CHISTOV   | 13   | PR D88 071103        | R. Chistov <i>et al.</i>    | (BELLE Collab.)              |
| AUBERT,B  | 05M  | PRL 95 142003        | B. Aubert <i>et al.</i>     | (BABAR Collab.)              |
| LESIAK    | 05   | PL B605 237          | T. Lesiak <i>et al.</i>     | (BELLE Collab.)              |
| Also      |      | PL B617 198 (errat.) | T. Lesiak <i>et al.</i>     | (BELLE Collab.)              |
| DANKO     | 04   | PR D69 052004        | I. Danko <i>et al.</i>      | (CLEO Collab.)               |
| LINK      | 02H  | PL B541 211          | J.M. Link <i>et al.</i>     | (FNAL FOCUS Collab.)         |
| CHAN      | 01   | PR D63 111102        | S. Chan <i>et al.</i>       | (CLEO Collab.)               |
| FRABETTI  | 98B  | PL B426 403          | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.)          |
| ALEXANDER | 95B  | PRL 74 3113          | J. Alexander <i>et al.</i>  | (CLEO Collab.)               |
| Also      |      | PRL 75 4155 (errat.) | J. Alexander <i>et al.</i>  | (CLEO Collab.)               |
| ALBRECHT  | 93B  | PL B303 368          | H. Albrecht <i>et al.</i>   | (ARGUS Collab.)              |
| FRABETTI  | 93C  | PRL 70 2058          | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.)          |
| HENDERSON | 92B  | PL B283 161          | S. Henderson <i>et al.</i>  | (CLEO Collab.)               |
| ALBRECHT  | 90F  | PL B247 121          | H. Albrecht <i>et al.</i>   | (ARGUS Collab.)              |
| BARLAG    | 90   | PL B236 495          | S. Barlag <i>et al.</i>     | (ACCMOR Collab.)             |
| ALAM      | 89   | PL B226 401          | M.S. Alam <i>et al.</i>     | (CLEO Collab.)               |
| AVERY     | 89   | PRL 62 863           | P. Avery <i>et al.</i>      | (CLEO Collab.)               |