

# QUARKS

The  $u$ -,  $d$ -, and  $s$ -quark masses are the  $\overline{MS}$  masses at the scale  $\mu = 2$  GeV. The  $c$ - and  $b$ -quark masses are the  $\overline{MS}$  masses renormalized at the  $\overline{MS}$  mass, i.e.  $\overline{m} = \overline{m}(\mu = \overline{m})$ . The  $t$ -quark mass is extracted from event kinematics (see the review “The Top Quark”).

 **$u$** 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_u = 2.16 \pm 0.07 \text{ MeV, CL} = 90\% \quad \text{Charge} = \frac{2}{3} e \quad I_z = +\frac{1}{2}$$

$$m_u/m_d = 0.462 \pm 0.020, \text{ CL} = 90\%$$

 **$d$** 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_d = 4.70 \pm 0.07 \text{ MeV, CL} = 90\% \quad \text{Charge} = -\frac{1}{3} e \quad I_z = -\frac{1}{2}$$

$$m_s/m_d = 17\text{--}22$$

$$\overline{m} = (m_u + m_d)/2 = 3.49 \pm 0.07 \text{ MeV, CL} = 90\%$$

 **$s$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_s = 93.5 \pm 0.8 \text{ MeV, CL} = 90\% \quad \text{Charge} = -\frac{1}{3} e \quad \text{Strangeness} = -1$$

$$m_s / ((m_u + m_d)/2) = 27.33^{+0.18}_{-0.14}, \text{ CL} = 90\%$$

 **$c$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_c = 1.2730 \pm 0.0046 \text{ GeV, CL} = 90\% \quad \text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

$$m_b - m_c = 3.45 \pm 0.05 \text{ GeV}$$

 **$b$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_b = 4.183 \pm 0.007 \text{ GeV, CL} = 90\% \quad \text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

 **$t$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Top} = +1$$

$$\text{Mass (direct measurements)} \quad m = 172.56 \pm 0.31 \text{ GeV}^{[a,b]} \quad (S = 1.6)$$

$$\text{Mass (from cross-section measurements)} \quad m = 162.5^{+2.1}_{-1.5} \text{ GeV}^{[a]}$$

$$\text{Mass (Pole from cross-section measurements)} \quad m = 172.4 \pm 0.7 \text{ GeV}$$

$$m_t - m_{\bar{t}} = -0.15 \pm 0.20 \text{ GeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma = 1.42_{-0.15}^{+0.19} \text{ GeV} \quad (S = 1.4)$$

$$\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.957 \pm 0.034 \quad (S = 1.5)$$

### **$t$ -quark EW Couplings**

$$F_0 = 0.693 \pm 0.013$$

$$F_- = 0.315 \pm 0.010$$

$$F_+ = -0.005 \pm 0.007$$

$$F_{V+A} < 0.29, \text{ CL} = 95\%$$

<b><math>t</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
$Wq(q = b, s, d)$			—
$Wb$			—
$e\nu_e b$	$(11.10 \pm 0.30) \%$		—
$\mu\nu_\mu b$	$(11.40 \pm 0.20) \%$		—
$\tau\nu_\tau b$	$(10.7 \pm 0.5) \%$		—
$q\bar{q}b$	$(66.5 \pm 1.4) \%$		—
$\gamma q(q=u, c)$	$[c] < 9.5$	$\times 10^{-6}$	95% —
$aq(q=u, c)$	$< 1$	$\times 10^{-3}$	95% —
<b><math>\Delta T = 1</math> weak neutral current (<math>T1</math>) modes</b>			
$Zq(q=u, c)$	$T1$ $[d] < 1.2$	$\times 10^{-4}$	95% —
$Hu$	$T1$ $< 1.9$	$\times 10^{-4}$	95% —
$Hc$	$T1$ $< 3.4$	$\times 10^{-4}$	95% —
$\ell^+ \bar{q} \bar{q}' (q=d, s, b; q'=u, c)$	$T1$ $< 1.6$	$\times 10^{-3}$	95% —
<b>Lepton Family number (<math>LF</math>) violating modes</b>			
$e^\pm \mu^\mp c$	$LF$ $< 8.9$	$\times 10^{-7}$	95% —
$e^\pm \mu^\mp u$	$LF$ $< 7$	$\times 10^{-8}$	95% —
$\mu^\pm \tau^\mp q$	$LF$ $< 8.7$	$\times 10^{-7}$	95% —

### **$b'$ (4<sup>th</sup> Generation) Quark, Searches for**

$$\text{Mass } m > 190 \text{ GeV, CL} = 95\% \quad (p\bar{p}, \text{ quasi-stable } b')$$

$$\text{Mass } m > 1390 \text{ GeV, CL} = 95\% \quad (\text{B}(b' \rightarrow Zb) = 1)$$

$$\text{Mass } m > 1350 \text{ GeV, CL} = 95\% \quad (\text{B}(b' \rightarrow Wt) = 1)$$

$$\text{Mass } m > 1570 \text{ GeV, CL} = 95\% \quad (\text{B}(b' \rightarrow Hb) = 1)$$

$$\text{Mass } m > 46.0 \text{ GeV, CL} = 95\% \quad (e^+ e^-, \text{ all decays})$$

## **$t'$ (4<sup>th</sup> Generation) Quark, Searches for**

$$\begin{aligned}
 m(t'(2/3)) &> 1280 \text{ GeV, CL} = 95\% & (\text{B}(t' \rightarrow Z t) = 1) \\
 m(t'(2/3)) &> 1295 \text{ GeV, CL} = 95\% & (\text{B}(t' \rightarrow W b) = 1) \\
 m(t'(2/3)) &> 1310 \text{ GeV, CL} = 95\% & (\text{singlet } t') \\
 m(t'(2/3)) &> 1350 \text{ GeV, CL} = 95\% & (t' \text{ in a weak isospin doublet } (t', b')) \\
 m(t'(5/3)) &> 1.460 \times 10^3 \text{ GeV, CL} = 95\% & (t'(5/3) \rightarrow t W^+)
 \end{aligned}$$

## **Free Quark Searches**

All searches since 1977 have had negative results.

## NOTES

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review “The Top Quark.”
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at  $\sqrt{s} = 7$  TeV. Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of  $173.2 \pm 0.9$  GeV. See the note “The Top Quark” in the Quark Particle Listings of this *Review*.
- [c] This limit is for  $\Gamma(t \rightarrow \gamma q)/\Gamma(t \rightarrow W b)$ .
- [d] This limit is for  $\Gamma(t \rightarrow Z q)/\Gamma(t \rightarrow W b)$ .